

THE DIFFERENTIATION OF BOVINE STAPHYLOCOCCI  
BY  
BACTERIOPHAGE

by  
PING HANG FUNG, B.Sc.  
THE UNIVERSITY OF MANITOBA



A Thesis  
submitted to  
The Faculty of Graduate Studies and Research  
The University of Manitoba

In partial fulfilment  
of the requirements for the degree  
of  
Master of Science

1960

## ACKNOWLEDGMENTS

The author is pleased to express his thanks to the late Dr. T.M.B. Payne, Professor and Chairman of the Department of Microbiology, under whose direction this study was carried out.

Grateful acknowledgment is likewise expressed to the Provincial Laboratory of Animal Pathology under the direction of Dr. J. Isa, for sympathetic encouragement and advice.

Finally sincere appreciation is extended to Dr. Norman James, Professor Emeritus of Microbiology, for his advice and assistance in the preparation of the manuscript.

THE DIFFERENTIATION OF BOVINE STAPHYLOCOCCI BY  
BACTERIOPHAGE

By Ping Hang Fung

ABSTRACT

A preliminary study was carried out on the optimum time for incubating phages produced by staphylococci. This was done by comparing the growth of each of several strains of Staphylococcus with the growth of the same strain plus its homologous phage in trypticase soy broth at 37°C at 15-minute intervals for nine hours, with an additional final colorimetric reading at 22 hours. A titer of the concentration of phage particles in the lysed culture was made at one hour and thereafter at half-hour intervals until the reading on the colorimeter remained stationary for one hour. The optimum time proved to be different for different groups of phages. In one group the titer remained at a high level overnight; in a second the titer was lowered appreciably; and in a third group the titer was practically lost. On the basis of this finding the time of incubation in the studies reported hereafter was varied from 2 to 6 hours, the time in each case depending on the reaction with the individual phage.

Three hundred and twenty-five isolates were obtained from udders of cows in nine areas adjacent to Winnipeg. Of these 166 were typable by using the 21 phages in the basic set together with six others. Most of the typable isolates were susceptible to phage 81. The majority of these isolates were susceptible also to one or more of four other phages—42B, 47B, 80 and 82. One phage-type was predominant in each area, but different predominant phage-types occurred in different areas. The majority of the isolates could not be grouped on the basis of the basic set alone.

Seven of 45 isolates obtained from pooled (cans) milk in one of these areas were typable. Two of the types from the pooled milk were the same as two of the 18 types obtained from individual cows in that area. Two other types were different from the types isolated from the milk of individual cows. In a second area investigated none of the isolates from the pooled milk was typable.

One of the seven isolates from bulk (tank) milk obtained from the latter two areas was typable. This type was the same as the predominant type isolated from herds in the first of the two areas.

In trials with 30 bottles of milk staphylococci were not found.

## TABLE OF CONTENTS

	PAGE
INTRODUCTION	1
HISTORICAL	4
MATERIAL AND METHODS	14
1. Milk samples	14
2. Isolation of staphylococci	15
3. Sources of bacteriophages	15
4. Culturing the propagating strains of staphylococci	16
5. Broth method of propagating bacteriophage	17
6. Plate method of propagating bacteriophage	18
7. Optimum incubation time for phage production	20
8. Titration of phage-containing broth	21
9. Testing of bacteriophage	22
10. Typing of isolates	23
<u>EXPERIMENTAL</u>	26
1. Optimum time of incubation for phage production	26
2. Phage-types of staphylococci	31
3. Phage-types of staphylococci from seven areas	33
4. Phage-types of staphylococci from other two areas	37
5. Isolates from pooled and bulk milk samples	40
DISCUSSION	42
REFERENCES	46
APPENDIX	51

## INDEX OF TABLES

TABLE		PAGE
I	Phage-types of staphylococci from udder milk	32
II	Susceptibility to certain other phages by staphylococci from milk lysed by phage 81	34
III	Phage-types of staphylococci from udder milk--Tuxedo herds	38
IV	Phage-types of staphylococci from pooled milk	40

## INDEX OF FIGURES

FIGURE		PAGE
1	Spotting of isolates with phages	24
2	Lysis of a strain (3A) of <u>Staphylococci aureus</u> by its homologous phage in trypticase soy broth at 37°C measured with a Beckman colorimeter	27
3	Lysis of a strain (77) of <u>Staphylococci aureus</u> by its homologous phage in trypticase soy broth at 37°C measured with a Beckman colorimeter	28
4	Lysis of a strain (3C) of <u>Staphylococci aureus</u> by its homologous phage in trypticase soy broth at 37°C measured with a Beckman colorimeter	30

INTRODUCTION



## INTRODUCTION

Staphylococci are widespread. Their presence in air and in the immediate environment of man is common. Under normal conditions they may be present in the nose, the throat or on the skin of the human and certain domestic animals. Due to their high adaptability staphylococcal infections have become of increasing concern, particularly during the past decade. They can adjust themselves to new environments such as human tissue bathed in antimicrobials, and most pathogenic strains presently predominant are antibiotic-resistant (16).

Of the acute infections in man staphylococcal infections present the most urgent need for basic treatment, especially following operations (2). This need results from the fact that penicillin-resistant strains have steadily risen from 50 to 80% of the strains currently carried by selected groups of individuals within hospitals, and to a lesser extent in the population at large (16). Staphylococcal

infections are endemic in the general population, since outbreaks have been reported repeatedly (2,19).

Immunization against staphylococcal diseases has not proven promising. Further, the development of strains resistant to antibiotics and drugs, currently accepted as effective, limits optimism that a permanent effective antibiotic or synthetic drug will appear in the immediate future. Therefore, control remains of primary importance. A prime requisite in this control is the ability to differentiate pathogenic from non-pathogenic strains.

The heterogeneity of this group of microorganisms confounds both the medical man and the biologist. In the medical field, a narrow view widely held is based on the dogma of the fixity of bacterial species, allowing plasticity within the species as regards virulence but failing to recognize variation due to genetic instability. The biologist, caring little about disease, aims to fit a species into a phylogenetical scheme and is bedevilled by the overlapping of characters such as coagulase reaction, toxin production,

pigment formation and tolerance to high salt concentration. However, methods of differentiating strains on the basis of one or more of these characters have not proven completely satisfactory (22). Bacteriophage typing appears to be stable and consistent. It provides an arbitrary method for the segregation of strains. The staphylococci that cause an epidemic in one area always fall into one phage pattern (17,19).

This presentation deals with an attempt to establish types of staphylococci by the bacteriophage method. It involves the flora in milk aseptically drawn from the cow and in milk at various stages of handling. The finding at a stage of handling of a type not present in the aseptically drawn sample could provide a basis for measuring sanitation in the handling of milk, since the new type could represent contamination. Conceivably, as an alternate probability, the new type could represent a genetical variant.

HISTORICAL

### HISTORICAL

The first investigation involving results that seemed to suggest that strains of staphylococci could be divided into groups according to their susceptibility to a series of bacteriophages was reported by Epstein and Fejgin (6). These investigators noted that staphylococci from lesions and from other sources did not respond in the same way to the phages studied. Burnet and Lush (3) divided different staphylococcal phages into different groups according to lytic action on different staphylococci. Williams and Timmins (24) attempted to use this phenomenon to differentiate the strains of staphylococci causing acute osteomyelitis.

However, it was Fisk (7,8) who initiated the present practice of phage typing of staphylococci. This investigator demonstrated that lysogenic staphylococci are widespread. He used the cross culture method to obtain different phages from the lysogenic strains of staphylococci and found that the reaction of these phages on different strains of staphylococci is selective. This reaction of a

staphylococcal strain to a given phage is not readily altered by changes in environment. Soon this method of typing proved of value in two small epidemiological studies (9,11).

Except for the addition of new phages and certain modifications in technique, the current methods used in phage typing of staphylococci are based on those used by Wilson and Atkinson (25). These workers demonstrated that phages could be separated from cocci by filtration. The earlier method depended upon inhibition of the cocci by the quaternary ammonium disinfectant, zephiran (7). They isolated 18 phages and devised a method for routine testing of different types of staphylococci on the basis of phage pattern. They divided their typable staphylococci into 21 types.

A comprehensive review of the methods currently in use has been reported by Williams and Rippon (23).

The main use of phages from human staphylococci has been in tracing the sources of epidemics of infections in man, but their use has been extended to the typing of staphylococci of animal origin.

MacDonald (12) was the first to report the

typing of staphylococci of bovine origin by phages. This investigator, working with phages obtained from Wilson and Atkinson, reported that the majority of the coagulase positive strains isolated from 280 samples of accredited milk, and all the 34 strains from bovine mastitis, were fully susceptible to phage 42D.

Smith (20), noting the value of the phage-typing in the investigation of outbreaks of staphylococcal infections in human subjects, attempted to ascertain whether the method of Wilson and Atkinson (25) might be of value in the epidemiological study of staphylococcal infections in animals. This investigator obtained 21 phages from G. S. Wilson, and isolated 11 additional phages. He reported that 25 of these could be grouped, as follows:

3A Group: Phages 3A, 3B, 3C and 51.

6 Group: Phages 6, 7, 42B, 47 and 47C.

29 Group: Phages 29, 29A, 31, 42C, 44, 52 and 52A.

42D Group: Phages 42D, E174/16, 14/94, 1363/14,  
129/16, 30/16 and P42D/E193.

88 Group: Phages 88 and 88A.

He showed that 93.3% of 1016 isolates obtained from

milk were typable by this grouping, whereas the remaining 6.7% proved to be susceptible to the 32 test phages. Over 83% were susceptible to one or more of the 42D group, and 9.5% to one or more of the 29 group. He divided the typable staphylococci from bovinemilk into 25 types. Strains isolated from the milk of cows suffering from acute or chronic mastitis did not differ in phage type from the strains isolated from the milk of cows with apparently normal udders. One phage type was predominant in the strains of staphylococci examined from one herd but in other herds the predominant types were different. Different phage types in different quarters of the same cow were not unusual but strains isolated from one quarter were generally of the same type, with few exceptions.

With a larger number of phages, Smith (20) typed a greater percentage of strains of staphylococci than did MacDonald (12) who used only the phages obtained from Wilson and Atkinson. Further, the percentage of strains fully susceptible to 42D alone (now allocated to Group IV) was not as large as that reported by MacDonald (12).

Price et al (14) studied the phage types of



staphylococci isolated from cows in England. These investigators divided most<sup>of</sup> the 896 isolates from 276 quarters of 15 herds into 12 fairly characteristic patterns when phages at routine test dilutions were used. These were designated by the letters A to M, as follows;

Code Letter	Phage susceptibility pattern determined with phages at routine test dilution
A	42D, 7/42D/44/44A, and other combinations of these four phages
B	Numerous phages of groups I and III
C	Lysed by one or more phages of group I and also by 42B
D	7 or 7/52A+
E	3C/7/29/42B/52/52A/54 (and 7/52/54+)
F	7/42B/54+

Code Letter	Phage susceptibility pattern determined with phages at routine test dilution
H	47A
I	42C/42D/44/44A/54+
J	54+
K	6/47/53/54+
L	6/7/47/53/54+
M	53/54

Three hundred and thirty nine colonies fell into pattern C; 183 into A; and 181 into L. In heavily infected herds infections were due to more than one phage type. Two phage types obtained from one quarter were uncommon, though different types from different quarters of the same cow were encountered quite frequently. This finding was in agreement with that obtained by Smith (20).

Barnum and Fuller (1) used the bacteriophage typing of staphylococci initiated by Smith (20) to

differentiate pathogenic from non-pathogenic strains in milk, along with other characteristics of the strains, in studies on the epidemiology and the efficacy of treatment and control measures.

These investigators isolated phages from 42 lyso-genic cultures of staphylococci. Using these phages, they tested 648 isolates of staphylococci from milk, and found that most of the isolates belonged to one of five groups. One hundred and ninety five belonged to Group A; 179 to Group A<sub>1</sub>; 76 to Group B; 45 to Group B<sub>1</sub>; 17 to Group B<sub>2</sub>; 123 were not susceptible to any of the 42 phages; and 13 were unclassifiable by these phages. Further, this method of phage typing did not distinguish between strains isolated from healthy and diseased udders. This result confirmed the finding of Smith (20). However, it did afford definite evidence that cows served as carriers of strains that could cause infection in the mammary gland. One phage group of staphylococci was predominant in one herd and other groups in different herds. This is in agreement with the results obtained by other workers (14).

Edwards (4) found that most strains from 35 cows of one herd were lysed by phage 42D at routine test dilution, but also were lysed by stronger concentrations

of phages 54, 77 and 79. Later this investigator working with Rippon (5) reported similar findings, in this case stronger concentrations of a larger number of phages proving to be lytic. They divided the pathogenic staphylococci isolated from these herds into five types, as follows:

Type	Diluted Phage	Phage Pattern undiluted
I	42D	29/52/79/42D/54/77
1a	42D	42D/6/7/42E/47
2	—	3A/3B/3C/71
3	52B	7/75/77/52B
4	52A	52/52A
5	78	54/76/77/78

All the above types were found in cases of clinical mastitis in cows. This finding confirmed the results of Price et al (14) mentioned earlier. These results likewise confirmed those of Smith (20) and Price et al (14) that one type was predominant in each herd but the predominant type in one herd differed from that in

another.

In a study of 57 isolates of staphylococci from milk of a mastitis-affected herd, Rountree et al (17) found that 33 were lysed by phage 42D or by other phages not attacking human strains. Twenty two isolates belonged to phage pattern 29/42D/52.

Thatcher and Simon (21) reported that the predominant type of staphylococci isolated from dairy products in Canada was 42D.

In Wisconsin, Seto and Wilson (18) isolated six staphylococcal phages designated as S1, S2, S3, S4, S5, and S6. The phages were propagated on micrococci of bovine origin. These investigators found that of 379 coagulase positive staphylococcal cultures tested 93% were sensitive to one or more of the above six phages and to phage 42D. All cultures of phage type S2 were sensitive to phage 44A-- this type was reported by McLean (13) to be the predominant type in staphylococci isolated from milk in Australia-- and 80% of the 44A phage type cultures were sensitive to phage S2. Thus, the predominant type of staphylococci of bovine origin isolated in Wisconsin was the same as that in Australia but differed from the predominant type of bovine origin isolated in England (20) and in Canada (21). In the

latter two countries the predominant type was 42D.

In a subsequent study in Wisconsin, Reid and Wilson (15) found the same predominant phage type of staphylococci in cows with acute mastitis, with chronic mastitis and with clinically normal udders as that reported by Seto and Wilson (18). Forty-four per cent of the typable isolates belonged to phage type 44A, and 30% belonged to type 42D. Further, type 44A staphylococci occurred more frequently in udders of acute and chronic mastitis-affected cows than did type 42D. However, the 42D type occurred slightly more frequently in normal udders than did the 44A type. A greater number of strains was phage typable from cows with acute mastitis than from those with chronic mastitis or from those with normal udders.

## MATERIAL AND METHODS