UNIVERSITY
OF MANITOBA

MAX RADY COLLEGE OF MEDICINE

Rady Faculty of Health Sciences

Bachelor of Science in Medicine Degree Program End of Term Final Report

Office of Graduate and
Advanced Degrees
Education in Medicine Max
Rady College
of Medicine
Rady Faculty of
Health Sciences
University of Manitoba

Student Name: Antonia Zhu

Date: 08/02/18

Project Title: Scleraxis as a prognostic marker of myocardial fibrosis in hypertrophic cardiomyopathy (SPARC) study

Primary Supervisor Name: Dr. Davinder S. Jassal

Department: Department of Internal Medicine

Co-Supervisor Name: Dr. Michael Czubryt

Department: Department of Physiology and Pathophysiology

Summary (250 words max single spaced):

Background: Hypertrophic cardiomyopathy (HCM) is the leading cause of sudden cardiac death (SCD) in individuals under 40 years of age. Evaluation of risk factors, including myocardial fibrosis characterized by late gadolinium enhancement on cardiac magnetic resonance (LGE-CMR), is critical for identifying high-risk HCM patients who require an implantable cardioverter defibrillator (ICD) for primary prevention of SCD. Given the high cost and limited availability of CMR in Canada, development of a serum biomarker of myocardial fibrosis would be a useful method for early risk stratification of HCM patients for SCD.

Objective: To determine whether scleraxis can serve as a biomarker of myocardial fibrosis that correlates with disease severity in HCM.

Methods: Between 2017 and 2018, adults with HCM at St. Boniface Hospital were recruited for the study. Patient characteristics, including demographics and SCD risk factors, were obtained. Cardiac function and myocardial fibrosis were assessed using transthoracic echocardiography (TTE) and CMR. Serum scleraxis was quantified using an enzyme-linked immunosorbent assay.

Results: A total of 46 patients were enrolled [58 ± 14 years (31 males)]. The average 5-year SCD risk was $2.3 \pm 1.3\%$. TTE and CMR confirmed evidence of asymmetric hypertrophy with preserved left ventricular ejection fraction, consistent with HCM. On CMR, LGE was present in 70% of participants. As compared to controls, serum scleraxis levels were up-regulated 5-fold in the HCM population.

Conclusions: This study supports the utility of scleraxis as a biomarker in HCM patients. Future studies are warranted to investigate scleraxis as a prognostic marker for identifying high-risk HCM patients who need an ICD.

Student Signature

Antonia Zhu

Primary Supervisor Signature

[Signature]

Acknowledgments: I gratefully acknowledge the sole or partial funding support from the following sponsors;

H.T. Thorlakson Foundation
Dean, College of Medicine
Research Manitoba

Manitoba Medical Service Foundation (MMSF)
Vice-Dean, Research Rady FHS
Health Sciences Centre Research Foundation
Heart and Stroke Foundation

Sponsorship if different than above;

Diagnostic Services of Manitoba; MMCF - Dr. Walter Nakielny Research Scholarship

MD/PHD MD/MSc. BSc. (MED) MED II Summer Research Program
Joe Doupe Annual Event Undergraduate Medical Student Research Symposium

INTRODUCTION

Sudden cardiac death (SCD) is a leading cause of mortality, accounting for 40 000 deaths in Canada annually.¹ It is defined as the sudden and unexpected loss of cardiac function, breathing, and consciousness.² In individuals less than 40 years of age, hypertrophic cardiomyopathy (HCM) is the leading cause of SCD.³

HCM is the most common inherited cardiovascular disease in Canada, affecting 1 in every 500 individuals. It is inherited as an autosomal dominant condition primarily caused by mutations in cardiac sarcomere genes.⁴ HCM is characterized by unexplained and asymmetric wall thickness >15 mm in at least one myocardial segment of the left ventricle, histological myocyte disarray, and coronary microvascular dysfunction.⁵ Another characteristic feature is abnormal myocardial fibrosis, which refers to the excessive deposition of extracellular matrix (ECM) components within the myocardium. Over time, fibrosis alters the stiffness and compliance of the heart and disrupts electrical signalling between cells. Consequently, myocardial fibrosis predisposes HCM patients to arrhythmias, heart failure, and an increased risk of SCD.⁶

Since the incidence of SCD is known to be higher in individuals with HCM (~1% per year), studies have focused on identifying risk factors for SCD and producing algorithms for risk assessment of HCM patients. In 2014, a multi-centre, retrospective, longitudinal cohort study developed and validated the most recent SCD risk prediction model, called HCM Risk-SCD, which has since been included in the European Society of Cardiology's guidelines.⁷ This model calculates the 5-year risk of SCD for HCM patients by considering several risk factors, including: age, maximum left ventricular (LV) wall thickness, left atrial (LA) diameter, maximum left ventricular outflow tract (LVOT) gradient, family history of SCD, non-sustained ventricular tachycardia, and unexplained syncope. Evaluation of these risk factors is critical for identifying high-risk HCM patients who require an implantable cardioverter defibrillator (ICD) for primary prevention of SCD. Based on the HCM Risk-SCD model, individuals with a 5-year SCD risk of: i) <4% are generally not indicated for an ICD; ii) 4 to 6% may be considered for an ICD; and iii) ≥6% are recommended to receive an ICD.⁷

Multimodality cardiac imaging, including transthoracic echocardiography (TTE) and cardiac magnetic resonance imaging (CMR), are commonly used in the diagnosis and management of individuals with HCM. TTE is used to delineate asymmetric left ventricular hypertrophy (LVH), which preferentially involves the interventricular septum (Figure 1A). TTE can also identify systolic anterior motion of the mitral valve leaflets, which results in LVOT obstruction and mitral regurgitation. When LVOT obstruction occurs, TTE can be used to measure the LVOT gradient at rest and with Valsalva. With intravenous gadolinium administration, contrast CMR can be used to non-invasively characterize the extent of myocardial fibrosis in HCM patients. In more than half of the HCM population, late gadolinium enhancement on CMR (LGE-CMR) appears in a patchy mid-wall pattern on hypertrophic segments (Figure 1B).⁴ The degree of LGE-CMR in HCM strongly correlates with the risk of SCD.⁸⁻¹³ In a multi-centre prospective study, LGE of ≥15% of LV mass demonstrated a two-fold increase in SCD risk.¹² This supports the role of LGE-CMR as an additional risk assessment tool for SCD. Although LGE-CMR provides significant prognostic information for HCM, it cannot be routinely used in the clinical setting due to its high cost and limited availability throughout Canada.

Development of a serum biomarker for myocardial fibrosis would be highly advantageous, as it would serve as a cost-effective alternative to LGE-CMR for early risk stratification of HCM patients for SCD. Previous studies have investigated serum biomarkers of myocardial fibrosis including type I collagen, matrix metalloproteinase-1 (MMP-1), and tissue inhibitor of

metalloproteinase-1 (TIMP-1). Type I collagen is a major component of the ECM, which when excessively accumulated, results in fibrosis. Matrix metalloproteinases (MMPs), including MMP-1, are proteinases involved in the degradation of ECM components, whereas tissue inhibitors of metalloproteinases (TIMPs), such as TIMP-1, bind to and inhibit MMPs. When the ratio of MMPs:TIMPs is low, fibrosis develops. Despite being associated with myocardial fibrosis in HCM, these biomarkers have not been found to correlate with SCD risk.^{14–16} As a result, there remains a need for a more prognostic biomarker of myocardial fibrosis that is capable of predicting SCD.

Scleraxis, a transcription factor, is a strong candidate for a novel biomarker of myocardial fibrosis in HCM. Fibrosis occurs when resident cardiac fibroblasts convert to myofibroblasts, which are pathologic cells that excessively synthesize ECM components including fibrillar collagens. Scleraxis has been shown to control this conversion process through direct transcription regulation of genes that encode matrix constituents (collagen and proteoglycans) and markers of myofibroblasts (α -smooth muscle actin and fibronectin).^{17,18} Furthermore, scleraxis has been shown to potentiate the TGF β /Smad3 signaling pathway, which is a key regulator of ECM production.¹⁷ Transforming growth factor β (TGF β) is an important pro-fibrotic agent involved in chronic fibrotic diseases. It induces expression of ECM genes and depends on Smad proteins for transduction of its signaling pathway. Scleraxis has been found to be required for TGF β /Smad3-mediated gene expression, as it facilitates transcription complex formation.¹⁷ *Ex vivo* data demonstrate that addition of scleraxis to rat cardiac fibroblasts induces their conversion to myofibroblasts, while inhibition of scleraxis expression or activity promotes reversal of the myofibroblast phenotype.¹⁷ *In vivo* deletion of the scleraxis gene results in significant loss of ECM in the murine heart.¹⁷ Altogether, these data implicate scleraxis as a critical regulator of fibrosis.

A preliminary study by Dr. Czubryt's collaborator, Dr. Moises Selman (Instituto Nacional de Enfermedades Respiratorias), demonstrated that scleraxis was detectable in the serum samples of 204 study participants [n=21 with idiopathic pulmonary fibrosis (IPF), n=126 with scleroderma, and n=57 controls] using an enzyme-linked immunosorbent assay (ELISA). Furthermore, serum scleraxis was found to differentiate between disease entities, as scleraxis levels were elevated by 60% in IPF patients and 35% in scleroderma patients in comparison to controls (p<0.001). However, it is unknown as to whether scleraxis can similarly serve as a serum biomarker of myocardial fibrosis that correlates with disease severity in HCM patients. This study is the first to address this significant knowledge gap.

OBJECTIVE

The objective of the study is to determine whether scleraxis can serve as a serum biomarker of myocardial fibrosis that is detectable and that correlates with disease severity in individuals with HCM.

METHODS

i. Study population

This is a retrospective study involving a total of 46 individuals with HCM. From June 2017 to July 2018, adults with HCM receiving care at St. Boniface Hospital were recruited for the study. Study participants must have been identified as having both TTE and CMR performed as part of routine clinical care (Figure 2). Exclusion criteria included contraindication to CMR and/or unwillingness to give informed consent. Baseline characteristics, including age, cardiovascular risk factors, HCM risk factors for SCD, medication use, and prior hospitalization for cardiovascular events (heart failure or acute coronary syndrome), were obtained through patient interview and chart review. In addition, each study participant's most recent 12-lead electrocardiogram (ECG)

analysis was acquired from the in-hospital Marquette Universal System for Electrocardiology (MUSE) database or community cardiologists in Winnipeg.

ii. Cardiovascular imaging (TTE and CMR)

All study participants received a TTE study (GE Vivid 7, Milwaukee, WI, USA). Interventricular septal thickness, posterior wall thickness, left ventricular ejection fraction (LVEF), and LA diameter were determined from 2-dimensional images.¹⁹ Using pulsed wave Doppler echocardiography, transmitral LV filling velocity at the tips of the mitral valve leaflets was obtained to measure diastolic parameters including E wave and deceleration time. Images were analyzed offline using GE Echopac (Vivid 7, Version 11.2, GE Medical Systems, Milwaukee, WI, USA) by one medical trainee and one level 3 TTE expert who were blinded to the clinical data. Measurements were then averaged for each parameter and verified by another level 3 TTE expert blinded to the clinical data.

All study participants underwent CMR imaging using a 1.5-T scanner (Avanto or Aera; Siemens Healthineers, Erlangen, Germany). Breath-hold cine imaging was performed using a segmented TrueFISP sequence with ECG gating to achieve 25 images covering the entire cardiac cycle. LGE images were obtained after 10 minutes of 0.2 mmol/kg injection of Gadolinium (Gd-DTPA, Magnevist, Schering, Germany) using a T1-weighted IR-prepared multislice TurboFLASH sequence with magnitude and phase sensitive reconstruction. Images were acquired sequentially in the short axis, followed by horizontal and vertical long axis images (TR 700 ms, TE 3.36 ms, FA 25°, 8 mm slice thickness, 1.6 mm interslice gap, matrix 256x192). LV cavity dimensions and systolic function were determined as per standard Society of Cardiovascular MRI criteria.²⁰ Images were analyzed offline using a cardiac 3D workstation, either syngo.via (Siemens Healthineers, Erlangen, Germany) and/or CMR42 (Circle Cardiovascular Imaging Inc., Calgary, Alberta). LV internal diameter, LV wall thickness, and LA diameter were analyzed by two medical trainees. Subsequently, measurements were averaged for each parameter and verified by a level 3 CMR expert blinded to the clinical data. LV cavity dimensions, systolic function, and the degree of LGE, determined by planimetric assessment of all short axis slices, were determined by two level 3 CMR experts who were blinded to the clinical data.

iii. Serum biomarker: scleraxis

Blood samples were collected via venipuncture at a single time point in the outpatient hematology laboratory at St. Boniface Hospital. A total of 4x5 ml blood samples were acquired from each study participant using BD Vacutainer SST tubes. An ELISA for scleraxis was performed using a Human Scleraxis Homolog A ELISA Kit (abx571821; abbexa.com). This kit provided a 96-well strip plate format, which was read on a standard microplate reader (BioTek Cytation 5 Imaging Reader) in the Institute of Cardiovascular Sciences at St. Boniface Hospital Albrechtsen Research Centre (SBRC). For comparison with the HCM study population, blood samples were collected from 20 healthy volunteers at SBRC, who were recruited as controls.

iv. Statistical analysis

The data were summarized as mean $\pm$ SD or number (percentage). A minimum sample size of 30 individuals was required to achieve an adequate power of 0.80 with an alpha level of 0.05 to correlate the level of serum scleraxis with the degree of LGE-CMR. A two-tailed p-value of <0.05 was considered statistically significant. Student's t-test was used to compare scleraxis levels of the control and HCM populations. Linear regression analyses were performed to determine whether there was a correlation between scleraxis levels and disease severity in HCM. Imaging parameters, including maximum LV wall thickness, LA diameter, and extent of LGE-CMR, were designated as surrogate markers of disease severity in HCM, given that they have

each correlated with SCD risk in previous studies. Analyses were performed using the SAS software 9.4 (SAS Institute).

RESULTS

i. Baseline characteristics and SCD risk factors

Of the 118 HCM patients approached between June 2017 and July 2018, a total of 46 individuals with HCM were enrolled in the study. Baseline characteristics and SCD risk factors are summarized in Tables 1 and 2, respectively. The mean age of the HCM study cohort was 58 ± 14 years (31 males). The most prevalent risk factors for SCD included history of non-sustained ventricular tachycardia (24%) and enlarged LA (≥ 48 mm) diameter (20%). Using the HCM Risk-SCD prediction model, the mean 5-year SCD risk was $2.3 \pm 1.3\%$, with more than half of the participants receiving a risk score of $< 4\%$. The predominant cardiovascular risk factors included hypertension (57%), smoking (43%), and hyperlipidemia (37%). At the time of recruitment, the majority of the HCM patients (70%) were asymptomatic with a functional capacity of New York Heart Association (NYHA) Class I. The most commonly used medications were beta blockers (46%), angiotensin-converting enzyme inhibitors/angiotensin receptor blockers (41%), and diuretics (33%). There was a low rate of major cardiovascular events among the study participants. A total of 7 of the 46 HCM patients experienced a major cardiovascular event in their lifetime, with stroke being the most common event (7%). In this study cohort, 4 HCM patients (9%) had a pre-existing ICD. Of these 4 individuals, 2 received an ICD for primary prevention of SCD. None of these individuals experienced ICD discharges.

The most recent ECG findings for the study population are summarized in Table 3. A large majority of the study population (83%) was in sinus rhythm, while only 3 individuals (7%) had underlying atrial fibrillation. ECG analysis for 5 participants either demonstrated paced rhythms or had never been performed due to the diagnosis of HCM being rather recent. More than half (57%) of the study participants demonstrated left ventricular hypertrophy with repolarization abnormalities, and a total of 5 HCM patients (11%) demonstrated left atrial enlargement.

ii. Cardiovascular imaging (TTE and CMR)

Echocardiographic parameters of the study population are summarized in Table 4. The LVEF was normal at $60 \pm 10\%$. The mean septal wall thickness was increased (> 15 mm) with asymmetric hypertrophy consistent with the diagnosis of HCM. Systolic anterior motion of the mitral valve (SAM) was present in 15 (33%) individuals. Of those with SAM, the mean LVOT gradient was 40 ± 36 mmHg at rest and 68 ± 46 mmHg with Valsalva. The mean E/E' ratio, which estimates left ventricular filling pressure, was elevated for both lateral (> 6.0) and septal (> 7.8) walls, confirming diastolic dysfunction in the HCM patients.

CMR parameters of the study population are summarized in Table 5. The LVEF was normal at $68 \pm 13\%$. Similar to TTE, there was asymmetric hypertrophy of the interventricular septum as compared to the posterior wall, consistent with HCM. In addition, LA diameter was increased on CMR, suggesting left atrial enlargement. LGE was present in 32 (70%) study participants, in a distribution of 1 (20%), 2 (13%), or ≥ 3 myocardial segments (37%) (Figure 3).

iii. Serum biomarker: scleraxis

The results of the ELISA analysis for serum scleraxis in the HCM patients, compared to controls, is depicted in Figure 4. The average serum scleraxis level of the HCM patients (0.0402 ± 0.008 ng/ml) was more than 5-fold greater than that of controls (0.0076 ± 0.004 ng/ml). Using an unpaired two-tailed t-test, the difference between the scleraxis levels of these two groups was found to be statistically significant with a p-value of < 0.01 . Linear regression analyses

did not identify any correlation between scleraxis levels and markers of disease severity in HCM, including maximum LV wall thickness, LA diameter, and extent of LGE-CMR.

DISCUSSION

As the rate of SCD remains elevated among individuals with HCM, there continues to be a need for improvement in the risk stratification of this patient population. In this study, there are three significant findings to note, including: i) the demographics of the study population were representative of the overall HCM population; ii) the study population mostly consisted of HCM patients with a low risk for SCD; and iii) scleraxis levels were significantly elevated in HCM patients, as compared to healthy individuals.

Our study cohort of Manitobans with HCM was predominantly middle-aged, male, and Caucasian. These demographics are similar to the general HCM population, which has been studied comprehensively worldwide. HCM is a common genetic cardiovascular disease that has been reported in more than 50 countries, with a prevalence of 1 in 500 individuals (0.2%).²¹ Due to the age-dependent expression of HCM mutations and cardiac hypertrophy, the prevalence of HCM is expected, and has been observed, to be higher (0.29%) in individuals older than 60 years of age.²² As such, the average age at diagnosis of HCM is typically in the fifth decade of life.²³ Although the average age of the global HCM population has not been established, numerous studies have characterized cohorts of HCM patients to be between 45 and 50 years of age. Literature comparing the prevalence and distribution of HCM among various ethnic groups is limited. However, it has been established that Hutterites and Mennonites are genetically predisposed to HCM.²⁴ Many studies note a male predominance among individuals diagnosed with HCM, with a female predominance only being reported among patients over 60 years of age.²⁵ The mechanism(s) underlying this phenomenon has not yet been elucidated. However, it has been proposed that gender-related differences in penetrance of HCM mutations and in hormone levels (e.g. estrogen) may be contributing factors.^{26,27} A total of 46 of approximately 2500 HCM patients in Manitoba were enrolled in our study. The average age of our HCM cohort was 58±14 years, which is higher than the average age of cohorts in previous studies of HCM patients (45 to 50 years). The majority of the study participants were male (67%), which is consistent with previous studies. Of the study participants, 83% were Caucasian, which may reflect the genetic predisposition in Hutterites and Mennonites as well as the demographics of the Manitoban population.

In our study, the HCM cohort mainly consisted of individuals with a low risk for SCD. Using the HCM Risk-SCD prediction model developed by O'Mahony et al.,⁷ we calculated the SCD risk of each study participant in our HCM cohort. The average 5-year SCD risk for our study population was 2.3±1.3%, with more than half of the study participants receiving a risk score of <4%. According to the aforementioned prediction model, an ICD is generally not indicated in patients with a 5-year risk of SCD <4%.⁷ While it may be an incidental finding that our study participants have a lower risk for SCD, it also may be representative of the overall HCM population. A review of the literature demonstrated that although HCM is the most frequent cause of SCD in young adults (<40 years), the vast majority of affected individuals remain undiagnosed or are not substantially impacted by the disease. In addition, SCD risk has been observed to be lower in individuals beyond 40 years of age, as older patients more frequently experience other cardiac or non-cardiac comorbidities, as opposed to morbidity or mortality associated with HCM.²¹

The results of this study support scleraxis as a potential clinical biomarker of myocardial fibrosis in individuals with HCM. Previous studies have demonstrated that serum biomarkers of myocardial fibrosis, including type I collagen, MMP-1, and TIMP-1, are associated with diastolic

dysfunction in HCM.^{14–16} The literature also demonstrates that matrix metalloproteinase-9 (MMP-9) and a precursor of B-type natriuretic peptide (BNP) were increased in HCM patients with myocardial fibrosis detected by LGE-CMR.²⁸ Furthermore, MMP-9 has been shown to correlate negatively with exercise capacity, while BNP has been associated with the need for septal reduction therapy.^{28,29} However, no biomarkers have been found to predict the risk of SCD in HCM patients to date. In this study, scleraxis was significantly elevated in HCM patients as compared to controls. Scleraxis holds great potential as a biomarker in the HCM population, since it has been established as a key regulator that controls fibroblast-to-myofibroblast conversion and therefore the synthesis of cardiac fibrosis.¹⁷ In contrast to previously studied biomarkers, which are indicative of existing levels of fibrosis, scleraxis is unique as it may serve as an earlier biomarker, given its upstream regulation of ECM production. Although scleraxis levels were increased in HCM patients, they were not shown to correlate with the severity of HCM, as represented by maximum LV wall thickness, LA diameter, and extent of LGE-CMR, on linear regression analyses. Previous studies that have investigated correlations between biomarkers and HCM severity have involved sample sizes of several hundred individuals.²⁹ As a result, it is likely that the small size of our HCM cohort is not sufficient to determine whether there is a correlation between scleraxis levels and disease burden detected on imaging. A prospective study with a larger patient population and several years of follow-up would be required to evaluate whether scleraxis levels predict the risk of SCD in the HCM population.

Limitations

There are a number of limitations to the current study. First, the imaging tests and blood draws were not performed at the same time due to the retrospective design of the study. This difference in time may have interfered with the detection of any correlation between scleraxis and imaging parameters. Second, serum samples were only obtained at a single time point. A longitudinal study evaluating serial scleraxis levels could potentially provide more prognostic information regarding changes in myocardial fibrosis within each study participant. Finally, although serum scleraxis levels were obtained from both control and HCM populations, imaging parameters could not be compared. Future studies would incorporate an age- and sex-matched control population for both imaging and biomarker analyses.

Future directions

This study demonstrated that scleraxis levels are up-regulated in HCM patients as compared to controls. The diagnostic and prognostic abilities of scleraxis in the HCM setting have yet to be elucidated. A multi-centre randomized screening trial with a larger sample size may be conducted to determine whether scleraxis is capable of diagnosing HCM via blood test, which could then be verified by TTE. Furthermore, a longitudinal and prospective study may be performed to evaluate scleraxis as a prognostic biomarker that identifies individuals with HCM who need aggressive therapy and/or ICD implantation for primary SCD prevention.

CONCLUSION

This study supports that scleraxis is a detectable serum biomarker for myocardial fibrosis in HCM. Future studies are warranted to evaluate scleraxis not only as a diagnostic test, but also a prognostic biomarker for identifying high-risk HCM patients who require aggressive management, including ICD insertion, for the prevention of SCD.

Figure 1. Transthoracic echocardiography (TTE) and cardiac magnetic resonance imaging (CMR) for a patient with hypertrophic cardiomyopathy. **A)** A parasternal long axis view on TTE demonstrating asymmetric hypertrophy of the interventricular septum (arrow) as compared to the corresponding inferolateral wall. LV, left ventricle; Ao, aortic root; LA, left atrium. **B)** A short axis view on CMR, following intravenous gadolinium administration, showing multiple foci of late gadolinium enhancement (LGE). These foci indicate regions of fibrosis within the mid-myocardium of the septum (arrows). RV, right ventricle; LV, left ventricle.

Figure 2. Clinical protocol: A total of 46 HCM patients were enrolled in the study between June 2017 and July 2018 inclusive. Following consent, each study participant underwent blood collection. Participants were subsequently interviewed for baseline characteristics acquisition. TTE and CMR images were analyzed for each HCM patient. Finally, an enzyme-linked immunosorbent assay (ELISA) was performed on blood samples from all study participants.

Figure 3. A) Of the 46 HCM patients, 32 (70%) demonstrated LGE on CMR, in a distribution of 1, 2, or ≥3 myocardial segments. **B)** The distribution of LGE-CMR among the HCM patients is depicted in pie chart format.

Table 1. Baseline characteristics of study population

	HCM patients (n=46)
Age (yrs)	58±14
Male	31 (67)
Height (cm)	173±10
Weight (kg)	88±18
Ethnicity	
Caucasian	38 (83)
Black	2 (4)
Asian	5 (11)
Other	1 (2)
Positive for HCM genetic marker	7 (15)
Family history of HCM	9 (20)
Cardiovascular risk factors	
Diabetes	7 (15)
Hypertension	26 (57)
Smoking	20 (43)
Hyperlipidemia	17 (37)
Family history of CAD	6 (13)
NYHA Functional Class	
Class I	32 (70)
Class II	12 (26)
Class III	2 (4)
Class IV	0
Blood pressure (mmHg)	
Systolic	131±15
Diastolic	78±11
Medications	
ACE inhibitors/ARBs	19 (41)
Beta blockers	21 (46)
Calcium channel blockers	13 (28)
Diuretics	15 (33)
Warfarin	4 (9)
Intervention	
Alcohol ablation	0
Myectomy	2 (4)
ICD implantation	4 (9)
Major cardiovascular event	
ICD discharge	0
VT/VF	2 (4)
Hospitalization for HF	1 (2)
ACS	1 (2)
Stroke	3 (7)

Data presented as mean ± SD or number (percentage). HCM (hypertrophic cardiomyopathy); yrs (years); CAD (coronary artery disease); NYHA (New York Heart Association); ACE (angiotensin-converting enzyme); ARB (angiotensin receptor blocker); ICD (implantable cardioverter defibrillator); VT (ventricular tachycardia); VF (ventricular fibrillation); HF (heart failure); ACS (acute coronary syndrome).

Table 2. Risk factors for sudden cardiac death

	HCM patients (n=46)
Age (yrs)	58±14
Unexplained syncope	5 (11)
Family history of SCD	2 (4)
History of non-sustained ventricular tachycardia	11 (24)
Maximum LV wall thickness (mm)	17±5
Left atrial diameter (mm)	42±7
Maximum LVOT gradient (mmHg)	68±46
5-year SCD risk as per HCM Risk-SCD model (%)	2.3±1.3

Data presented as mean ± SD or number (percentage). HCM (hypertrophic cardiomyopathy); yrs (years); SCD (sudden cardiac death); LV (left ventricular); LVOT (left ventricular outflow tract).

Table 3. ECG findings of study population

	HCM patients (n=46)
Sinus rhythm	38 (83)
Atrial fibrillation	3 (7)
Left atrial enlargement	5 (11)
Left ventricular hypertrophy	26 (57)

Data presented as number (percentage). HCM (hypertrophic cardiomyopathy)

Table 4. Echocardiographic parameters of study population

	HCM patients (n=46)
<u>LV cavity dimensions and systolic function</u>	
Wall thickness (mm)	
Septal	16±4
Posterior	12±2
Septal/posterior ratio	1.4±0.4
LVEDV (ml)	112±57
LVESV (ml)	43±25
Stroke volume (ml)	69±34
LVEDD (mm)	46±8
LVESD (mm)	29±9
LVOT diameter (mm)	20±3
LA diameter (mm)	42±7
LVEF (%)	60±10
Presence of SAM	15 (33)
LVOT gradient at rest (mmHg)	40±36
Basal LVOT gradient >30 mmHg	5/15 (33)
LVOT gradient provoked (mmHg)	68±46
<u>Diastolic parameters</u>	
E (m/s)	0.7±0.2
E' Lat wall (m/s)	0.09±0.03
E' IVS (m/s)	0.06±0.02
E/E' Lat wall	10±5
E/E' IVS	14±6
E/E' average	12±5
MV Dec Time (ms)	234±81

Data presented as mean $\pm$ SD or number (percentage). HCM (hypertrophic cardiomyopathy); LVEDV (left ventricular end-diastolic volume); LVESV (left ventricular end-systolic volume); LVEDD (left ventricular end-diastolic diameter); LVESD (left ventricular end-systolic diameter); LVOT (left ventricular outflow tract); LA (left atrium); LVEF (left ventricular ejection fraction); SAM (systolic anterior motion); E (peak transmitral LV filling velocity during early diastole); E' (peak velocity of mitral annulus in diastole); Lat (lateral); IVS (interventricular septum); MV (mitral valve); Dec (deceleration).

Table 5. CMR parameters of study population

	HCM patients (n=46)
Wall thickness	
Septal (mm)	20 $\pm$ 4
Posterior (mm)	13 $\pm$ 4
LVEDV (ml)	136 $\pm$ 44
LVEDV indexed to BSA (ml/m ²)	68 $\pm$ 20
LVESV (ml)	46 $\pm$ 34
LVESV indexed to BSA (ml/m ²)	23 $\pm$ 16
LVSV (ml)	89 $\pm$ 24
LVSV indexed to BSA (ml/m ²)	44 $\pm$ 12
LVEDD (mm)	46 $\pm$ 8
LVESD (mm)	31 $\pm$ 7
LVEF (%)	68 $\pm$ 13
LA diameter (mm)	52 $\pm$ 11

Data presented as mean $\pm$ SD or number (percentage). HCM (hypertrophic cardiomyopathy); LVEDV (left ventricular end-diastolic volume); BSA (body surface area); LVESV (left ventricular end-systolic volume); LVSV (left ventricular stroke volume); LVEDD (left ventricular end-diastolic diameter); LVESD (left ventricular end-systolic diameter); LVEF (left ventricular ejection fraction); LA (left atrium).

Figure 4. ELISA analysis: The average serum scleraxis level of the HCM patients was more than 5-fold greater than that of controls. *p < 0.01.

REFERENCES

1. Cardiac Arrhythmia Network of Canada. Sudden Cardiac Death. <https://canet-nce.ca/our-work/research/sudden-cardiac-death/>.
2. Heart and Stroke Foundation of Canada. Cardiac Arrest. <http://www.heartandstroke.ca/heart/conditions/cardiac-arrest>.
3. Cooper RM, Raphael CE, Liebrechts M, et al. New developments in hypertrophic cardiomyopathy. *Can J Cardiol*. 2017;33(10):1254-1265.
4. Elliott PM, Anastakis A, Borger MA, et al. 2014 ESC guidelines on diagnosis and management of hypertrophic cardiomyopathy: The task force for the diagnosis and management of hypertrophic cardiomyopathy of the European Society of Cardiology (ESC). *Eur Heart J*. 2014;35(39):2733-2779.
5. Biagini E, Coccolo F, Ferlito M, et al. Dilated-hypokinetic evolution of hypertrophic cardiomyopathy: Prevalence, incidence, risk factors, and prognostic implications in pediatric and adult patients. *J Am Coll Cardiol*. 2005;46(8):1543-1550.
6. Maron BJ. Contemporary insights and strategies for risk stratification and prevention of sudden death in hypertrophic cardiomyopathy. *Circulation*. 2010;121:445-456.
7. O'Mahony C, Jichi F, Pavlou M, et al. A novel clinical risk prediction model for sudden cardiac death in hypertrophic cardiomyopathy (HCM Risk-SCD). *Eur Heart J*. 2014;35(30):2010-2020.
8. Bruder O, Wagner A, Jensen CJ, et al. Myocardial scar visualized by cardiovascular magnetic resonance imaging predicts major adverse events in patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2010;56(11):875-887.
9. O'Hanlon R, Grasso A, Roughton M, et al. Prognostic significance of myocardial fibrosis in hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2010;56(11):867-874.
10. Maron MS, Appelbaum E, Harrigan CJ, et al. Clinical profile and significance of delayed enhancement in hypertrophic cardiomyopathy. *Circ Heart Fail*. 2008;1(3):184-191.
11. Green JJ, Berger JS, Kramer CM, et al. Prognostic value of late gadolinium enhancement in clinical outcomes for hypertrophic cardiomyopathy. *JACC Cardiovasc Imaging*. 2012;5(4):370-377.
12. Chan RH, Maron BJ, Olivetto I, et al. Prognostic value of quantitative contrast-enhanced cardiovascular magnetic resonance for the evaluation of sudden death risk in patients with hypertrophic cardiomyopathy. *Circulation*. 2014;130(6):484-495.
13. Ho CY, López B, Coelho-Filho OR, et al. Myocardial fibrosis as an early manifestation of hypertrophic cardiomyopathy. *N Engl J Med*. 2010;363(6):552-563.
14. Ellims AH, Taylor AJ, Mariani JA, et al. Evaluating the utility of circulating biomarkers of collagen synthesis in hypertrophic cardiomyopathy. *Circ Heart Fail*. 2014;7(2):271-278.
15. Fassbach M, Schwartzkopff B. Elevated serum markers for collagen synthesis in patients with hypertrophic cardiomyopathy and diastolic dysfunction. *Z Kardiol*. 2005;94(5):328-335.
16. Lombardi R, Betocchi S, Losi MA, et al. Myocardial collagen turnover in hypertrophic cardiomyopathy. *Circulation*. 2003;108(12):1455-1460.
17. Bagchi RA, Roche P, Aroutiounova N, et al. The transcription factor scleraxis is a critical regulator of cardiac fibroblast phenotype. *BMC Biol*. 2016;14(1):3-6.
18. Roche P, Nagalingam R, Bagchi R, et al. Role of scleraxis in mechanical stretch-mediated regulation of cardiac myofibroblast phenotype. *Am J Physiol Cell Physiol*. 2016;311(2):297-307.
19. Lang RM, Bierig M, Devereux RB, et al. Recommendations for chamber quantification. *Eur J Echocardiogr*. 2006;7(2):79-108.
20. Grothues F, Smith G, Moon J, et al. Comparison of interstudy reproducibility of cardiovascular magnetic resonance with two-dimensional echocardiography in normal subjects and in patients with heart failure or left ventricular hypertrophy. *Am J Cardiol*.

- 2002;90(1):29-34.
21. Maron BJ, Maron MS. Hypertrophic cardiomyopathy. *Lancet*. 2013;381:242-255.
22. Marian A, Braunwald E. Hypertrophic cardiomyopathy: Genetics, pathogenesis, clinical manifestations, diagnosis, and therapy. *Circ Res*. 2017;121:749-770.
23. Maron MS, Hellawell JL, Lucove JC, et al. Occurrence of clinically diagnosed hypertrophic cardiomyopathy in the United States. *Am J Cardiol*. 2016;117(10):1651-1654.
24. Orton N, Innes A, Chudley A, et al. Unique disease heritage of the Dutch-German Mennonite population. *Am J Med Genet A*. 2008;146A:1072–1087.
25. Olivotto I, Maron MS, Adabag AS, et al. Gender-related differences in the clinical presentation and outcome of hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2005;46(3):480-487.
26. Kubo T, Kitaoka H, Okawa M, et al. Gender-specific differences in the clinical features of hypertrophic cardiomyopathy in a community-based Japanese population: Results from Kochi RYOMA study. *J Cardiol*. 2010;56(3):314-319.
27. Dimitrow PP, Czarnecka D, Jaszcz KK, et al. Sex differences in age at onset of symptoms in patients with hypertrophic cardiomyopathy. *J Cardiovasc Risk*. 1997;4(1):33-35.
28. Roldán V, Marín F, Gimeno JR, et al. Matrix metalloproteinases and tissue remodeling in hypertrophic cardiomyopathy. *Am Heart J*. 2008;156(1):85-91.
29. Geske JB, McKie PM, Ommen SR, et al. B-type natriuretic peptide and survival in hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2013;61(24):2456-2460.