

The University of Manitoba  
ISOLATION AND CHARACTERIZATION OF X-LINKED  
MUTANTS IN THE NEMATODE PANAGRELLUS REDIVIVUS

by  
Daniel J. Burke

A THESIS  
SUBMITTED TO THE FACULTY OF GRADUATE STUDIES  
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS  
FOR THE DEGREE OF MASTER OF SCIENCE

Department of Zoology  
Winnipeg, Manitoba  
May, 1978

ISOLATION AND CHARACTERIZATION OF X-LINKED  
MUTANTS IN THE NEMATODE PANAGRELLUS REDIVIVUS

BY

DANIEL J. BURKE

A dissertation submitted to the Faculty of Graduate Studies of  
the University of Manitoba in partial fulfillment of the requirements  
of the degree of

MASTER OF SCIENCE

© 1978

Permission has been granted to the LIBRARY OF THE UNIVERSITY OF MANITOBA to lend or sell copies of this dissertation, to the NATIONAL LIBRARY OF CANADA to microfilm this dissertation and to lend or sell copies of the film, and UNIVERSITY MICROFILMS to publish an abstract of this dissertation.

The author reserves other publication rights, and neither the dissertation nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

#### ACKNOWLEDGEMENTS

The author expresses his deep appreciation to:  
Dr. B.J. Backmann for donating bacterial stocks,  
Dr. P.K. Isaac for the use of his microscopy facilities,  
Dr. J.J. Pasternak for communicating unpublished  
results and to the members of the supervisory  
committee. I wish to thank Ms. Sandi Chapman for  
her excellent technical assistance and many hours  
of useful discussion during the course of this work.

Finally, thanks are extended to Dr. Martin Samoiloff  
for continued patience, guidance and supervision.  
In every aspect of our friendship he has earned my  
deepest respect.

## TABLE OF CONTENTS

	Page
Abstract	1
Introduction	2
Materials and Methods	5
Nematode cultures	5
Mutagenesis	8
Results	11
Discussion	18
Figure 1	26
Figure 2	27
Figure 3	28
Figure 4	29
Figure 5	30
References	31

## ABSTRACT

A protocol for obtaining mutants in the nematode Panagrellus redivivus following mutagenesis with the alkylating agent ethyl methanesulfonate was established. The induced mutation rate for the X chromosome is 0.79% with an average mutation rate per locus of  $4.4 \times 10^{-4}$ . The total number of genes on the X chromosome coding for indispensable functions is estimated as approximately 300.

Of the 24 X-linked mutants that were isolated only 8 could be maintained as homozygotes due to the severity of the phenotype and the mating inefficiency of males. The mutants define 6 complementation groups and map within a limited region of the X chromosome.

Reasons for the failure to obtain stable visible mutants in earlier studies are discussed.

## INTRODUCTION

Collection of mutations which perturb a particular biological process allows the investigator to dissect the process in a stepwise fashion, leading to an understanding of the controlling mechanisms. This approach has been useful in elucidating the mechanisms regulating gene expression in procaryotes (Retznikoff, 1972; Goldberger, 1974). Brenner (1974) demonstrated the utility of a free-living nematode as a model organism for the study of development. Utilizing the self-fertilizing hermaphrodite Caenorhabditis elegans, he isolated 300 ethyl methanesulfonate (EMS) induced visible mutants and defined 77 complementation groups. This laid the groundwork for intensive study, at the cellular and molecular levels, into the mechanisms controlling the temporal sequence and spatial organization observed during development (see Edgar and Wood, 1977; Riddle, 1978).

As an alternative to collecting mutants, the investigator can perturb the process utilizing external agents such as antibiotics, microwave radiation and microbeam surgery. Generally the agents are broad in their action and results are indicative but not conclusive. Brenner and co-workers; Babu (1974); Ward et al. (1975); Herman et al. (1976); Hirsch and Vanderslice (1976); Klass et al. (1976); Hodgkin and Brenner (1977); Lewis and Hodgkin (1977); Sulston and Horvitz (1977);

utilizing the genetic approach, have recently met with great success. Samoiloff and co-workers, utilizing the former approach, have been less successful in developing the dioecous nematode Panagrellus redivivus as a model organism for studying post-embryonic development.

The work on P. redivivus focussed on stage specific post-embryonic events including: growth and ecdysis (Samoiloff and Pasternak, 1968; Samoiloff and Pasternak, 1969; Samoiloff, 1970; Samoiloff, 1973 a,b), gonad development (Boroditsky and Samoiloff, 1973), mating behaviour (Cheng and Samoiloff, 1972; Samoiloff et al., 1973; Samoiloff et al., 1974; Balakanich and Samoiloff, 1974), chemotaxis and osmotic tolerance (Pollock and Samoiloff, 1976), and the nutritional requirements of development (Abdulrahman and Samoiloff, 1975). In all cases, the processes could be interrupted with various inhibitors of macromolecular synthesis or modified by physical means. The results indicated that bursts of gene activity were required to provide precursor material necessary to complete a particular phase of the life cycle. The experimental data were consistent with a model of nematode development proceeding in quantum steps where the bursts of gene activity coincide with the morphological event of ecdysis. The inherent limitations of this approach meant that the experiments were inconclusive. In most cases, further analysis would have required either sophisticated biochemical techniques or a collection of

mutants to facilitate a genetic approach.

One study reported a single behavioural mutant (Pollock and Samoiloff, 1976), the only mutant isolated after an extensive analysis of many chemically induced behavioural variants (Pollock, 1974). This result is surprising in view of Brenner's success with the hermaphrodite C. elegans and raises questions concerning the utility of Panagrellus for genetic studies.

The present work resolves the following questions:

- (1) Can stable visible mutants of Panagrellus be obtained by chemical mutagenesis?
- (2) Are the mutations inherited in a classical Mendelian fashion?
- (3) What is the range of visible phenotypes possible?
- (4) Can the induced mutants be utilized to enhance further genetic studies in Panagrellus?
- (5) What are the reasons for the failure to obtain many mutants of Panagrellus in earlier studies?

## MATERIALS AND METHODS

Nematode cultures:

NG agar (Brenner, 1974): Prepared by dissolving 3g NaCl, 2.5 g Bactopeptone, 17 g agar and 1 ml cholesterol in ethanol (5 mg/ml) in 975 ml glass distilled water. After autoclaving (15 psi, 15 min), 1 ml CaCl<sub>2</sub> (1.0M), 1 ml MgSO<sub>4</sub> (1.0M) and 25 ml potassium phosphate buffer (1.0 M) pH 6.0 were added in order. Finally, 5 ml of an aqueous solution saturated with Nystatin (a fungal inhibitor) was added.

M9 buffer: 5 g NaCl, 6 g Na<sub>2</sub>HPO<sub>4</sub>, 3 g KH<sub>2</sub>PO<sub>4</sub> and 0.25 g MgSO<sub>4</sub>·7 H<sub>2</sub>O per litre.

C. elegans Ringer's (Hirsch et al., 1976): 0.821 g anhydrous sodium acetate, 0.018 g NaCl, 0.146 g KCl, 0.103 g CaCl<sub>2</sub>, 0.080 g MgCl<sub>2</sub>·6 H<sub>2</sub>O, 0.206 g Sucrose, 5 ml 0.05 M solution of Hepes buffer (Gibco) in 100 ml glass distilled water.

The nematodes used were from strain C-15 of Panagrellus redivivus obtained as previously described (Pasternak and Samoiloff, 1970). After nine consecutive generations of brother-sister matings, a single gravid female from the F<sub>10</sub> was used to initiate the stocks used in genetic studies. Stocks were maintained at room temperature (22°C) by culturing in petri plates containing NG agar. Worms were grown monoxenically on a mutant of Escherichia coli K12 generously donated by Dr. B. Bachmann, Yale University. The bacterium, strain JEF-8 (Coli Genetic

Stock Center #5678) carrying chromosomal markers thr 31, car B8, rel A, and met B<sub>1</sub> has chosen for car B8, a  $\Delta$  (car B) - (formerly pyr A locus). The non-reverting mutant is an auxotroph requiring methionine, arginine and uracil. The bacterial stock was initiated from a single colony, grown to saturation in Nutrient Broth (BBL) and tested for nutritional requirements. A sample was then frozen in 30% glycerol and stored for future use. Stocks were tested periodically for requirements. Occasionally bacterial cultures were contaminated and new stocks were established from the frozen samples. Before use, NG plates were seeded with JEF-8, maintained at 37°C overnight and allowed to cool to room temperature before use. The growth of the auxotroph on NG agar was slow, preventing the bacteria from over-growing the worms. The generation time for P. redivivus under these conditions is 4.0 days.

Small plates (35 mm x 10 mm) inoculated with a single gravid female can support growth such that a fertile F<sub>1</sub> generation is obtained, The F<sub>2</sub> population overgrows the plate and these F<sub>2</sub> larvae do not complete development. Intermediate-sized plates (60 x 15) will support growth such that approximately half of the F<sub>2</sub> population reach sexual maturity and produce progeny. A complete F<sub>2</sub> population can only be obtained by culturing

on large plates (100 x 15) but the culture becomes crowded with early F<sub>3</sub> offspring before the entire F<sub>2</sub> population mature. The F<sub>3</sub> larvae do not complete development under these conditions.

Plates contaminated by molds and bacteria were rendered monoxenic according to the glutaraldehyde method (Brenner, 1974). The glutaraldehyde kills the worms and contaminants but not embryos developing in utero. These embryos complete embryogenesis, larvae devour the female and crawl into the bacterial lawn within 36 hours. Ten of these larvae were transferred to a fresh plate.

Panagrellus redivivus is ovoviviparous. The embryo and the first larval stage (L<sub>1</sub>) develop in utero. The second (L<sub>2</sub>), third (L<sub>3</sub>) and fourth (L<sub>4</sub>) larval stages and the adults are free swimming. Each stage is separated by a molt. It is impossible to distinguish male and female larvae by morphological parameters. The female is competent to mate immediately after the final larval molt. To obtain virgin females it is necessary to grow individual larvae in isolation and select adult females. Panagrellus is transparent and mature females are easily distinguished from adult males. The female is larger and filled with eggs, the male is smaller with a conspicuous crescent shaped copulatory apparatus (spicule) and a well developed ductus deferens. Larvae were grown

in isolation on small NG plates seeded with bacteria.

#### Mutagenesis:

Mutations were induced with ethyl methanesulfonate (EMS) according to the method of Brenner (1974). A synchronously growing culture was utilized to ensure that all animals were at the same stage of gonad development. Synchrony was obtained by placing 50-100 gravid females on a fresh plate and allowing them to produce larvae for six hours. The females were removed and all remaining larvae developed in good synchrony. After 72 hours the worms were washed off the plate with 3 ml M9 buffer into a scintillation vial. 1 ml of freshly prepared EMS (0.2 M in M9 buffer) was added and the suspension allowed to sit at room temperature for 4 hours. At completion of mutagenesis an equal volume of 5% sodium thiosulfate was added to neutralize the mutagen and five minutes later the suspension was washed into a 250 ml separatory funnel and filled with M9 buffer. The worms were allowed to sediment by gravity and were washed in separatory funnels with M9 buffer, and finally poured into an empty petri plate. Worms were transferred in pairs to NG plates and inspected for progeny.

#### Handling and Observation:

Worms on agar were manipulated individually with a sharp wooden applicator stick sterilized in alcohol. Mass transfers were achieved by streaking large numbers of animals from crowded plates onto fresh media with

sterile inoculating loops. Worms were manipulated in fluid with micropipettes prepared as needed. Observations were made using a Wild M4A transilluminating dissecting microscope.

Chromosomes were stained by Feulgen's method (Stowell, 1945) with a slight modification. Whole mounts of male nematodes were unsatisfactory. The male gonad was dissected from the male by amputating the anterior portion of the worm behind the pharynx. The gonad was dissected on an albuminized slide in C. elegans Ringer's solution. The tissue was fixed in situ with Carnoy's fluid after the excess Ringer's solution had been removed with tissue paper. A clean cover slip was placed on top and the gonad was squashed. The slide was frozen on dry ice, the cover slip removed and the preparation flooded with absolute ethanol. The preparation was hydrated, hydrolyzed in 1 N HCl (60°C) for 11 min and stained in Schiff's reagent for 15 min. After bleaching in sulfurous acid, the staining was intensified with 0.5% Azur B in 1% sodium metabisulfate for 6 minutes - a method developed for visualizing Paramecium cytoplasmic bacterioids (Samoiloff, unpublished). The preparation was rinsed in distilled water, dehydrated, counterstained in Fast Green, cleared in xylene and mounted in DPX.

Whole worms were photographed on thin agar blocks (Sulston and Horvitz, 1977).

All photographs were taken with a Zeiss universal photomicroscope equipped with Nomarski optics and a flash attachment.

Bacterial cultures:

Bacteria were maintained in liquid Nutrient Broth (BBL) and subcultured bi-weekly. Cultures were grown to saturation at 37°C and stored at 4°C for up to one week.

## RESULTS

Chromosome Constitution: Cytological observations (Fig. 1) confirm a previous report (Hechler, 1970) that spermatogenesis produces two types of sperm with either four or five chromosomes. Oogenesis produces gametes with five chromosomes. The results indicate that the haploid complement of autosomes is four, there is an XO mechanism of sex determination and males are the heterogametic sex. Males are effectively haploid (hemizygous) with respect to the X chromosome and recessive mutations will automatically be expressed as they cannot be masked in a heterozygous form.

Isolation of Mutants: The progeny of two mutagenized parents are the product of two independently mutagenized gametes. It is extremely unlikely that the mutagen would have affected the same locus in both parental gametes, so none of the autosomal mutations will appear in the  $F_1$  in a homozygous form. Two classes of visible mutants in the  $F_1$  are possible: dominant (or semi-dominant) mutations and one-half of the recessive mutations generated on the X chromosome of eggs. In practice only the  $F_1$  female progeny are considered as they contain two independently mutagenized X chromosomes. One-half of the former class and all of the latter class will appear in the  $F_1$  males and would not be isolated. To ensure virginity of  $F_1$  females, each  $F_1$  larva was picked onto a

small NG plate seeded with bacteria and previously inoculated with a single wild type male. Large numbers of  $F_1$  animals (500-1000) were used to establish individual lines. All progeny from a pair of mutagenized parents were designated by a lower case letter and each individual is identified with an arabic numeral. Ten larvae from parental pair a are designated a1, a2, a3, ..... a10. Any variant isolated was named according to the phenotype (see below) and given the designation of the line from which it was isolated.

A single X-linked mutation in any  $F_1$  female should be expressed in one half of the  $F_2$  male progeny. All plates with  $F_2$  progeny were screened for X-linked mutations by visual inspection. To obtain autosomal recessive mutants, the animals from  $F_2$  plates with no X-linked mutants were transferred en masse to large plates. A single autosomal recessive mutation should be expressed in 1/16 of the  $F_3$  progeny.

Phenotypes of mutants: Panagrellus mutants are similar in phenotype to those described for C. elegans (Brenner, 1974). A summary of phenotypes isolated is presented in Table 1.

"Uncoordinated" animals with defective movement are the most common class of mutants that can be isolated. The uncoordinated phenotype ranged from slight to extreme paralysis. In some instances the smooth sinusoidal movement of the wild type had a superimposed shaking.

Table I. Preliminary characterization of mutants  
with respect to phenotype, expression and  
chromosomal location:

dm - dominant; sd - semi-dominant;

vb - variable expression.

Preliminary mutant designation	Chromosome	Comment
uncoordinated (unc)		
unc s215	x	severe unc; ♂ did not mate
unc f215	x	severe unc; ♂ did not mate
unc b205	x	severe unc; ♂ did not mate
unc c16	x	moderate unc; ♂ effective
unc t80	x	moderate unc; ♂ effective
unc c6	x	moderate unc; ♂ effective
unc t71	x	shaking superimposed; ♂ did not mate
unc a33	x	shaking superimposed; ♂ did not mate
unc k10	x	shaking superimposed; ♂ did not mate
unc f20	x	slight unc; ♂ effective
unc i4	x	moderate unc; ♂ effective
unc m11	x	severe unc; ♂ did not mate
unc d38	x	moderate unc; ♂ effective
unc ul3	x	moderate unc; ♂ effective
unc u44	x	moderate unc; ♂ effective
unc e71	x	severe unc; ♂ did not mate
unc i2	x	moderate unc; ♂ effective
unc ul	x	severe unc; ♂ did not mate
unc a49	autosome	severe unc; ♂ did not mate
dumpy (dpy)		
dpy q68	autosome	sd; ♂ effective
dpy j22	x	♂ did not mate
dpy dl	autosome	dm; vb
dpy a49	autosome	extreme dpy; ♂ did not mate
dpy m50	autosome	extreme dpy; ♂ did not mate
dpy s225	autosome	sd
small (sma)		
sma s40	autosome	sd
sma d225	autosome	sd
sma r7	autosome	sd

Preliminary  
mutant  
designation

Chromosome

Comment

---

long (lon)

lon i3                   x

lon t36                   x

lon t56                   x

blistered (bli)

bli s100                   x           small blister; vb

bli g100                   x           small blister; vb

roller (rol)

rol a50                   autosome       very dpy; linked double;  
♂ did not mate

"Dumpy" individuals made up the second largest class of mutants. The animals had a short and fattened appearance compared to wild type (Fig. 2).

"Small" - these animals were shorter than the wild type but did not have a fattened appearance (Fig. 2).

"Long" - these animals were longer and thinner than the wild type (Fig. 2).

"Blistered" - these mutants had a fluid filled blister on the cuticle (Fig. 3).

"Roller" - in these mutants, the animals rotated around their long axes as they progressed forward.

Mutant Characterization: Homozygous sex-linked recessive mutations were obtained in the following manner: mutant males from the  $F_2$  plate were crossed to virgin wild type females and virgin female progeny (heterozygotes) were isolated. Large numbers of hemizygous males were obtained by streaking the remaining  $F_2$  animals onto a large plate and allowing them to segregate  $F_3$  progeny. Since one-half of the  $F_2$  females would be heterozygous for the mutation, a large number of  $F_3$  males (1/4 of the total) were mutant. The mutant males were crossed to the virgin heterozygous females and homozygous males and females were isolated. Mutations that were expressed throughout the life cycle permitted isolation of homozygotes with relative ease since mutants (male and female) could be isolated as larvae. When it was impossible to distinguish mutant from wild type larvae, it became necessary to grow

individuals in isolation. Dominant and semi-dominant mutants were isolated as homozygotes by recurrent selection of homozygous females that did not segregate any wild type progeny after three generations. Autosomal recessive mutants were followed by picking mutant gravid females onto a fresh plate and inspecting the progeny. If all of the adults were wild type in appearance the female had mated with a wild type male and all the progeny were heterozygotes. If one half of the progeny were mutant the female had mated with a heterozygous male. None of the females produced only mutant progeny. In both above-mentioned cases, gravid females were then transferred to a fresh plate where they segregated mutants in differing proportions. In all cases of autosomal recessive mutants, the male could not effect mating and a homozygous line could not be established. These mutants were not considered further.

Of the 52 variants initially isolated, 34 segregated the mutant phenotype in subsequent generations. Eleven homozygous lines were established including 3 autosomal and 8 X-linked mutations (Table II). The remaining mutants, 16 sex linked and 7 autosomal, could not be established as homozygous lines. In most cases the males were ineffective at mating but some mutants were discarded because of variable expression of the phenotype.

Table II. Chromosomal location of mutants rendered homozygous.

<u>Mutants homozygosed</u>	<u>Chromosomal location</u>
unc t80	X
unc i2	X
unc i4	X
unc d38	X
unc u13	X
unc f20	X
unc u44	X
dpy q68	autosome
lon i3	X
sma s225	autosome
sma s40	autosome

Back Crosses: Homozygous virgin female mutants were crossed to wild type males and the appearance of the mutation in the progeny was scored. Sex linked recessive mutations were defined as those mutations which appeared in the progeny with expression restricted to males only. Mutations which satisfied this criterion but gave a substantial proportion of wild type males were considered to have low penetrance and were discarded. Mutations that expressed in both males and females were considered semi-dominant since the heterozygote had an intermediate phenotype. In all such cases, the males expressed the intermediate phenotype indicating that none of the semi-dominant mutants were sex linked. The semi-dominant mutants were not analyzed further. A single dominant mutant (a dumpy) was isolated but had a variable phenotype and was discarded. Back cross data are presented in Table III.

Genetic Nomenclature: Homozygous mutants were named according to the system established for C. elegans (Horvitz, pers. comm.). Briefly: gene names refer to relatively broad phenotypic categories. The general name refers to the phenotype originally detected or most easily scored. Different genes within the same general category are distinguished by arabic numerals. Each mutation name consists of a lower case letter (b) followed by an arabic numeral. Suffixes indicating particular characteristics of a mutation follow the mutation name.



Table III. Detection of sex linked mutations.

Phenotypes of progeny from homozygous mutant females (phenotype in parentheses) mated to wild type males are presented. Preliminary mutation designations are used.

(r) - recessive

(sd) - semi-dominant

Parental Female	Progeny				Chromosomal Location
(uncoordinated)	unc♀	wt♀	unc♂	wt♂	
unc t80		120	104		X (r)
unc i2		140	116		X (r)
unc i4		239	208		X (r)
und d38		102	95		X (r)
unc ul3		148	171		X (r)
unc f20		164	180		X (r)
unc u44		147	119		X (r)
(Long)	lon♀	wt♀	lon♂	wt♂	
lon i3		132	119		X (r)
(Dumpy)		♀	♂		
dpy q68		174	192		autosomal (sd)
		all intermediate			
(Small)		♀	♂		
dpy s225		128	149		autosomal (sd)
sma s40		169	150		autosomal (sd)
		all intermediate			

These suffixes include: dm, dominant; sd, semi-dominant; vb, variable phenotype. All mutations are assumed to be recessive to wild type unless specified. A list of Panagrellus mutations and gene names is given in Table IV.

The rationale for assigning gene names was derived from results of complementation tests as described below.

Complementation: Only the sex linked uncoordinated mutants were complemented. Crosses of two homozygous mutants often resulted in a low number of progeny because of the mating inefficiency of uncoordinated males. The number of progeny could be increased by utilizing a heterozygous female mated to a large number of hemizygous males. The heterozygous female was constructed by crossing three mutant larvae (all of the uncoordinated mutants isolated expressed the phenotype as larvae) with 10 wild type males. All normally moving progeny were heterozygous females and could be distinguished from males at the larval stages, hence the phenotypes allowed easy isolation of virgin females. Three heterozygous females (+/unc-a) were crossed with 20 males (unc-b) and the appearance of mutant females in the progeny indicated that the mutations did not complement. All complementation tests were confirmed with homozygotes. Mutations b1, b2 and b7 were non-complementing and were considered allelic. All other mutations complemented each other. Results are summarized in Table V.

Table IV. Genetic Nomenclature. Allelic mutations  
were determined by complementation tests.

x - X chromosome

a - autosome