

THE UNIVERSITY OF MANITOBA

SUBNIVEAN ACCUMULATION OF CO₂ AND
ITS EFFECTS ON WINTER DISTRIBUTION OF SMALL MAMMALS

by

CHERYL EVELINE PENNY

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE
OF MASTER OF SCIENCE

DEPARTMENT OF ZOOLOGY

WINNIPEG, MANITOBA

April 1978

SUBNIVEAN ACCUMULATION OF CO₂ AND
ITS EFFECTS ON WINTER DISTRIBUTION OF SMALL MAMMALS

BY

CHERYL EVELINE PENNY

A dissertation submitted to the Faculty of Graduate Studies of
the University of Manitoba in partial fulfillment of the requirements
of the degree of

MASTER OF SCIENCE

© 1978

Permission has been granted to the LIBRARY OF THE UNIVER-
SITY OF MANITOBA to lend or sell copies of this dissertation, to
the NATIONAL LIBRARY OF CANADA to microfilm this
dissertation and to lend or sell copies of the film, and UNIVERSITY
MICROFILMS to publish an abstract of this dissertation.

The author reserves other publication rights, and neither the
dissertation nor extensive extracts from it may be printed or other-
wise reproduced without the author's written permission.

ABSTRACT

Measurements of subnivean CO₂ during two winters (1974 - 1975 and 1975 - 1976) in six different habitats in the taiga 280 km northeast of Winnipeg, Manitoba revealed that subnivean CO₂ accumulated consistently in four habitats while not in two others. The CO₂ concentrations increased up to maxima of five times ambient levels. The accumulation of CO₂ was affected primarily by density and hardness of the snow.

A reduction in small mammal numbers was associated with the fall critical period. Accumulation of subnivean CO₂ was frequently associated with changes in small mammal distribution, involving reductions in the numbers of animals present at the sites of greater CO₂ concentrations.

ACKNOWLEDGEMENTS

This study was supported by a NRC grant from 1974 to 1976. I received an NRC bursary in 1973 to 1974.

At this time I would like to acknowledge the assistance of the following people without whose direction the project would not have got off the ground: Dr. D. Schindler, Dr. Neil Arnason, Dr. Terry Dick and especially Dr. W. O. Pruitt Jr. who created the Taiga Biological Station. Fellow students and friends such as Ed and Bonny Hill, Richard Leonard, Glenn Sutherland, Keith Yonge and Suzanne Levine were of great help. Especial thanks to Bill Conley and Don McDonald without whose support life would have been much more difficult, and lastly thanks to the many people who helped whenever they were up at Taiga Biological Station.

TABLE OF CONTENTS

ABSTRACT	i
ACKNOWLEDGEMENTS	ii
LIST OF TABLES	v
LIST OF FIGURES	viii
INTRODUCTION	1
MATERIALS AND METHODS	6
STUDY AREA	6
SECTION 1. CO ₂ SAMPLING SITES	21
CO ₂ SAMPLING SITES	21
CO ₂ ANALYSIS	22
SECTION 2. ENVIRONMENTAL PARAMETERS	24
MICROTOPOGRAPHY	24
CLIMATE	24
SNOW CONDITIONS	25
SUBNIVEAN TEMPERATURE	25
STATISTICAL ANALYSIS	26
SECTION 3. SMALL MAMMAL TRAPPING	28
RESULTS	32
SECTION 1. CO ₂ ACCUMULATION	32
SECTION 2. Factors influencing CO ₂ accumulation	38
SECTION 3. EFFECT OF CO ₂ ON SMALL MAMMAL DISTRIBUTION	56
SMALL MAMMAL NUMBERS	56
MOVEMENT OF SMALL MAMMALS IN RELATION TO CO ₂ ACCUMULATION	62
STATISTICAL ANALYSIS OF CHANGES IN SMALL MAMMAL DISTRIBUTION	76
DISCUSSION	89
SECTION 1. CO ₂ ACCUMULATION	89

SECTION 2. FACTORS INFLUENCING CO ₂ ACCUMULATION	91
SECTION 3. EFFECT OF CO ₂ ON SMALL MAMMAL DISTRIBUTION	94
SUMMARY	98
SECTION 1. CO ₂ ACCUMULATION	98
SECTION 2. FACTORS INFLUENCING CO ₂ ACCUMULATION	98
SECTION 3. EFFECT OF CO ₂ ON SMALL MAMMAL DISTRIBUTION	99

LIST OF TABLES

Table 1	Definition of environmental parameters as used in statistical analysis	27
Table 2	Monthly totals of CO ₂ samples (%/volume) for the winters of 1973-76.	33
Table 3	Measured CO ₂ concentrations (%/volume) and accumulations from six habitats during winters 1973-74, 1974-75, 1975-76 . . .	36
Table 4	Table of CO ₂ concentrations (%/volume) in soil (10 cm. below surface) and adjacent CO ₂ testing tubes in aspen plot taken once on each date in 1975-76.	39
Table 5	CO ₂ concentrations (%/volume) measured once on each date within the snow profile at the ground-snow interface, 20 cm. above the ground and 40 cm. above the ground in the alder tamarack bog at station 17	40
Table 6	Hourly sampling of subnivean CO ₂ (%/volume) concentrations at two stations on 28 and 29 March 1975	41
Table 7	Discriminant analysis of winter 1974-75 site-dependent and site-independent environmental factors with grouped CO ₂ . . .	42
Table 8	Discriminant analysis of winter 1975-76 site-dependent and site-independent environmental factors with grouped CO ₂ . . .	43
Table 9A	Means, standard deviations and F ratios of all environmental parameters in three different CO ₂ groups for winter 1975-76.	45
Table 9B	Within groups correlation matrix of all environmental parameters except CO ₂ for 1975-76.	46
Table 10A	Means, standard deviations and F ratios of all environmental parameters in three different CO ₂ groups for winter 1974-75.	47
Table 10B	Within groups correlation matrix of all environmental parameters except CO ₂ for 1974-75.	48

Table 11	Frequency of occurrence of variables as 3 significant variables in step-wise discriminant analysis of 6 study plots in winters 1974-75 and 1975-76.	50
Table 12	Mean, standard deviation and correlation coefficient of snow hardness (g/cm^2) and snow density (g/cm^3) in both winters	51
Table 13	Snowcover thickness (cm) with standard error and range in each habitat for 1974-75 and 1975-76.	52
Table 14	Mean, standard deviation and Pearson correlation coefficient of CO_2 (%/volume) and subnivean temperature ($^{\circ}\text{C}$.) in the winters of 1974 to 1975 and 1975 to 1976 using SPSS.	54
Table 15	Subnivean mean temperatures ($^{\circ}\text{C}$.) with standard error and range in each habitat for 1974-75 and 1975-76	55
Table 16	Number of small mammals live-trapped in the summers of 1974, 1975 and 1976 and throughout the study	57
Table 17	Significance of difference of number of captures of small mammals in 1973-76.	59
Table 18	Total number of live-trapped captures/100 trap nights on six study plots throughout the study.	60
Table 19	Significance of change in number of small mammals trap captures and trap failures before and after hiemal threshold	63
Table 20	Significance of number of captures in 3 small mammal trapping lines in alder tamarack bog before and after hiemal threshold	77
Table 21	Significance of number of captures in 4 lines of small mammal traps in alder tamarack bog before and after hiemal threshold	78
Table 22	Fienberg's contingency table test for black spruce bog 1975-76: observed values with A_1 capture of A_2 failure, B = number of the line (i.e., B_1 = line 1, B_2 = line 2, etc.) and C_{1+} = before and C_{2+} = after the hiemal threshold. A^+ , B^+ , or C_{1+} = combined values of A&/or B &/or C	80
Table 23	Fienberg's contingency table test for black spruce bog 1975-76: X^2 values.	81
Table 24	Fienberg's contingency table test for alder ridge ecotone 1975-76: observed values	83

Table 25 Fienberg's contingency table test for alder ridge ecotone
1975-76: χ^2 values 84

Table 26 Fienberg's contingency table test for aspen upland 1975-76
observed values 86

Table 27 Fienberg's contingency table test for aspen upland 1975-76
 χ^2 values 87

LIST OF FIGURES

Figure 1	Map of the study area	8
Figure 2	Black spruce bog plot	10
Figure 3	Aspen upland plot	12
Figure 4	Alder tamarack bog plot	14
Figure 5	Alder ridge ecotone plot looking from the ridge into the adjacent bog	16
Figure 6	Jackpine ridge plot	18
Figure 7	Jackpine sandplain plot	20
Figure 8	Distribution of small mammal captures in the aspen upland from October 1975 to April 1976	65
Figure 9	Distribution of small mammal captures in the alder ridge ecotone from October 1975 to May 1976.	68
Figure 10	Distribution of small mammal captures in the alder tamarack bog from November 1975 to April 1976.. . . .	71

INTRODUCTION

A disagreement was found in the literature as to whether or not CO₂ increases in concentration or accumulates to levels greater than ambient (.03%/volume CO₂) under the snowcover and to what extent small mammal distributions are affected by such accumulations.

Bashenina (1956) found CO₂ levels concentrations increased up to 4% near Moscow . In the areas where greater concentrations of CO₂ were recorded she found subnivean mouse nests raised higher above the ground than in areas with lesser concentrations of CO₂. Pichler (in Geiger 1965) also mentioned that CO₂ was present in the subnivean area of a rye crop in concentrations greater than expected. Kelley et al (1968) and Kelley and Weaver (1969) studied CO₂ accumulation under the snow in the arctic tundra and quoted a range of 0.034 to 0.1 percent by volume CO₂. Havas and Mäenpää (1972) found subnivean CO₂ to accumulate to levels of .03 to 0.105 percent/volume in a Hylocomium-Myrtillus type forest in Finland.

In contrast, Fuller and Holmes (1972) stated that subnivean CO₂ did not increase sufficiently to affect small mammal distributions in taiga. Reiners (1968) found soil CO₂ production to be approximately zero in January in a cedar swamp, fen and oak forest. Kelley et al (1968)

and Coyne and Kelley (1971) found subnivean CO_2 levels decreased to ambient once a snowcover of 20 cm. was established.

Therefore, the focus of this thesis is to answer the questions:

1. Does CO_2 accumulate in the subnivean space?
2. If it does, where and why do the accumulations occur?
3. Do any accumulations influence small mammal distributions?

CO_2 could accumulate by an increase either in production or in the strength of factors influencing retention. CO_2 is produced primarily by the decomposition of leaf litter and by plant and root respiration. It is also released from air pockets or reservoirs in the soil. Respiration from small mammals has been discounted as a significant contributor to amounts of subnivean CO_2 (Kelley et al 1968). Micro-organisms are the main decomposers of leaf litter (Benoit et al 1972) and have been shown to produce up to 28% of the CO_2 evolved on a woodland soil surface (Witkamp and Frank 1969).

Root respiration along with CO_2 released from small air pockets in the soil contributed the remaining 72% of CO_2 measured (Witkamp and Frank 1969). Root respiration was also reported by other authors as an important contributor (40.5%) of the over all CO_2 measured (Reiners 1968 Kosonen 1969, Brown and Macfayden 1969, and Anderson 1973). Root and micro-organism respiration are interrelated ($r=0.72$) with decomposition (Kucera and Kirkham 1971). Also, their combined contribution is the most significant in summer. Little soil testing has been conducted outside the summer months except for work done by Reiners

(1968). Therefore in winter another factor, plant respiration, may become more important. Both Hagerup (in Kalela 1962) and Havas and Mäenpää (1972) found significant plant respiration under the snow. So while one source of CO₂ may decline, another may increase in importance.

The production of CO₂ varies due to environmental conditions such as temperature, moisture levels, pH, wind as well as age and amount of litter. Many authors have found a positive correlation between plant or bacterial respiration and soil temperature, where respiration ceases at approximately -7°C. (Flanagan and Scarborough 1972, Benoit et al 1972). However, the effect of temperature was observed only within critical limits of moisture, litter age and depth and all were interdependent (Witkamp 1963, 1966). The effect of pH on CO₂ production was evident only outside the range 5 to 7.5 (Flanagan and Scarborough 1972). Kelley et al (1968) found that wind negatively influenced CO₂ accumulation.

Differences from summer to winter in the distribution of small mammals have been noted by a number of authors (Iverson and Turner 1972, Beer 1961, Andrzejewski and Mazurkiewicz 1976, Kalela 1962, Buckner 1966, Riewe 1973 and Morris 1969). These differences are in response to changes in the environment (Stickel 1960, McNab 1963). The subnivean environment is dark (Evernden 1966), with a relative humidity of 100%, quiet (Pruitt 1959a, 1960), and has a relatively constant warm temperature (0 to -7°C.) when compared to the

supranivean environment (Pruitt 1957, Fuller et al 1969).

Such a relatively constant environment develops when the snowcover reaches the hiemal threshold (15 to 20 cm.), and small mammals utilize or enter this as soon as it forms (Pruitt 1960). Without this environment small mammals could not survive northern winters (Pruitt et al 1961).

Once the hiemal threshold is established small mammals still can change their distributions within the habitat (West 1977). This distribution change is not necessarily correlated with temperature (Fuller 1977) or food availability (Gorecki and Gebczynska 1962, Grodzinski 1963, Chitty et al 1968, Flowerdew 1972, Andrzejewski and Mazurkiewicz 1976, Pernetta 1976, and Fairbairn 1977). Therefore something else in the subnivean environment may be triggering the changes in small mammal distribution.

Bashenina (1956) and others suggested CO_2 accumulation may influence small mammal distribution. The effect of CO_2 can be measured because small mammals respond by increased ventilation to increases in its concentrations (Galantsev and Tumanov 1969). The concentrations at which CO_2 could be detected by or affect small mammals is under study by a number of authors. Aquatic mammals have quite high tolerances to CO_2 , with 10% causing little effect on ventilation or heart rates (Irving in Soholt et al 1973). Burrowing and hibernating mammals normally experience up to 2.5 to 3% by volume CO_2 in their chambers (Kennerly 1964, Studier and Proctor

1971, Williams and Rausch 1973). These concentrations are very high compared to the ambient levels. But, non-diving, non-fossorial, non-hibernating rodents have a much greater sensitivity to CO₂ (Soholt et al 1973, Darden 1971) and genera such as Apodemus, Microtus and Clethrionomys exhibited a reduction in heart rate of 12 to 25% when exposed to CO₂ levels as low as 1 to 1.5% by volume (Galantsev and Tumonov 1969). Also, lengthy exposure to low levels of CO₂ (2%) led to increased ventilation and decreased rectal temperatures in Merriam's kangaroo rats (Dipodomys merriami) (Soholt et al 1973). Withers (1975) working with small (20 gm.) semi-fossorial Pseudomys albocinereus, found a decreasing rectal temperature in response to all levels of CO₂ greater than ambient. Baudinette (1974) found the effect of CO₂ was greater when the animal was exposed to CO₂ outside thermoneutral temperatures of 20 to 30°C. (Dawson 1955). The subnivean temperatures outside the nest (-7 to 0°C.) are definitely outside the small mammals thermoneutral zone. Nests allow increases in temperatures from 7 to 24 C. greater than environmental conditions (Hayward 1965) but would also probably be areas of greater CO₂ concentrations because of occupation.

Therefore, small mammals can respond to increased concentrations of CO₂. Temperature or food are not necessarily correlated with changes in small mammal distribution. The response to increases in CO₂ concentrations may be related to changes in the distribution of small mammals.

MATERIALS AND METHODS

STUDY AREA

Field research was conducted around the Taiga Biological Station at 51°05' N lat.; 95° 20'W long. (Figure 1). The area is within the taiga or boreal coniferous forest on the Precambrian Shield. Habitats investigated were black spruce bog (Figure 2), aspen upland (Figure 3), alder tamarack bog (Figure 4), alder ridge ecotone (Figure 5), jackpine ridge (Figure 6), and jackpine sandplain (Figure 7).

Average precipitation is 50 to 60 cm. rainfall equivalent with 40 cm. falling as rain and the rest as snow (Atmospheric and Environmental Services 1970). Temperatures range from - 50 to +40°C with maximum temperatures in June and July and minima in January and February. The first major snowfall, establishing the hiemal threshold (snowcover thickness of 15 - 20 cm.), occurs in late November to early December, and snow disappears in late April or May.


Square grids 0.4 ha. in size were established in each of the six habitat types. Each grid was composed of 100 small mammal trapping sites spaced equidistantly from each other in a 10 x 10 pattern

FIGURE 1

Map of the study area

△ Taiga Biological Station

• Study Plot



34 35 36 37 38
20'

AIKENS LAKE

MANITOBA-ONTARIO

EAST OF PRINCIPAL MERIDIAN - EST DU MERIDIEN PRINCIPAL

Scale 1:50,000 Échelle




FIGURE 2

Black spruce bog plot



FIGURE 3

Aspen upland plot



FIGURE 4

Alder tamarack bog plot



FIGURE 5

Alder ridge ecotone plot looking from
the ridge into the adjacent bog



FIGURE 6

Jackpine ridge plot




FIGURE 7

Jackpine sandplain plot



each site about 6.5 m. apart. One axis of the grid was labelled A to J and the other axis was numbered 1 to 10. Each grid point was permanently marked by a wooden 2.5 cm. x 2.5 cm. x 1.5 m. stake with alternate yellow and black decimeter markings.

SECTION 1

CO₂ SAMPLING SITES

The same trapping areas were sampled for CO₂ and small mammals. At each CO₂ sampling site a 2 m. piece of rubber tubing (0.6 cm. inside diameter) was taped to one of the wooden stakes of the grid. The upper end was taped to the wooden stake and the lower end lay on the ground-air interface away from the regular grid pathways. CO₂ testing sites were established to sample as much as possible of the microtopographical and habitat variation.

Each CO₂ measuring site was sampled once per month from mid December 1973 to early April 1974 unless it was plugged or otherwise inoperable. This inability to measure CO₂ at some sites led to missing data. One site was set up at each of the most elevated and at the most depressed sites of the grid in the aspen upland and jackpine ridge plots. Two adjacent stations were chosen in each of the relatively flat alder bog and black spruce bog plots.

The sites for 1974 to 1975 consisted of lines of sampling stations along a microtopographical gradient from the highest to the lowest points on

the jackpine ridge (five stations), aspen upland (ten stations), alder ridge ecotone (ten stations) and black spruce bog (five stations). A line of 10 CO₂ testing stations was also established across the alder tamarack bog and included all the microtopographical variation in the plot. Two CO₂ stations were established, one in the open, the other in the more densely forested part of the jackpine sandplain. Additional stations in differing microhabitats were placed within the aspen plot so that all soil and vegetation types could be sampled. On the alder ridge ecotone additional testing sites were located on the ridge area of the plot. No extra readings were taken in the bog because the alder tamarack bog plot and the ecotone plot are adjacent for half of their length. All stations were tested at least three times a month from early December to mid April.

CO₂ testing sites for 1975 to 1976 were the same as those of 1974 to 1975 with the addition of two subterranean (10 cm. deep) testing stations (one each in the greater CO₂ and lesser CO₂ accumulation areas in thick soil as determined from 1974 to 1975 data) in the aspen plot and two in presumed mouse holes in the alder ridge ecotone. These stations were checked three times a month or more.

CO₂ ANALYSIS

CO₂ concentrations were measured using a Dräger multigas analyzer

and analyzing tubes with a range of 0.01 to 0.3 \pm .005%/volume. The meter and tubes were calibrated against a gas chromatograph (see Appendix A). The multigas analyzer is essentially a small hand pump with a one-way valve. During the CO₂ measurement, a redox reaction takes place in the measuring tube, causing a white to purple color change. The procedure for utilizing the analyzer in the field is found in Appendix B.

One problem encountered during the readings was the shattering of the glass measuring tubes in the analyzer. The CO₂ analyzer was inoperative during January 1976 and caused the loss of a number of weeks of data before it could be replaced. CO₂ was usually measured once a week at each station from the onset of the hiemal threshold until the snowcover was no longer continuous or the testing station was exposed.

CO₂ readings were taken at two stations from 24 March 1975 to 27 March 1975 in an attempt to see if any variation occurred on day to day measurements. Kelley et al. (1968) found the least daily variation to occur between 1000 and 1600 hours. On 28 March 1975 hourly CO₂ readings were taken from 1200 to 2000 (dark) and 0600 to 1000 29 March 1975 to note CO₂ variation. Since the color change was not readily discernable when read in the light of propane or kerosene lamps, no further readings were taken after dark.

SECTION 2

ENVIRONMENTAL PARAMETERS

MICROTOPOGRAPHY

Accurate data were collected to find microtopographical gradients within the plots. This was done by establishing line transects down the A to J plot lines and along the #1 baseline of the plots. I measured the distance to the ground from a levelled string every 50 cm. along the transect lines. These readings were later corrected to make the lowest point equal to 0 and thus the total microtopographic variation in the plot could be visualized. Maps showing major features such as bare rock, large drops between lines and depressions between lines were also made.

CLIMATE

Temperature, precipitation, phase of the moon, wind speed, cloud cover and barometric pressure were recorded each day at TBS. Temperature readings were taken at 0800 hours and daily maxima were also recorded. Precipitation was recorded by the weather station in Bissett. Barometric pressure was recorded at about 1500, often 0800 and the time of the temperature maximum each day at TBS.

SNOW CONDITIONS

Snow profiles were taken twice monthly adjacent to rather than on each of the plots to decrease human disturbance, using the methods and instruments outlined in Klein et al. (1950). These measurements included snow thickness, number of snow layers, thickness of each layer, hardness (g/cm^2), density (g/cm^3), temperature of each snow layer, presence of ice layers, crystal type within each snow layer as well as any peculiarities noted in the profile.

In 1973 to 1974 snow thickness was measured adjacent to each station for each CO_2 reading throughout that testing. In 1974 to 1975 snow thickness was measured only in March and April at each station for each CO_2 reading. In 1975 to 1976 the number of ice layers present in the profile as well as snow thickness were measured at each station for each CO_2 reading taken from December to April.

SUBNIVEAN TEMPERATURE

Thermistors installed at 12 CO_2 testing stations during the summer were used to measure subnivean temperatures in the winters of 1973 to 1974 and 1975 to 1976. The thermistors were read using Sanwa Electric Instrument Company type P-3 ohmmeter. Their accuracy was not consistent so that temperatures in 1975 to 1976 were measured by using an ordinary thermometer ($\pm 0.5^\circ\text{C}$) in a protective metal case. These temperature readings were taken at least 50 cm. from the subnivean

opening of the CO₂ testing tube. This allowed individual temperature readings to be taken at each CO₂ station for every CO₂ reading taken. If ice layers were encountered, a wire was used to make a hole and then the thermometer was inserted to the ground level. The hole made by the thermometer was filled in after each measurement.

STATISTICAL ANALYSIS

To carry out the statistical analysis parameters were defined, numbered and entered as in Table 1. To test the effect of subnivean temperature on CO₂ concentration a Pearson correlation coefficient was determined for both winters of 1974 to 1976 using the Statistical Package for the Social Sciences (SPSS).

To test the relationship of snow hardness and snow density a Pearson correlation coefficient was determined for both winters of 1974 to 1976 using SPSS. The means, standard errors and ranges of subnivean temperature and snowcover thickness were listed for each of the six plots using the Condescriptive SPSS file.

A step-wise Discriminant analysis (from SPSS) was used to determine those environmental parameters which were most closely associated with CO₂. A classification of CO₂ concentrations into three ranges was used: less than ambient (less than 0.0175%), ambient (0.0175% to 0.025%) and above ambient (more than 0.025%). The means and standard deviations of each variable within each of these groups

TABLE 1

DEFINITION OF ENVIRONMENTAL PARAMETERS AS USED IN STATISTICAL ANALYSIS

Y_1 : CO_2 station number where Y_1 to Y_{10} represent the alder tamarack bog, Y_{11} to Y_{20} the alder ridge ecotone, Y_{21} to Y_{30} the aspen upland, Y_{31} to Y_{35} the black spruce bog, Y_{36} to Y_{40} the jackpine ridge and Y_{41} to Y_{42} the jackpine sandplain.

Y_2 : the change in microtopography where 0 is flat, 1 shows that the difference in height on both sides of the stake is larger than 100 cm, 2 shows that the difference in height on both sides of the stake is less than 100 cm.

Y_3 : thickness of soil measured in cm up to 1 m.

Y_4 : the percent of green vegetation in the ground cover of a 0.25 m quadrat at each CO_2 testing station prior to the first snowfall (not measured in 1974 to 1975).

Y_1 to Y_4 are site dependent factors that are constant over time.

Y_5 : the chronological sequence of CO_2 measurements 1 to 15 (1974 to 1975) and 1 to 11 (1975 to 1976).

Y_6 : CO_2 in %/volume $\times 10^4$.

Y_7 : snowcover thickness in cm.

Y_8 : subnivean temperature in $^{\circ}C$.

Y_9 : number of ice layers in the snow adjacent to the CO_2 station (not measured at all stations in 1974 to 1975).

Y_{10} : total number of hardness readings in gm/cm^2 , measured every two weeks.

Y_{11} : total number of density readings in g/cm^3 , measured every two weeks.

Y_{12} : range of snow temperature in the profile $\times 10$, measured every two weeks.

Y_{13} : ambient air temperature in $^{\circ}C$.

Y_{14} : barometric pressure in mmHg.

Y_{15} : change in barometric pressure.

is shown for both winters. The differences between groups is explained by the analysis. The minimum F statistic (comparing variances) was determined by the Wilks Method for Discriminant analysis.

A step-wise Discriminant analysis was done separately on the data from the different winters. Analysis of CO₂ was first carried out against all parameters, then against site-dependent parameters, site-independent parameters, the snow parameters and separately against subnivean temperatures and two snow parameters measured at each site. Separate analysis was done against the snow parameters because they were only measured once every two weeks. This led to missing cases which SPSS deleted in a listwise fashion and thus decreased the numbers eligible to each other analysis. Analysis was performed with the day effect and then without it. There was at least a 10% decrease in predictive value (percentage of "grouped" variables correctly classified by significant variables) in 1975 to 1976 in contrast to no effect in 1974 to 1975 when the day effect was removed. Therefore to make comparisons more realistic this day effect was removed before the two years were compared.

SECTION 3

SMALL MAMMAL TRAPPING

Small mammals were live-trapped throughout the study and toe-

clipped for later recognition. Sherman traps were used for all summer trapping and one trap was set at each of the 100 stakes on each plot. Sherman traps were not used before May nor after September because of the potential heat loss from their metal sides and the corresponding increased small mammal mortality. Trapping grids were established in June, July and August 1973 and were live-trapped immediately after they were established. The black spruce bog, the alder tamarack bog, the aspen upland and the jackpine ridge were each trapped with three trap checks per day. In order to get a broader idea of the number of type of species present than three-night trapping periods allowed (Smith 1966), I trapped these plots for 11 days each.

Summer trapping was carried out from May to September of both 1974 and 1975. Plots in the alder ridge ecotone and jackpine sandplain were established and trapped in September 1974. A three-night trapping schedule was followed and the traps were put on the plots in the early afternoon (Gentry and Odum 1957). Trapping was discontinued during any period of cold or rainy weather to prevent trap deaths due to exposure.

Masonite or Jolly-Board traps were used for all the winter trapping. These wooden traps transferred less heat than metal Shermans and thus reduced trap mortality. Also they were smaller in size than the Shermans and the animal had a smaller volume to keep warm. Winter trapping was done using wooden trap chimneys (Pruitt

1959b).

In the winter of 1973 to 1974 trapping was carried out in the alder tamarack bog and the black spruce bog at trap chimneys at every second stake for one three-night period.

In the winter of 1974 to 1975 trapping was carried out once a month in the alder tamarack bog, the alder ridge ecotone, the black spruce bog and the aspen upland from December to April. A trap chimney was placed at every fourth stake giving 25 trap chimneys in each habitat. All chimneys were pre-baited with raw oatflakes at least two different times before trapping began in December. This was intended to induce small mammals to include trap chimneys in their food search pattern and thus open subnivean tunnels to them in the winter. Prior to the hiemal threshold trapping was not normally carried out because it disrupts the normal subnivean space formation. In 1974, however, trapping was started in December even though the threshold was not reached because at that time it was felt that the data being lost were worth more than the disruptions of the subnivean space. Initially traps were checked every four hours; however, this proved too arduous so traps then were checked every six hours.

In the winter of 1975 to 1976 trapping was again carried out in the alder tamarack bog, the alder ridge ecotone, the black spruce bog and the aspen upland from October to April. The original pattern of the 1974 to 1975 trap chimneys was retained with 10 new chimneys added

to the alder tamarack bog and aspen upland habitats alternating A - J, #4 and #5 trap markers, five new chimneys were added to the black spruce bog at B10, D10, F10, H10 and J10, and five to the alder ridge ecotone at A4, C4, E4, G4 and I4.

During the winter of 1975 to 1976 extra bait was added to the traps and they were checked every eight hours. During this year there was no snow until December and the hiemal threshold was then quickly reached so that trapping was possible for all but a few weeks in December.

The bait used for small mammal trapping was a combination of peanut butter, rolled oats and occasionally chopped raisin. Weight, sex and species were recorded for each animal to determine population structure.

RESULTS

SECTION 1.

CO₂ ACCUMULATION

Carbon dioxide accumulations were considered to be important to small mammals when subnivean CO₂ concentrations were greater than ambient levels for two or more consecutive weekly samplings. This meant that CO₂ had been greater than ambient for at least one week and not just a momentary increase. It is important to note that I consistently measured ambient CO₂ at TBS to be 0.02%/volume whereas other studies have reported 0.028-0.035%/volume CO₂ and that I used the former value to define ambient conditions.

CO₂ accumulated at five of seven CO₂ sampling stations on the four habitats in 1973 to 1974, 22 of 39 in the six habitats in 1974 to 1975, and 23 of 44 in the six habitats in 1975 to 1976. Accumulations were recorded primarily in December, March to April and February to March, respectively in the winters of 1973-74, 1974-75 and 1975-76 (Table 2). The proportion of CO₂ measurements within the concentration ranges differed between habitats, but was the same from year to year within a given habitat. This was especially true in the aspen upland, alder tamarack bog, alder ridge ecotone and jackpine ridge (Table 3). The black spruce bog and jackpine sandplain exhibited a greater variation in CO₂ measurements between the two later years (Table 3).

TABLE 2

Monthly totals of CO₂ samples (%/volume) for the winters of 1973-76.

Plot and period	<0.015	0.015-0.025	0.025-0.045	0.045-0.065	>0.065	. . Number of CO ₂ measurements . .					
Black spruce bog											
1973-74: December	0	1	0	0	1						
January	0	0	1	0	0						
February	0	0	1	0	0						
March	0	1	0	1	0						
April	0	0	1	0	0						
1974-75: December	0	13	3	0	0						
January	0	14	1	0	0						
February	0	10	5	0	0						
March	0	12	8	0	0						
April	0	3	7	0	0						
1975-76: December	4	4	0	0	0						
January	4	11	0	0	0						
February	0	4	10	1	0						
March	0	7	6	7	0						
April	0	0	3	0	0						
Aspen upland											
1973-74: December	0	0	0	0	2						
January	0	1	1	0	0						
February	0	0	1	1	0						
March	0	2	0	2	0						
April	0	0	2	0	0						
1974-75: December	2	16	9	0	0						
January	1	14	11	1	0						
February	1	11	11	3	1						
March	1	12	16	3	0						
April	0	4	8	2	0						
1975-76: December	3	9	1	0	0						
January	1	19	2	0	0						
February	0	11	14	4	1						
March	0	13	16	9	0						
(without soil)											

. . . continued

TABLE 2 (CONTINUED)

Plot and period	. . . Number of CO ₂ measurements . . .				
	<0.015	0.015-0.025	0.025-0.045	0.045-0.065	>0.065
Alder tamarack bog					
1973-74: December	2	0	3	0	0
January	0	1	1	0	0
February	0	2	0	0	0
March	1	3	0	0	0
April	0	2	0	0	0
1974-75: December	0	24	3	0	0
January	2	24	1	0	0
February	2	22	3	0	0
March	1	19	7	0	0
April	2	21	5	0	0
1975-76: December	5	5	0	0	0
January	0	23	0	0	0
February	0	36	1	0	0
March	0	21	9	0	0
Alder ridge ecotone					
1974-75: December	0	20	5	0	0
January	3	18	4	0	0
February	3	19	5	0	0
March	0	19	7	0	0
April	1	14	6	2	0
1975-76: December	5	5	0	0	0
January	3	19	3	0	0
February	0	23	7	0	0
March	0	26	10	3	0
Jackpine ridge					
1973-74: December	0	2	0	0	0
January	0	2	0	0	0
February	0	1	1	0	0
March	0	2	1	0	0
April	0	2	0	0	0

. . . continued

TABLE 2 (CONTINUED)

Plot and period	<0.015	0.015-0.025	0.025-0.045	0.045-0.065	>0.065
. . Number of CO ₂ measurements . .					
Jackpine ridge (continued)					
1974-75: December	0	9	0	0	0
January	1	10	1	0	0
February	0	9	1	0	0
March	0	18	2	0	0
April	0	8	0	0	0
1975-76: December	0	5	0	0	0
January	1	13	1	0	0
February	1	12	2	0	0
March	0	13	2	0	0
April	0	0	3	0	0
Jackpine sandplain					
1974-75: December	0	6	0	0	0
January	0	5	1	0	0
February	0	3	1	0	0
March	1	5	2	0	0
April	0	4	0	0	0
1975-76: December	0	2	0	0	0
January	1	4	1	0	0
February	0	3	3	0	0
March	0	4	2	0	2

TABLE 3

Measured CO₂ concentrations (%/volume) and accumulations from six habitats during winters 1973-74, 1974-75 and 1975-76.

Plot and period	CO ₂ concentrations (%/volume)										Total no. of readings	No. of stations with accumulation	No. of stations without accumulation
	<0.015		0.015-0.025		0.025-0.045		0.045-0.065		>0.065				
	No.	%	No.	%	No.	%	No.	%	No.	%			
Black spruce bog													
1973-74	0	0	2	29	3	43	1	14	1	14	7	1	0
1974-75	0	0	52	68	24	32	0	0	0	0	76	3	2
1975-76	8	13	26	43	19	31	8	13	0	0	61	5	0
Aspen													
1973-74	0	0	3	25	4	33	3	25	2	17	12	2	0
1974-75	5	4	57	45	55	43	9	7	1	1	127	8	1
1975-76*	4	3	59	47	37	30	15	12	10	8	125	11	1
1975-75**	4	4	52	50	33	32	13	13	1	1	103	9	1
1975-76***	4	4	45	49	29	32	12	13	1	1	91	8	1
Alder tamarack bog													
1973-74	3	20	8	53	4	27	0	0	0	0	15	1	1
1974-75	7	5	110	81	19	14	0	0	0	0	136	3	6
1975-76	5	5	85	85	10	10	0	0	0	0	100	0	10
Alder ridge ecotone													
1974-75	7	6	90	71	27	21	2	2	0	0	126	7	2
1975-76	8	8	73	70	20	19	3	3	0	0	104	4	6
Jackpine ridge													
1973-74	0	0	9	82	2	18	0	0	0	0	11	1	1
1974-75	1	2	54	92	4	7	0	0	0	0	59	0	5
1975-76	2	4	43	81	8	15	0	0	0	0	53	1	4
Jackpine sandplain													
1974-75	1	4	23	82	4	14	0	0	0	0	28	1	1
1975-76	1	5	13	59	6	27	0	0	2	9	22	2	0

*With soil

**Without soil

***Without A6 and soil

The 1973 to 1974 CO₂ sampling stations were used as a pilot study to determine the location and number of sites to be used in the following two winters. Comparison between winter 1973 to 1974 and those of 1974 to 1975 and 1975 to 1976 was not emphasized because of the small sample size in 1973 to 1974 (only two sampling stations/plot and both those were not always in operation for the monthly samplings). The winter of 1973 to 1974 was also different in that the maximum snowcover thickness was double that found during the other two winters.

The aspen upland exhibited a progressive increase in CO₂ concentrations (Table 2). The same progressive increase can be seen in both years in the alder ridge ecotone, but the maximum CO₂ concentrations were less. The black spruce bog exhibited this increase in concentration in the winter 1975 to 1976. The jackpine sandplain showed a less pronounced but still evident increase in concentrations in 1975 to 1976.

In neither the alder tamarack bog nor jackpine ridge did CO₂ concentrations show an increase as the year progressed nor was there any obvious accumulation of CO₂ during the year (Table 2). The black spruce bog and jackpine sandplain did not show marked CO₂ increases in the winter 1974 to 1975.

CO₂ measurements taken in the soil at two stations in the aspen upland habitat were always equal to or greater than adjacent

measurements taken at the soil-snow interface (Table 4).

CO₂ measurements taken within the snow profile were found to be consistent within the measurable variation of the CO₂ meter, except on 12 January 1974 when the range was from 0.0175 to 0.03%/volume CO₂ (Table 5).

CO₂ measurements taken hourly from 1200 to 2000 March 28, and 0600 to 1000 29 March, 1975 ranged from 0.01 to 0.02%/volume which I considered to be insignificant (Table 6). Daily measurements taken at the same stake and same time of day from 24 to 27 March 1975 were found to be consistent at 0.0175 to 0.02%/volume CO₂.

CO₂ measurements of the air in two mouse holes were found to be less than or equal to ambient and were discontinued after two months because no occupation was evident.

SECTION 2.

FACTORS INFLUENCING CO₂ ACCUMULATION

The overall ability of a model based on environmental parameters to predict the CO₂ concentrations in 1974 to 1975 was slightly less than half that in 1975 to 1976 (38.6%, 61.3%) (Table 7 and 8). The lack of consistency between the predictive values of the two winters may be due to the variability of different lag effects of time on CO₂

TABLE 4

Table of CO₂ concentrations (%/volume) in soil (10 cm. below surface) and adjacent CO₂ testing tubes in aspen plot taken once on each date in 1975-76

	15 December 1975	23 December 1975	10 January 1976	16 January 1976	2 February 1976	9 February 1976	14 February 1976	24 February 1976	4 March 1976	13 March 1976	19 March 1976	31 March 1976
Station C7												
Subnivean		0.0175-0.03	inoperative meter	0.015-0.0175	0.04-0.045	0.04-0.045	0.04	0.055-0.06	0.035-0.04	0.035-0.04	0.06	0.04
Soil		0.025		0.04	0.06-0.065	0.07-0.075	0.06-0.07	0.10-0.11	0.08	0.07-0.075		
Station F4												
Subnivean	0.01-0.015	0.01-0.015		0.01-0.015	0.0175-0.02	0.02-0.025	0.035-0.04	0.04	0.02	0.02	0.04	0.02
Soil	0.02	0.015-0.0175		0.015-0.0175	0.02	0.02-0.025	0.04-0.045	0.04	0.02-0.025	0.02-0.025	0.05-0.06	0.08

TABLE 5

CO₂ concentrations (%/volume) measured once on each date within the snow profile at the ground-snow interface, 20 cm. above the ground and 40 cm. above the ground in the alder tamarack bog at station 17

Date of Measurement	Subnivean CO ₂	20 cm above ground CO ₂	40 cm above ground CO ₂
19 December 1973	0.0100	0.0125	0.0125
27 December 1973	0.0375-0.0400	0.0375-0.0400	0.0375-0.0400
12 January 1974	0.0200-0.0250	0.0175-0.0200	0.0250-0.0300
23 February 1974	*	0.0200-0.0250	0.0200-0.0250
17 March 1974	0.0200-0.0250	0.0200-0.0250	0.0200-0.0250
7 April 1974	0.0175-0.0200	0.0175-0.0200	0.0175-0.0200

*Blocked.

TABLE 6

Hourly sampling of subnivean CO₂ (%/volume)
 concentrations at two stations on 28 and 29
 March 1975

Date and time	Station #1	Station #2
28 March: 1200	0.0175-0.02	0.02
1300	0.010-0.015	0.0175-0.02
1400	0.0175-0.02	0.02
1500	0.015-0.0175	0.02
1600	0.0175-0.02	0.02
1700	0.015-0.0175	0.0175-0.02
1800	0.0175-0.02	0.02
1900	--	0.02
2000	0.02	0.02
2100 - 0600	not enough daylight to continue readings.	
29 March: 0600	0.01-0.015	0.0175-0.02
0700	0.015-0.0175	0.02
0800	0.015-0.0175	0.02
0900	0.0175-0.02	0.02
1000	0.0175-0.02	0.02

TABLE 7

Discriminant analysis of winter 1974-75
site-dependent and site-independent environmental factors
with grouped CO₂

Analysis	No. of cases	First 3 significant variables			Percent of "grouped" variables correctly classified by significant variables
With Y ₅ : Y ₆ vs. Y ₂ , Y ₃ , Y ₅ , Y ₇ -Y ₁₅	585	Y ₇	Y ₁₄	Y ₃	38.63
Y ₆ vs. Y ₂ , Y ₃ , Y ₅	585	Y ₃	-	-	20.00
Y ₆ vs. Y ₇ -Y ₉ , Y ₁₃ -Y ₁₅	585	Y ₇	Y ₁₄	Y ₁₃	36.41
Y ₆ vs. Y ₁₀ -Y ₁₂	585	Y ₁₂	Y ₁₁	Y ₁₀	50.77
Without Y ₅ : Y ₆ vs. Y ₂ , Y ₃ , Y ₅ , Y ₇ -Y ₁₅	585	Y ₇	Y ₁₄	Y ₃	40.00
Y ₆ vs. Y ₂ , Y ₃ , Y ₅	585	Y ₃	-	-	20.00
Y ₆ vs. Y ₇ -Y ₉ , Y ₁₃ -Y ₁₅	585	Y ₇	Y ₁₄	Y ₁₃	36.41
Y ₆ vs. Y ₁₀ -Y ₁₂	585	Y ₁₂	Y ₁₁	Y ₁₀	50.77
Alder tamarack bog (without Y ₅)					
Y ₆ vs. Y ₂ , Y ₃ , Y ₇ -Y ₁₅	135	Y ₇	Y ₁₂	Y ₁₃	40.74
Y ₆ vs. Y ₂ , Y ₃	135	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	135	-	-	-	-
Y ₆ vs. Y ₇ -Y ₉	135	Y ₇	-	-	31.85
Alder ridge ecotone (without Y ₅)					
Y ₆ vs. Y ₂ , Y ₃ , Y ₇ -Y ₁₅	135	Y ₁₀	Y ₉	Y ₂	42.96
Y ₆ vs. Y ₂ , Y ₃	135	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	135	Y ₁₀	Y ₁₁	Y ₁₂	21.85
Y ₆ vs. Y ₇ -Y ₉	135	Y ₇	Y ₉	-	39.26
Aspen upland (without Y ₅)					
Y ₆ vs. Y ₂ , Y ₃ , Y ₇ -Y ₁₅	135	Y ₃	Y ₁₀	Y ₇	25.19
Y ₆ vs. Y ₂ , Y ₃	135	Y ₃	-	-	39.26
Y ₆ vs. Y ₁₀ -Y ₁₂	135	Y ₁₀	-	-	*
Y ₆ vs. Y ₇ -Y ₉	135	Y ₈	Y ₇	-	45.93
Black spruce bog (without Y ₅)					
Y ₆ vs. Y ₂ , Y ₃ , Y ₇ -Y ₁₅	71	Y ₁₄	Y ₁₃	Y ₁₅	74.65
Y ₆ vs. Y ₂ , Y ₃	71	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	71	Y ₁₂	Y ₁₁	-	66.20
Y ₆ vs. Y ₇ -Y ₉	71	-	-	-	-
Jackpine ridge (without Y ₅)**					
Y ₆ vs. Y ₂ , Y ₃ , Y ₇ -Y ₁₅	70	Y ₁₃	Y ₇	Y ₁₂	71.43
Y ₆ vs. Y ₂ , Y ₃	70	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	70	Y ₁₀	-	-	*
Y ₆ vs. Y ₇ -Y ₉	70	-	-	-	-
Jackpine sandplain (without Y ₅)					
Y ₆ vs. Y ₂ , Y ₃ , Y ₇ -Y ₁₅	-	-	-	-	-
Y ₆ vs. Y ₂ , Y ₃ ***	-	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	-	-	-	-	-
Y ₆ vs. Y ₇ -Y ₉	-	-	-	-	-

*Reduced space dispersion matrix cannot be inverted.

**Groups 1 and 2 only.

***Groups 2 and 3 only.



TABLE 8

Discriminant analysis of winter 1975-76
site-dependent and site-independent environmental factors
with grouped CO₂

Analysis	No. of cases	First 3 significant variables			Percent of "grouped" variables correctly classified by significant variables
With Y ₅ : Y ₆ vs. Y ₂ -Y ₅ , Y ₇ -Y ₁₅	462	Y ₅	Y ₁₄	Y ₁₂	61.26
Y ₆ vs. Y ₂ -Y ₅	462	Y ₅	Y ₂	Y ₃	53.38
Y ₆ vs. Y ₇ -Y ₉ , Y ₁₃ -Y ₁₅	462	Y ₁₄	Y ₉	Y ₁₅	47.40
Without Y ₅ : Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅	462	Y ₁₁	Y ₁₄	Y ₇	51.08
Y ₆ vs. Y ₂ -Y ₄	462	Y ₂	-	-	21.43
Y ₆ vs. Y ₁₀ -Y ₁₂	462	Y ₁₁	Y ₁₂	Y ₁₀	60.17
Alder tamarack bog (without Y ₅)					
Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅ *	100	Y ₇	Y ₁₅	Y ₁₃	69.00
Y ₆ vs. Y ₂ -Y ₄	100	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	100	Y ₁₁	Y ₁₀	-	69.00
Y ₆ vs. Y ₇ -Y ₉	100	Y ₇	-	-	18.00
Alder ridge ecotone (without Y ₅)					
Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅	110	Y ₁₁	Y ₁₄	Y ₄	35.43
Y ₆ vs. Y ₂ -Y ₄	110	Y ₄	Y ₃	Y ₂	45.45
Y ₆ vs. Y ₁₀ -Y ₁₂	110	Y ₁₁	Y ₁₀	-	51.82
Y ₆ vs. Y ₇ -Y ₉	110	Y ₈	Y ₉	-	47.27
Aspen upland (without Y ₅)					
Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅	110	Y ₁₁	Y ₁₃	Y ₇	64.55
Y ₆ vs. Y ₂ -Y ₄	110	Y ₃	Y ₂	-	43.64
Y ₆ vs. Y ₁₀ -Y ₁₂	110	Y ₁₁	Y ₁₂	Y ₁₀	54.55
Y ₆ vs. Y ₇ -Y ₉	110	Y ₇	Y ₈	-	56.36
Black spruce bog (without Y ₅)					
Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅	55	Y ₈	Y ₉	Y ₇	54.55
Y ₆ vs. Y ₂ -Y ₄	55	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	55	Y ₁₁	Y ₁₀	Y ₁₂	56.36
Y ₆ vs. Y ₇ -Y ₉	55	Y ₈	Y ₉	Y ₇	78.18
Jackpine ridge (without Y ₅)					
Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅ **	49	Y ₃	Y ₁₄	-	73.47
Y ₆ vs. Y ₂ -Y ₄	49	Y ₃	-	-	61.22
Y ₆ vs. Y ₁₀ -Y ₁₂	49	-	-	-	-
Y ₆ vs. Y ₇ -Y ₉	49	Y ₇	Y ₉	Y ₈	46.94
Jackpine sandplain (without Y ₅)					
Y ₆ vs. Y ₂ -Y ₄ , Y ₇ -Y ₁₅ **	19	Y ₇	Y ₉	Y ₄	78.95
Y ₆ vs. Y ₂ -Y ₄	19	-	-	-	-
Y ₆ vs. Y ₁₀ -Y ₁₂	19	-	-	-	-
Y ₆ vs. Y ₇ -Y ₉	19	Y ₇	Y ₉	-	78.95

*Groups 1 and 2 only.

**Groups 2 and 3 only.

accumulation. The same parameters reacted differently in different winters. This might lead to lower predictive values, especially when many parameters are closely related and their effects cannot be disassociated from one another (Table 9 and 10). A 30% predictive value is expected from random distribution so that 60% predictive value is significant given the time lag associated between the measured variables and their effects on CO₂ production and accumulation (Neil Arnason, personal communication). The largest predictive values were found in the black spruce bog and jackpine ridge in 1974 to 1975 and the jackpine ridge and jackpine sandplain in 1975 to 1976. However, in both years these plots had the fewest number of cases of the six plots. In 1974 to 1975 the day effect was closely associated with the number of ice layers, snow hardness, density and ambient air temperature, which are themselves interrelated. In 1975 to 1976 the only strong association is seen between the day effect and snow density, with a lesser association between the day effect and number of ice layers in the snow profile.

To get a more accurate idea of what was associated with CO₂ accumulation the day effect was removed. Then, in the analysis where all parameters except date were analyzed against CO₂, snow characteristics were found to be of first significance in the alder tamarack bog (snowcover thickness) and the alder ridge ecotone (snow hardness) in 1974 and 1975, and in the alder tamarack bog (snowcover thickness), alder ridge ecotone (snow density), aspen upland (snow density) and jackpine sandplain (snowcover thickness) in 1975 to

TABLE 9A

Means, standard deviations and F ratios of all environmental parameters
in three different CO₂ groups for winter 1975-76

	Y ₂	Y ₃	Y ₄	Y ₅	Y ₇	Y ₈	Y ₉	Y ₁₀	Y ₁₁	Y ₁₂	Y ₁₃	Y ₁₄	Y ₁₅
Group 1*													
\bar{X}	3.6970	3.3939	2.6970	1.6364	24.6060	-70.9691	0.0000	13.6364	152.3636	29.6970	-122.1212	2,973.3938	1.3030
SD	0.6386	1.9361	1.2371	0.9293	7.3184	27.2275	0.0000	5.6282	13.2338	21.9353	93.3661	13.2665	0.4667
Group 2**													
\bar{X}	3.4364	3.3151	2.4452	5.4589	33.7123	-49.8630	0.4658	331.3354	195.3904	37.7877	-137.3972	2,967.4041	1.5479
SD	1.0568	1.8672	1.2207	2.8506	10.7979	27.7982	0.6233	751.6003	37.7193	21.4965	90.1039	15.2659	0.5763
Group 3***													
\bar{X}	3.0444	3.5556	2.3778	7.7333	37.0444	-26.3555	0.5778	309.0000	227.9556	24.7778	-50.0000	2,943.7554	1.9556
SD	1.2424	1.7781	1.4815	2.4346	10.7004	30.0874	0.4995	830.8787	45.7451	24.6096	85.3602	29.9333	0.5623
F ratio	4.0100	0.2941	0.6638	53.6762	14.6424	21.8578	11.8977	2.7200	39.5533	6.6197	16.3898	34.0000	14.3814

*Group 1: <0.015%/volume CO₂

**Group 2: 0.015-0.025%/volume CO₂

***Group 3: >0.025%/volume CO₂

TABLE 9B

Within groups correlation matrix of all environmental parameters except CO₂ for 1975-76

	Y ₂	Y ₃	Y ₄	Y ₅	Y ₇	Y ₈	Y ₉	Y ₁₀	Y ₁₁	Y ₁₂	Y ₁₃	Y ₁₄	Y ₁₅
Y ₂	1.0000												
Y ₃	0.2238	1.0000											
Y ₄	0.2859	0.2780	1.0000										
Y ₅	0.0830	-0.1339	0.1196	1.0000									
Y ₇	-0.0687	0.2558	0.0250	0.1815	1.0000								
Y ₈	-0.0246	0.0500	-0.1290	0.3157	0.1781	1.0000							
Y ₉	0.0707	-0.0661	0.0122	0.6484	0.1057	0.1885	1.0000						
Y ₁₀	0.1384	0.1707	0.2392	0.3508	0.0335	0.1200	0.4884	1.0000					
Y ₁₁	0.0570	-0.0865	0.0771	0.8739	0.1108	0.4690	0.4688	0.2125	1.0000				
Y ₁₂	-0.0119	0.2266	-0.0789	-0.1283	0.2045	-0.3227	-0.0080	-0.0354	-0.2006	1.0000			
Y ₁₃	-0.1111	-0.1620	-0.1184	0.0811	-0.1043	0.2529	-0.0189	-0.0994	0.2514	-0.2275	1.0000		
Y ₁₄	0.0469	0.1501	0.1093	0.0115	0.0280	-0.4429	0.2932	0.3896	-0.2114	0.3479	-0.4994	1.0000	
Y ₁₅	-0.0305	-0.0528	0.0454	0.0861	0.1417	0.1246	-0.0896	-0.3395	0.0369	0.0818	0.2224	-0.0497	1.0000

TABLE 10A

Means, standard deviations and F ratios of all environmental parameters
in three different CO₂ groups for winter 1974-75

	Y ₂	Y ₃	Y ₅	Y ₇	Y ₈	Y ₉	Y ₁₀	Y ₁₁	Y ₁₂	Y ₁₃	Y ₁₄	Y ₁₅
Group 1*												
\bar{X}	3.5333	3.0000	8.9333	38.2000	-56.6667	0.4000	221.8000	231.2000	20.0000	-118.0000	2,957.1333	1.6000
SD	1.0601	1.9640	4.0614	10.2901	19.3342	0.5071	387.2166	53.6646	11.0195	106.5833	36.9401	0.7368
Group 2**												
\bar{X}	3.6066	3.0328	9.5574	38.8524	-52.0492	0.5246	243.6721	239.0164	25.2623	-93.5246	2,966.8032	1.7377
SD	0.9179	1.9146	3.3542	13.0203	17.9670	0.5035	403.9604	62.9431	17.8987	92.0614	20.6559	0.7938
Group 3***												
\bar{X}	3.4375	3.8750	10.3750	30.1250	-43.7500	0.5625	439.3125	241.3750	17.8125	-32.8125	2,976.2500	2.0625
SD	1.2093	1.7464	4.2564	12.1758	23.6291	0.5123	641.7971	71.0069	17.0263	128.5557	20.4222	0.9287
F ratio	0.1920	1.3270	0.6193	3.1502	1.8597	0.4666	1.3157	0.1183	1.5438	3.1107	3.1102	1.4232

*Group 1: <0.015%/volume CO₂

**Group 2: 0.015-0.025%/volume CO₂

***Group 3: >0.025%/volume CO₂

TABLE 10B

Within groups correlation matrix of all environmental parameters except CO₂ for 1974-75

	Y ₂	Y ₃	Y ₅	Y ₇	Y ₈	Y ₉	Y ₁₀	Y ₁₁	Y ₁₂	Y ₁₃	Y ₁₄	Y ₁₅
Y ₂	1.0000											
Y ₃	0.1707	1.0000										
Y ₅	0.1498	-0.0234	1.0000									
Y ₇	-0.0533	0.1830	-0.1562	1.0000								
Y ₈	-0.2020	-0.3537	0.3707	-0.3317	1.0000							
Y ₉	0.0547	-0.1750	0.8625	-0.1452	0.3571	1.0000						
Y ₁₀	0.1884	-0.0437	0.7142	-0.4876	0.4343	0.5419	1.0000					
Y ₁₁	0.1413	-0.0280	0.8244	-0.3614	0.3665	0.7008	0.7459	1.0000				
Y ₁₂	0.0192	0.3089	-0.5215	0.4172	-0.4816	-0.4731	-0.4805	-0.6133	1.0000			
Y ₁₃	0.1507	-0.0885	0.8244	-0.3786	0.4673	0.6784	0.7421	0.8134	-0.6774	1.0000		
Y ₁₄	-0.0095	-0.2040	0.3746	-0.2149	0.1748	0.2587	0.2797	0.2192	-0.1559	0.1149	1.0000	
Y ₁₅	0.1320	0.0194	0.4355	-0.4393	0.3540	0.3906	0.5542	0.6319	-0.5433	0.7236	-0.1965	1.0000

1976. The variables most often included in 1974 to 1975 and 1975 to 1976 were snow hardness and snow density. Snow parameters accounted for 15 of 23 significant variables in 1974 to 1975 and 18 of 33 significant variables in 1975 to 1976 (Table 11).

There appeared to be a good relationship between snowcover density and hardness in 1974 to 1975 ($r=.72$), but this decreased in 1975 to 1976 ($r=.27$, Table 12). The variability of mean snowcover thickness measured in 1974 to 1975 was less variable than, and included in the range of the 1975 to 1976 measurements (Table 13). The alder tamarack bog and aspen upland had the thickest snowcover in both winters. Snowcover thickness while being the first significant variable chosen is greatly influenced by changes in barometric pressure and ambient temperature (Table 10B). The effects of these were closely correlated to hardness and density (Table 10) and possibly masked the snow station factors. Also the snowcover measurements were closely associated with time (Table 9 and 10).

These could be indicative of the overall effect of weather on snowcover development. In the analysis of the snow station parameters, F values sufficiently large to be analyzed were obtained in four of six cases in both 1974 to 1975 and 1975 to 1976. Their predictive values were similar to those found for the overall grouping probably because they were often included in the first three significant variables.

TABLE 11

Frequency of occurrence of variables as 3 significant variables
in step-wise discriminant analysis of 6 study plots
in winters 1974-75 and 1975-76

Year and variable	No. of times included in first 3 significant variables (excluding Y ₅)
1974-75: Y ₂	1
Y ₃	2
Y ₇	3
Y ₈	0
Y ₉	1
Y ₁₀	5
Y ₁₁	2
Y ₁₂	4
Y ₁₃	3
Y ₁₄	1
Y ₁₅	1
Total	23
1975-76: Y ₂	2
Y ₃	4
Y ₄	3
Y ₇	4
Y ₈	1
Y ₉	2
Y ₁₀	4
Y ₁₁	6
Y ₁₂	2
Y ₁₃	2
Y ₁₄	2
Y ₁₅	1
Total	33

TABLE 12

Mean, standard deviation and correlation coefficient
of snow hardness (g/cm^2) and snow density (g/cm^3) in both winters

	Number of readings	\bar{X}	SD	Correlation coefficient (r)
1974-75				
Hardness	269	226.7700	373.9700	0.7218
Density	267	237.8800	51.1800	
1975-76				
Hardness	265	265.3018	671.0449	0.2728
Density	265	192.7207	41.1680	

TABLE 13

Snowcover thickness (cm) with standard error and range
in each habitat for 1974-75 and 1975-76

	Number of readings	Missing readings	\bar{X}	SE
1974-75				
Alder tamarack bog	99	36	35.5	1.9
Alder ridge ecotone	98	37	28.7	1.5
Aspen upland	94	41	35.4	1.7
Black spruce bog	51	24	32.4	2.1
Jackpine ridge	51	24	28.6	1.9
Jackpine sandplain	22	8	31.3	3.5
1975-76				
Alder tamarack bog	100	10	38.5	0.9
Alder ridge ecotone	104	6	31.0	1.1
Aspen upland	102	8	36.6	1.0
Black spruce bog	55	0	29.5	1.0
Jackpine ridge	53	2	22.9	1.7
Jackpine sandplain	22	0	28.5	1.7

Three of the factors thought to have the greatest influence on CO₂ accumulation: subnivean temperature, snowcover thickness and ice layers in the snow profile; were tested against the CO₂ groups in each of the six plots in both years (Table 7 and 8). In only one case, the black spruce bog, did the predictive value increase markedly (55 to 78%) but in the alder tamarack bog there was a marked decrease (69 to 18%) in predictive value. Therefore the combination of these three factors was not effective in predicting CO₂ concentrations. In only one case was subnivean temperature found to have significant effect on the percentage of "groups" correctly classified. Subnivean temperature was not as significant a contributor to a predictive value as the snow characteristics especially snowcover thickness.

The mean subnivean temperatures for both winters were very similar, but the correlation coefficients showing the relationship between it and CO₂ concentrations were quite different for the two winters (Table 14). So some other environmental factor must have influenced the correlation. The winter of 1974 to 1975 had colder mean temperatures than 1975 to 1976 in all habitats except the alder ridge ecotone where it was 0.3°C. warmer (Table 15). This is within the region of error of the thermometer. The aspen upland had the highest mean subnivean temperatures in both winters. The black spruce bog and jackpine ridge had the lowest subnivean temperatures in 1974 to 1975 and 1975 to 1976.

TABLE 14

Mean, standard deviation and Pearson correlation coefficient of CO₂ (%/volume) and subnivean temperature (°C.) in the winters of 1974 to 1975 and 1975 to 1976 using SPSS

	Number of Measurements	\bar{x}	SD	Correlation coefficient
1974-75				
CO ₂ (%/volume)	550	.0259	.0132	0.1990
Subnivean temperature (°C)	250	-4.88	2.10	
1975-76				
CO ₂ (%/volume)	434	.0263	.0116	0.5415
Subnivean temperature (°C)	436	-4.20	3.21	

TABLE 15

Subnivean mean temperatures ($^{\circ}\text{C}.$) with standard error and range
in each habitat for 1974-75 and 1975-76

	No. of readings.	Missing readings	\bar{X}	SE
1974-75				
Alder tamarack bog	75	60	-5.21	0.24
Alder ridge ecotone	76	59	-4.53	0.23
Aspen upland	44	91	-3.20	0.22
Black spruce bog	24	51	-6.65	0.38
Jackpine ridge	14	61	-5.89	0.44
Jackpine sandplain	17	13	-5.94	0.30
1975-76				
Alder tamarack bog	110	0	-4.32	0.27
Alder ridge ecotone	106	4	-4.83	0.34
Aspen upland	108	2	-2.98	0.25
Black spruce bog	50	5	-4.46	0.49
Jackpine ridge	42	13	-5.13	0.59
Jackpine sandplain	20	2	-4.28	0.69

The site-dependent factors (change in microtopography, thickness of soil and percentage of green ground cover) did not influence the predictive value consistently between plots or years when tested alone or with other environmental parameters, although they should have been what determined CO₂ production. This suggests that the most important site-dependent parameters were not measured. Site-independent factors contributed to the predictive values to the greatest degree. The formation of the snow layer was important in contributing to the predictive value. Snow density, hardness and thickness were especially important in association with CO₂ accumulation.

SECTION 3.

EFFECT OF CO₂ ON SMALL MAMMAL DISTRIBUTION

SMALL MAMMAL NUMBERS

The total numbers of small mammals of the eight species caught during this study are shown in Table 16. Microsorex hoyi is not listed because it was not possible to differentiate between it and Sorex cinereus in the field. All individuals of these two species were labelled Sorex cinereus. The most frequently captured species was Clethrionomys gapperi; its numbers were greater than all the other species combined. Microtus pennsylvanicus was the second most common species in 1975 to 1976. For the other trapping periods

TABLE 16

Number of small mammals live-trapped in the summers of 1974, 1975 and 1976 and throughout the study.

SPECIES	JUNE-SEPT. 73	MAY-SEPT. 74	MAY-SEPT. 74 & DEC. 74-Apr. 75	MAY-SEPT. 75	MAY-SEPT. 75 and OCT. 75 - APR.76
<u>Clethrionomys</u> <u>gapperi</u>	44	109	109	198	239
<u>Microtus pennsylvanicus</u>	1	2	2	3	36
<u>Peromyscus</u> <u>maniculatus</u>	6	11	11	11	11
<u>Zapus hudsonius</u>	0	1	1	2	2
<u>Sorex cinereus</u>	2	17	17	2	6
<u>S. arcticus</u>	0	0	0	0	1
<u>Blarina brevicauda</u>	10	20	20	3	3
<u>Synaptomys borealis</u>	0	0	0	2	2
number live-trapped	63	160	160	221	300
number trap nights	4000	5400	7100	8700	10,960

(summer 1973 and 1974 to 1975) Peromyscus maniculatus, S. cinereus and Blarina brevicauda were the next most common species and had greater numbers captured than Microtus.

The total number of individuals captured in 1975 to 1976 was greater than in either of the other two years. Clethrionomys and Microtus increased markedly in number from 1973 to 1976. Synaptomys borealis and Sorex arcticus were captured for the first time in 1975 to 1976. Peromyscus numbers were similar in 1975 to 1976 to the other years. Zapus hudsonius was rarely captured. Sorex cinereus and Blarina captures dropped markedly from their greatest numbers in 1974 to 1975. The decreases and stationary numbers are even more significant because there was an increase in trap nights (Table 17; $p < 0.005$) No new animals were added to the total during the winter of 1974 to 1975, in contrast to the winter of 1975 to 1976 when there was a marked increase in new individuals trapped, particularly Clethrionomys and Microtus (Table 16).

The numbers of small mammals captured per 100 trap nights in each month in each habitat can be seen in Table 18. C. gapperi and P. maniculatus were captured in all plots. M. pennsylvanicus was captured in the alder tamarack bog, alder ridge ecotone and aspen upland. S. cinereus and B. brevicauda were captured in all habitats except the jackpine sandplain. S. arcticus was captured once in the alder tamarack bog. Synaptomys borealis was captured rarely in the alder tamarack bog and alder ridge ecotone. Z. hudsonius was

TABLE 17

Significance of difference of number of captures of small mammals in 1973-76

	Trap nights	Observed captures	Expected captures	χ^2
1973	4,000	63	63.00	0 (p = 0, df = 2)
1974-75	7,100	160	111.80	20.780 (p<0.005, df = 2)
1975-76	10,960	300	172.62	93.996 (p<0.005, df = 2)

TABLE 18

Total number of live-trapped captures/100 trap nights
on six study plots throughout the study

	<i>Clethrionomys gapperi</i>	<i>Microtus pennsylvanicus</i>	<i>Peromyscus maniculatus</i>	<i>Sorex cinereus</i>	<i>S. arcticus</i>	<i>Synaptomys borealis</i>	<i>Blarina brevicauda</i>	<i>Zapus hudsonius</i>
Jackpine sandplain								
September 1974	1.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0
May 1975	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0
June 1975	0.0	0.0	0.7	0.0	0.0	0.0	0.0	0.0
July 1975	0.0	0.0	1.7	0.0	0.0	0.0	0.0	0.0
August 1975	0.0	0.0	1.7	0.0	0.0	0.0	0.0	0.0
September 1975	0.0	0.0	6.7	0.0	0.0	0.0	0.0	0.0
Jackpine ridge								
August 1973	4.3	0.0	0.0	0.1	0.0	0.0	0.1	0.0
June 1974	6.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
July 1974	2.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
August 1974	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
September 1974	10.0	0.0	1.0	1.7	0.0	0.0	0.7	0.0
May 1975	4.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
June 1975	3.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
July 1975	8.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
August 1975	3.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
September 1975	6.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Black spruce bog								
August 1973	3.3	0.0	0.0	1.0	0.0	0.0	1.0	0.0
June 1974	0.7	0.0	0.3	0.0	0.0	0.0	0.0	0.0
July 1974	6.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0
August 1974	5.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
December 1974	6.7	0.0	0.0	1.3	0.0	0.0	0.0	0.0
February 1975	4.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
March 1975	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
July 1975	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
August 1975	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
September 1975	4.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
November 1975	25.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0
January 1976	2.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
February 1976	2.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
April 1976	3.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Alder tamarack bog								
September 1973	2.2	0.0	0.0	0.0	0.0	0.0	1.4	0.0
May 1974	7.0	0.0	0.0	0.7	0.0	0.0	0.0	0.0
June 1974	4.3	0.0	0.0	0.7	0.0	0.0	0.0	0.0
July 1974	10.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
August 1974	9.0	0.0	0.0	0.0	0.0	0.0	1.7	0.0
September 1974	9.0	0.0	0.0	1.7	0.0	0.0	3.3	0.0
December 1974	17.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
January 1975	2.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
February 1975	2.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
March 1975	4.0	1.3	0.0	0.0	0.0	0.0	0.0	0.0
April 1975	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
May 1975	5.3	0.0	0.0	0.0	0.0	0.0	0.3	0.0
June 1975	7.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
July 1975	19.0	0.3	0.0	0.3	0.0	0.0	0.0	0.0
August 1975	7.7	0.7	1.3	0.0	0.0	0.0	0.0	0.0
September 1975	12.7	2.0	0.0	0.7	0.0	0.7	0.0	0.0
November 1975	9.3	0.0	0.0	0.7	0.0	0.0	0.0	0.0
December 1975	3.7	1.3	0.0	0.0	0.0	0.0	0.0	0.0
January 1976	1.7	2.7	0.0	0.0	0.3	0.0	0.0	0.0
February 1976	1.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0
March 1976	3.0	2.0	0.0	0.0	0.0	0.0	0.0	0.0
April 1976	1.7	1.7	0.0	0.0	0.0	0.0	0.0	0.0

. . . continued

TABLE 18 (CONTINUED)

	<i>Clethrionomys gapperi</i>	<i>Microtus pennsylvanicus</i>	<i>Peromyscus maniculatus</i>	<i>Sorex cinereus</i>	<i>S. arcticus</i>	<i>Synaptomys borealis</i>	<i>Blarina brevicauda</i>	<i>Zapus hudsonius</i>
Alder ridge ecotone								
September 1974	5.3	0.3	1.3	0.3	0.0	0.0	1.3	0.0
December 1974	6.7	0.0	0.0	1.3	0.0	0.0	0.0	0.0
February 1975	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
April 1975	2.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
May 1975	4.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
June 1975	11.0	0.3	0.0	0.0	0.0	0.3	0.0	0.3
July 1975	8.3	1.3	0.0	0.0	0.0	0.0	0.0	0.0
August 1975	12.0	3.3	0.7	0.3	0.0	0.0	0.0	0.0
September 1975	6.3	0.7	0.0	0.3	0.0	0.0	0.3	0.0
October 1975	25.3	1.3	0.0	0.0	0.0	0.0	0.0	0.0
November 1975	13.3	2.2	0.0	0.0	0.0	0.0	0.0	0.0
December 1975	5.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0
January 1976	0.0	2.2	0.0	0.0	0.0	0.0	0.0	0.0
February 1976	2.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
April 1976	5.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Aspen upland								
July 1973	2.1	0.1	2.7	0.0	0.0	0.0	0.1	0.0
July 1974	3.7	0.0	2.7	0.0	0.0	0.0	0.0	0.0
August 1974	3.3	0.0	1.7	0.0	0.0	0.0	1.7	0.0
September 1974	1.7	0.0	1.7	0.3	0.0	0.0	2.0	0.0
December 1974	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
January 1975	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
May 1975	0.0	0.0	2.0	0.0	0.0	0.0	0.0	0.0
June 1975	0.0	0.0	1.7	0.0	0.0	0.0	0.0	0.0
July 1975	4.7	0.0	2.0	0.0	0.0	0.0	0.0	0.0
August 1975	7.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0
October 1975	20.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
November 1975	16.1	2.8	0.0	1.9	0.0	0.0	0.0	0.0
December 1975	11.4	2.8	0.0	0.0	0.0	0.0	0.0	0.0
January 1976	3.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0
February 1976	5.6	2.8	0.0	0.0	0.0	0.0	0.0	0.0
March 1976	5.6	7.6	0.0	0.0	0.0	0.0	0.0	0.0
April	2.8	3.8	0.0	0.0	0.0	0.0	0.0	0.0

captured rarely in the alder ridge ecotone and black spruce bog.

There is a significant decrease ($p < 0.05$) in the numbers of captures after the fall critical period (onset of the hiemal threshold) in some plots both years and in only one winter in others (Table 19). The aspen upland and black spruce bog showed no significant change in the number of captures in 1974 to 1975, there being few captures during this period in these two plots. The alder tamarack bog and alder ridge ecotone showed a significant decrease ($p < 0.01$) in numbers of captures before and after the hiemal threshold. In the winter of 1975 to 1976 all plots, except the aspen upland ($p < 0.05$), showed a significant decrease ($p < 0.01$) in numbers of captures during the fall critical period.

MOVEMENT OF SMALL MAMMALS IN RELATION TO CO_2 ACCUMULATION

Figures 8 to 10 show the distribution of small mammal captures in the aspen upland, alder ridge ecotone and the alder tamarack bog. The microtopography of the aspen upland is varied with areas of rock, thick soil (> 25 cm) and shallow soil (< 25 cm). The thick soils, particularly from A5-A8 to D5-D8, were associated with the greatest CO_2 accumulation. Before the CO_2 concentrations started to increase in January there was a significant number of captures in that area whereas after accumulation there were no captures and those on the rock, shallow and remainder of the deep soil (zones of lesser CO_2) were relatively increased. Once the snow melted and subnivean air mixed with environmental air, a marked animal that had frequented that area, before increased CO_2 concentrations, was recaptured there.






TABLE 19

Significance of change in number of small mammals' trap captures and trap failures before (included captures in the months before in which the hiemal threshold was broken) and after hiemal threshold (included all months when the hiemal threshold was established)

habitat and period	No. captures		No. failures		Significance	χ^2	df
	Before hiemal threshold		After hiemal threshold				
Aspen upland							
1974-75	4	146	1	224	p=0	3.4	1
1975-76	57	333	27	288	p<0.05 (almost p<0.01)	6.0	1
Alder tamarack bog							
1974-75	17	133	8	217	p<0.01	8.7	1
1975-76	54	171	31	194	p<0.01	7.7	1
Alder ridge ecotone							
1974-75	7	143	1	224	p<0.01	7.7	1
1975-76	44	301	4	266	p<0.01	32.1	1
Black spruce bog							
1974-75	6	144	4	221	p=0	1.7	1
1975-76	26	244	4	266	p<0.01	17.1	1


FIGURE 8

DISTRIBUTION OF SMALL MAMMAL CAPTURES IN THE ASPEN UPLAND
FROM OCTOBER 1975 TO APRIL 1976.


-  rock
-  shallow soil (25 cm)
-  deep soil (25 cm)
-  capture
-  trap site

Concentric circles = number of captures per site







OCTOBER



NOVEMBER




DECEMBER




JANUARY


HIEMAL THRESHOLD ESTABLISHED



FEB.



MARCH



APRIL

HIEMAL THRESHOLD BROKEN

FIGURE 9.

DISTRIBUTION OF SMALL MAMMAL CAPTURES IN THE ALDER RIDGE ECOTONE FROM OCTOBER 1975 TO MAY 1976. (THE MONTH OF MARCH 1976 WAS OMITTED FROM THE FIGURE BECAUSE NO CAPTURES WERE MADE IN THAT MONTH).



□ rock ridge

▨ bog


○ small mammal capture

● trap site


Concentric circles = number of captures per site




DECE




OCTOBER




NOVEMBER




DECEMBER



JANUARY



FEBUARY



APRIL


FIGURE 10.

DISTRIBUTION OF SMALL MAMMAL CAPTURES IN THE ALDER TAMARACK BOG
FROM NOVEMBER 1975 TO APRIL 1976.

○ small mammal capture


● trap site

Concentric circles = number of captures per site




NOVEMBER

FALL CRITICAL PERIOD




DECEMBER


HIEMAL THRESHOLD ESTABLISHED




JANUARY



FEBRUARY



MARCH



APRIL

HIEMAL THRESHOLD BROKEN

In the aspen plot four animals were caught quite frequently from October to January or later. Clethrionomys #530 was a young female caught first in August 1975. During August and November this animal was found in the area of the plot which later would have the greatest CO₂ concentration. In October it was captured in areas with thin soil. In February and March it was captured only in one area with thin soil at the edge of a bare rock. The area with thick soil and greater CO₂ concentrations was part of its home range in the late autumn prior to the fall critical period but not once CO₂ had started to accumulate.

Clethrionomys #9 was a young female first captured in October 1975. She was eventually captured 12 times from October to December in the area where an unusually great CO₂ concentration would eventually be measured. In January when CO₂ started to accumulate in the areas of thick soil this vole was captured on the rock ridge where no CO₂ accumulation occurred. Unfortunately she died in the trap.

Clethrionomys #41 was a female captured first in November. She was caught nine times in November and December in the area which later showed the greatest CO₂ accumulation but in January and March she was captured a total of three times only on rock ridges. In April she was captured once on a ridge and once back in the area where she had been captured originally nine times before CO₂ had started to accumulate.

Clethrionomys #1 was a female first captured in October 1975. In October and November she was captured four times in the area where CO₂ would accumulate to the greatest extent. In December she was captured once in the area with thin soil and in January on a rock ridge.

These four voles avoided the area where CO₂ was to accumulate to its maximum extent as soon as it started to do so. One vole was found dead, two others disappeared entirely from the plot and the one that survived through the spring critical period returned to the area that had once held the greatest amount of CO₂. The area of greatest CO₂ concentration was part of their home range but was utilized only when CO₂ was low.

The alder ridge ecotone was generally divided into two areas, ridge and bog (Figure 9). CO₂ did accumulate to some extent in 1975 to 1976. The number of small mammals captured after the fall critical period markedly decreased (November: 7 animals, December: 5 animals; critical period; January and February: 1 animal each; March: 0). In April the number of voles captured increased to four, one of which, Clethrionomys #508, a juvenile male who was first captured in October, was recaptured in the same area it had vacated after the November trapping period. It is concluded that the alder ridge was not a good overwintering habitat because only one of the animals regularly captured was there during the winter. Clethrionomys #0 was a young female who appeared in February after not being captured

since October. The other vole was captured only in January (Microtus #40-0 juvenile male). Two of the Clethrionomys that left the ridge in November or December had been captured in the bog as well and after the critical period were captured only there. The other voles may have moved into a different part of the adjacent bog that was not part of the study area.

The effects of CO_2 , habitat and the hiemal threshold were significant ($p < 0.01$) in relation to small mammal captures in the alder ecotone plot (Figure 9). Prior to the hiemal threshold (during the fall critical period) small mammal captures were high in this area. The adjacent bog plot reached the hiemal threshold before the ridge and small mammals were captured there more often than in the ecotone. In January there were two captures in the ecotone area that later had more CO_2 but these captures occurred prior to the increased CO_2 measurements. In February the same area still had high concentrations of CO_2 which small mammals avoided. The two captures recorded were on the ridge areas with reduced amounts of CO_2 . Most of the greater concentrations of CO_2 occurred in March and no small mammals were captured then. The small mammals may have moved into the adjacent bog area with relatively lower CO_2 . There was a corresponding increase in numbers from February to March in the alder tamarack bog (5 to 12 captures). In April the snowcover disappeared and small mammals were captured again in the alder ecotone plot.

The alder tamarack bog never showed a significant CO_2

accumulation, nor marked microtopographical variation, and small mammals exhibited no distinct change of areas occupied in the habitat. (Table 3, 20 and 21, Figure 10). Prior to the onset of the hiemal threshold (November to December) there was a decrease in numbers of captures probably due to the large temperature fluctuations of the fall critical period (Pruitt 1957 ;Fuller et al. 1969).

The black spruce bog has no distinct microtopographic differences but in 1975 to 1976 there were sufficient captures to show a significant effect ($p < 0.01$) of the fall critical period. Unfortunately there were insufficient small mammal captures after the hiemal threshold was reached to show the exact effect of CO_2 . Once the hiemal threshold was reached small mammal captures were very low (0 to 2.2 captures/100 trap nights). In March, during the period of greatest CO_2 accumulation, there were no captures of small mammals in this area. When the snowcover melted in April captures increased to higher numbers (3.3 captures/100 trap night) than those seen after the hiemal threshold was reached (2.2 captures/100 trap nights).

STATISTICAL ANALYSIS OF CHANGES IN SMALL MAMMAL DISTRIBUTION

In the alder tamarack bog a significant decrease ($p < 0.01$) was seen in the total numbers of small mammals captured before and after the hiemal threshold was established in both 1974 to 1975 and 1975 to

TABLE 20

Significance of number of captures in 3 small mammal trapping lines
in alder tamarack bog before and after hiemal threshold

Date	1			2			3		
	o*	e*	x ²	o*	e*	x ²	o*	e*	x ²
December 1974	1	4.8	3.0083	7	4.8	0.300	5	2.4	2.8167
January 1975	0	0.8	0.8000	2	0.8	1.800	0	0.4	0.4000
February 1975	1	0.8	0.0500	1	0.8	0.050	0	0.4	0.4000
March 1975	2	1.6	0.1000	1	1.6	0.225	1	0.8	0.0500
April 1975	0	1.6	1.6000	2	1.6	0.100	2	0.8	1.8000
Totals			5.5583			2.475			5.4667

df = 8 (non-significant)

1 = lines A - J #2-3, 2 = lines A - J #6-7, 3 = line A - J #10

o* = observed captures

e* = expected captures

TABLE 21

Significance of number of captures in 4 lines of small mammal traps
in alder tamarack bog before and after hiemal threshold

Date	1			2			3			4		
	o	e	x^2	o	e	x^2	o	e	x^2	o	e	x^2
November 1975	14	12.0	0.3333	9	12.0	0.7500	7	6.0	0.1667			
December 1975	8	4.3	3.1837	4	4.3	0.0209	2	2.1	0.0048	1	4.3	2.5325
January 1976	6	3.7	1.4297	2	3.7	0.7811	3	1.9	0.6368	2	3.7	0.7811
February 1976	0	1.4	1.4000	1	1.4	0.1143	2	0.7	2.4142	2	1.4	0.2571
March 1976	8	3.7	4.9973	3	3.7	0.1324	1	1.9	0.4263	1	3.7	1.9703
April 1976	1	2.9	1.2448	6	2.9	3.3138	2	1.4	0.2571	1	2.9	1.2448
Totals			12.5888			5.1125			3.9059			6.7858

df = 12 (p = 0), 1 = lines A - J #2-3, 2 = lines A - J #6-7, 3 = line A - J #10, 4 = lines A - J #4-5

1976 (Table 19). There was no marked microtopographical or vegetation zones in this plot so to test for change in distribution of small mammals each trapping line (A - J) was designated as a different area.

No significant change was observed in small mammal distribution either from December 1974 to April 1975 or November 1975 to April 1976 (Tables 20 and 21).

The black spruce bog exhibited a CO₂ accumulation at all five sampling stations as well as a significant decrease ($p < 0.01$) in the numbers of small mammals captured after the hiemal threshold in 1975 to 1976 (Tables 22 and 23). There were insufficient captures in 1974 to 1975 to allow statistical testing (no zeros can occur in marginal cells) and specific areas of CO₂ accumulation or lack of it could not be defined due to the homogeneity of the habitat.

Table 22 shows the setup of the test with the observed values and totals for 1975 to 1976. Table 23 shows the test itself using χ^2 . To get the expected values in this case:

$$e_{\text{cell 111}} = \frac{A_1 \cdot B_1 \cdot C_1}{h^2} = \frac{30 \cdot 540 \cdot 80}{(1,620)^2} = 5$$

This can be calculated for any number of models that are constructed. Model 1 (1 + A + B + C) of total independence for the black spruce bog was disproved. Model 2 (1 + A + B + C + AB) explaining the distribution due to the effect of the lines accounted for an insignificant amount

TABLE 22

Fienberg's contingency table test for black spruce bog 1975-76: observed values with A_1 capture of A_2 failure, B = number of the line (i.e., B_1 = line 1, B_2 = line 2, etc.) and C_1 = before and C_2 = after the hiemal threshold. A^+ , B^+ or C^+ = combined values of A &/or B &/or C

	C_1	C_2	C^+
$A_1 B_1$	11 observed value 111 cell number	1 112	12
$A_1 B_2$	8 121	2 122	10
$A_1 B_3$	7 131	1 132	8
$A_2 B_1$	259 211	269 212	528
$A_2 B_2$	262 221	268 222	530
$A_2 B_3$	263 231	269 232	532
$A_1 B^+$	26	4	30
$A_2 B^+$	784	806	1,590
$A^+ B_1$	270	270	540
$A^+ B_2$	270	270	540
$A^+ B_3$	270	270	540
$A^+ B^+$	810	810	1,620

TABLE 23

Fienberg's contingency table test for black spruce bog 1975-76: χ^2 values

Cell	Observed no. of captures	Model 1*		Model 2**		Model 3***	
		Expected no. of captures	χ^2	Expected no. of captures	χ^2	Expected no. of captures	χ^2
111	11	5	7.2000	6	4.1667	10.4	0.0346
112	1	5	3.2000	6	4.1667	1.6	0.2250
121	8	5	1.8000	5	1.8000	8.7	0.0563
122	2	5	1.8000	5	1.8000	1.3	0.3769
131	7	5	0.8000	4	2.2500	6.9	0.0014
132	1	5	3.2000	4	2.2500	1.1	0.0091
211	259	265	0.1358	264	0.0947	260.3	0.0065
212	269	265	0.0604	264	0.0947	267.7	0.0063
221	262	265	0.0340	265	0.0340	261.3	0.0019
222	268	265	0.0340	265	0.0340	268.7	0.0018
231	263	265	0.0151	266	0.0338	262.3	0.0003
232	269	265	0.0604	266	0.0338	269.7	0.0018
			18.3397 (df = 7)	16.0365 (df = 1)		0.7219 (df = 1)	
			$\chi^2 = 1.5813$ (df = 2, p = 0)		$\chi^2 = 16.0365$ (df = 1, p = 0.01)		

*Model 1: 1 + A + B + C
 **Model 2: 1 + A + B + C + AB
 ***Model 3: 1 + A + B + C + AB + AC

of the variation ($\chi^2 = 1.5813$, $df = 2$). Model 3 (I + A + B + C + AB + AC) explaining the variation in distribution due to the onset of the hiemal threshold accounted for a significant portion ($p < 0.01$) of the variation ($\chi^2 = 16.0364$, $df = 1$) with a non-significant remainder. This showed that the critical period was very significant in determining small mammal distribution. It can be seen from Table 18 that almost all the small mammals disappeared from the plot at that time. Thus the distribution or lack of small mammal captures after the hiemal threshold was established was due to the fall critical period.

If it had been possible to get more accurate and greater numbers of small mammal captures along the lines then a test of the effect of CO_2 on small mammals' distribution would have been possible.

The alder ridge ecotone showed a significant effect ($p < 0.01$) of the fall critical period on numbers of captures in both 1974 to 1975 and 1975 to 1976. It also had an accumulation of CO_2 at certain times (Table 6) and a marked microtopographical division into two areas. Given these factors a test for the effect of CO_2 on distributions of small mammals could be set up for 1975 to 1976. There were insufficient captures in 1974 to 1975. Fienberg's (1970) test, as explained for the black spruce bog, was used. Greater CO_2 concentrations are defined as readings greater than ambient levels more than 33.3 % of the time in the entire area of the habitat being studied. Table 24 gives the observed values and Table 25 shows the test. Model 2 showing the effect of CO_2 on habitat accounted for a

TABLE 24

Fienberg's contingency table test for alder ridge ecotone 1975-76:
observed values

	C ₁	C ₂	C ⁺
A ₁ B ₁	2 111	Number in that cell Cell Number 112	0 2
A ₁ B ₂	6 121	6 122	12
A ₂ B ₁	70 211	66 212	136
A ₂ B ₂	42 221	192 222	234
A ₁ B ⁺	8	6	14
A ₂ B ⁺	112	258	370
A ⁺ B ₁	72	66	138
A ⁺ B ₂	48	198	246
A ⁺ B ⁺	120	264	384

A₁ = trap captures, A₂ = Trap failure, B₁ = CO₂ > .02%/volume, B₂ = CO₂ ≤ .02%/volume, C = habitats (C₁ bog, C₂ rock), A⁺, B⁺, C⁺ - combined values of letter involved.

TABLE 25

Fienberg's contingency table test for alder ridge ecotone 1975-76: χ^2 values

Cell	Observed no. of captures	Expected no. of captures		Expected no. of captures		Expected no. of captures		Expected no. of captures	
		Model 1*	χ^2	Model 2**	χ^2	Model 3***	χ^2	Model 4****	χ^2
111	2	1.6	0.1000	2.6	0.1385	4.8	1.6333	1.6	0.1000
112	0	3.5	3.5000	2.4	2.4000	1.5	1.5000	0.4	0.4000
121	6	2.8	3.6571	1.8	9.8000	3.2	2.4500	6.4	0.0250
122	6	6.2	0.0065	7.2	0.2000	4.5	0.5000	5.6	0.0286
211	70	41.6	19.3885	69.4	0.0052	67.2	0.1167	70.4	0.0023
212	66	91.4	7.0586	63.6	0.0906	64.5	0.0349	65.6	0.0024
221	42	74.1	13.9057	46.3	0.3994	44.8	0.1750	41.6	0.0038
222	192	163.0	5.1595	190.8	0.0170	193.5	0.0116	192.4	0.0008
		52.7759 (df = 4)		13.0502 (df = 3)		6.4215 (df = 2)		0.5629 (df = 1)	
		$\chi^2 = 39.7252$ (df = 1, p = 0.01)		$\chi^2 = 6.6292$ (df = 1, p = 0.05)		$\chi^2 = 5.8585$ (df = 1, p = 0.05)			

- *Model 1: 1 + A + B + C
 **Model 2: 1 + A + B + C + BC
 ***Model 3: 1 + A + B + C + BC + AC
 ****Model 4: 1 + A + B + C + BC + AC + AB

significant amount ($p < 0.01$) of the variation ($x^2 = 39.7$, $df = 1$).

That is, CO_2 was closely related to habitat. Model 3 ($1 + A + B + C + BC + AC$) showing the effect of CO_2 on captures again accounted for a significant portion ($p < 0.01$) of the variation ($x^2 = 6.6$, $df = 1$).

When the effect of habitat on captures was included (Model 4: $1 + A + B + C + BC + AC + AB$) it accounted for a significant part ($p < 0.05$) of the variation ($x^2 = 5.9$, $df = 1$); an insignificant remainder of variation was left. CO_2 and habitat were related and both has significant effects ($p < 0.01$) on numbers of captures.

In the aspen upland the fall critical period had no significant effect on numbers of captures in 1974 to 1975 but did ($p < 0.05$) in 1975 to 1976. CO_2 accumulation occurred in both winters (Tables 3 and 4) in this plot and a microtopographical variation was obvious. This variation resulted in three distinct types of areas: rock, shallow soils and deep soils. There were not enough captures in 1974 to 1975 to allow statistical analysis. For the 1975 to 1976 data Fienberg's (1970) test was used (Table 26 and 27). Model 1 ($1 + A + B + C$) of complete independence was disproved. Model 2 ($1 + A + B + C + BC$) shows the relationship of CO_2 with habitat which accounted for a significant amount of the variation ($x^2 = 10.6$, $df = 3$). Model 3 ($1 + A + B + C + BC + AC$) showing the effect of CO_2 on captures again accounted for a significant portion of the variation ($x^2 = 7.5$, $df = 2$). When the effect of habitat on captures was also included (Model 4: $1 + A + B + C + BC + AC + AB$) it accounted for a significant part of the variation ($x^2 = 5.8$, $df = 1$). CO_2 levels and habitat was related

TABLE 26

Fienberg's contingency table test for aspen upland 1975-76:
observed values

	C_1	C_2	C_3	C^+
$A_1 B_1$	9 111	8 112	6 113	23
$A_1 B_2$	13 121	1 122	11 123	25
$A_2 B_1$	94 211	58 212	15 213	167
$A_2 B_2$	140 221	98 222	73 223	311
$A_1 B^+$	22	9	17	48
$A_2 B^+$	234	156	88	478
$A^+ B_1$	103	66	21	190
$A^+ B_2$	153	99	84	336
$A^+ B^+$	256	165	105	526

A_1 = trap captures, A_2 = trap failures, B = CO_2 (B_1 high, B_2 low),
 C_1 = habitat (C_1 deep, C_2 shallow, C_3 rock), A^+ , B^+ , C^+ - total
numbers in each of those letters.

TABLE 27

Fienberg's contingency table test for aspen upland 1975-76: χ^2 values

Cell	Observed no. of captures	Model 1*		Model 2**		Model 3***		Model 4****	
		Expected no. of captures	χ^2	Expected no. of captures	χ^2	Expected no. of captures	χ^2	Expected no. of captures	χ^2
111	9	8.4	0.0429	9.4	0.0170	8.9	0.0011	12.5	0.9800
112	8	5.4	1.2519	6.0	0.6667	3.6	5.3778	5.1	1.6490
113	6	3.4	1.9882	1.9	8.8474	3.4	1.9882	5.3	0.0924
121	13	14.9	0.2423	14.0	0.0714	13.1	0.0008	9.5	1.2895
122	1	9.6	7.7042	9.0	7.1111	5.4	3.5852	3.9	2.1564
123	11	6.1	3.9361	7.7	1.4143	13.6	0.4971	11.7	0.0418
211	94	83.7	1.2675	93.6	0.0017	94.1	0.0001	90.5	0.1354
212	58	54.0	0.2963	60.0	0.0667	62.4	0.3103	60.9	0.1381
213	15	34.3	10.8598	19.1	0.8801	17.6	0.3841	15.7	0.0312
221	140	148.0	0.4324	139.0	0.0072	139.9	0.0001	143.5	0.0854
222	98	95.4	0.0709	90.0	0.7111	93.6	0.2068	95.1	0.0884
223	73	60.7	2.4924	76.3	0.1427	70.4	0.0960	72.3	0.0068
		30.5849	(df = 8)	19.9374	(df = 5)	12.4476	(df = 3)	6.6944	(df = 2)
		$\chi^2 = 10.6474$ (df = 3, p = 0.05)		$\chi^2 = 7.4898$ (df = 2, p = 0.05)		$\chi^2 = 5.7810$ (df = 1, p = 0.05)			

*Model 1: 1 + A + B + C

**Model 2: 1 + A + B + C + BC

***Model 3: 1 + A + B + C + BC + AC

****Model 4: 1 + A + B + C + BC + AC + AB

with both having significant effects on the numbers of captures but some other factor causing the significant remainder of unexplained variation ($\chi^2 = 6.7$, $df = 2$) was also present. This could have been due to "trap addict" animals, predators, nonfunctioning traps or any number of unknowns.

DISCUSSION

SECTION 1.

CO₂ ACCUMULATION

In this study, I found a consistent variation between habitats in the accumulation of subnivean CO₂. This could explain why some authors found CO₂ accumulated (Bashenina 1956, Kelley et al 1968, and others) while others did not (Fuller and Holmes 1972, Reiners 1968, and Coyne and Kelley 1971). The CO₂ concentrations measured in this study agreed with those found by other authors (Coyne and Kelley 1971, Kelley et al. 1968, Bashenina 1956). Havas and Mäenpää (1972) recorded accumulations of three to four times the atmospheric levels in their area (0.028%/volume CO₂) while I measured maximum concentrations up to four or five times the atmospheric levels that I recorded (0.02 %/volume) at the Taiga Biological Station and the Experimental Lakes Area near Kenora, Ontario. The reduced amounts of atmospheric CO₂ in my study may be due to the large amount of continually green vegetation (conifers) available for photosynthesis above the snow and to the lack of air pollution in the area. The atmospheric levels recorded by Havas and Mäenpää (1972) are less than those of other authors from areas without evergreens (0.035%/volume CO₂, Kelley et al 1968).

I found that CO₂ concentrations in some areas such as the aspen

upland and alder ridge ecotone showed sustained accumulations until the spring critical period and ambient mixing. In contrast CO_2 concentrations in the alder tamarack bog and jackpine ridge sometimes increased above ambient and then usually returned quickly to ambient concentrations. Reiners (1968), Coyne and Kelley (1971) and Kelley et al (1968) found CO_2 concentrations decreased to ambient levels or less once the hiemal threshold had been reached but they studied a few habitats intensively and thus may not have encountered habitats with sustained accumulations.

Kelley et al (1968) found subnivean CO_2 levels to be most consistent from 1000 to 1600 hours. I found that subnivean concentrations of CO_2 changed little over a number of days and changed very little throughout the daylight hours in an area with ambient levels of subnivean CO_2 . However, as there may have been diel variation at sites with accumulating CO_2 , all measurements were taken between 1000 and 1600 hours.

The CO_2 concentrations that I found to be much lower than ambient may have been due to moss growth (Hagerup in Kalela 1962) or other plant photosynthesis that can occur at temperatures as low as 0°C . (Havas and Mäenpää 1972). This, with the light available under a snowcover of 15 cm. (Evernden 1966), could allow photosynthesis to use up subnivean CO_2 and create the reduced CO_2 concentrations. The fact that these reduced CO_2 levels occurred only occasionally after there was thicker snow, supports this conclusion.

SECTION 2.

FACTORS INFLUENCING CO₂ ACCUMULATION

The consistency of accumulation or lack of accumulation of CO₂ in five of six plots suggests that CO₂ accumulation is habitat or site-dependent. The site-dependent factors measured in this study do not uphold that assumption because neither soil depth, microtopographic change nor the amount of green vegetation contributed significantly to the predicted value of CO₂ concentrations. Generally, areas with deeper, well-developed soils (> 25 cm., aspen upland) exhibited the greatest CO₂ accumulations and greater concentrations occurred in the top 10 cm. of soil than occurred in the subnivean space. Areas with very shallow soils (< 10 cm., jackpine ridge) did not have this mass for root respiration and thus did not accumulate CO₂. Areas without green vegetation or soil such as bare rock in the aspen upland never accumulated CO₂ since there was no biological source to produce it. But, areas with little green vegetation did accumulate CO₂ so plant respiration is not necessarily a major producer of subnivean CO₂.

General site-dependent factors such as moisture may be of greater importance than I initially thought. Wet, but not saturated organic soils produce more CO₂ than drier ones (Polyakova 1970, Coyne and Kelley 1971). This was true in the alder ridge ecotone and aspen upland. The saturated peat of the alder tamarack bog did not

accumulate CO_2 because there was little air space available for gas exchange and thus there was a decrease in respiration. The black spruce bog was drier than the alder tamarack bog; this might have accounted for its greater CO_2 concentrations.

I agreed with Van der Drift and Witkamp, (in Witkamp 1963) that pH had little effect on decomposition rates, rather than the decreased rates found by Flanagan and Scarborough (1972) when pH was outside the range 5 to 7.5. Three habitats had pH values less than 5 and two of these accumulated subnivean CO_2 .

The presence of wind is rarely felt on the floor of the boreal forest, so that the effect of wind stated by Kelley et al (1968) was not evident.

The age and depth of litter mentioned by Reiners (1968) and Witkamp (1966) did not seem to influence CO_2 accumulation. The aspen upland and the alder tamarack bog had the greatest litter fall of the six habitats (deciduous versus primarily coniferous) but one had the greatest accumulation of CO_2 while the other had no accumulation.

The mean subnivean temperatures in the six habitats were rarely less than the critical -7°C . that was correlated with the cessation of CO_2 production from decomposition (Flanagan and Scarborough 1972). The aspen upland had the warmest subnivean temperatures and the greatest CO_2 accumulation. The alder tamarack bog was warmer, but accumulated

less CO_2 , than the black spruce bog. Also, there was no strong correlation between CO_2 concentrations and subnivean temperatures while subnivean temperatures only contributed significantly to the predictive value of CO_2 concentrations once in one habitat during the two year study.

Of the environmental parameters measured in this study site-independent factors, particularly snow factors of hardness, density, thickness and number of ice layers, had the greatest effect on subnivean CO_2 accumulation in both winters. They made up over half (54%, 65%) of the significant variables contributing to the prediction of CO_2 concentrations in both winters while being only 4 of 15 parameters. The most important snowcover parameters were density and hardness. The relationship between these two parameters was not clear but the fact that they affect CO_2 accumulations is not surprising because they characterize the blanket or container that confines the subnivean CO_2 . Denser, harder and thicker layers of snow would impede the diffusion of CO_2 from the subnivean space.

The production of CO_2 on one habitat and the lack of it on another would reasonably depend upon the site-dependent variables in that habitat. Unfortunately this was not the case in the three site-dependent variables I measured in this study. The effect of all parameters was obscured because of the lag time from their onset to their effect on the CO_2 production. A controlled experiment where different factors could be removed so as to remove or control lag

effects better would allow relationships to be stronger and more obvious.

SECTION 3.

EFFECT OF CO₂ ON SMALL MAMMAL DISTRIBUTION

Number of small mammals captured increased from 1973 to 1976. This is especially evident in the last year of the study where more individuals were captured over the winter than had been captured the previous summer. Such an increase could have been the result of subnivean breeding (Fuller 1977) or increased juvenile survival. However, no markedly juvenile voles were captured.

Clethrionomys gapperi was the small mammal captured most frequently in all six habitats. Fuller (1977) also found C. gapperi to be the boreal small mammal captured most commonly. Microtus pennsylvanicus can occupy the same type of habitat as C. gapperi (Iverson and Turner 1972, Payne 1974, Riewe 1973) and was increasing in numbers towards the end of the study. The remaining small mammal species in the study area were captured rarely if ever in winter.

The significant decrease in numbers of captures before and after the hiemal threshold, in all cases where there were large enough numbers of captures to be significant, is related to the fall critical period or time between thermal overturn (when the ground temperatures

become colder than the air temperatures) and onset of the hiemal threshold (Pruitt 1957, Fuller et al 1969). It is a time of environmental stress and small mammals without home ranges, food stores, nests or those that are not in peak physical condition are subject to its adverse effects. Many small mammals do not appear to live through a second winter (Hamilton 1942, Pernetta 1976); this would be expressed as a decreased post-snowfall capture rate (Gunderson in Stebbins 1976). The fall critical period was especially important in determining survival of small mammals in all four of the habitats trapped in winter.

The aspen upland and the alder ridge ecotone exhibited the greatest diversity of microtopography, significant CO₂ accumulations and significant effects of CO₂ ($p < .01$) on small mammal distributions. The specific movements of marked small mammals away from greater CO₂ concentrations and their return once the concentrations had lessened in the aspen upland, were indicative of the effect of CO₂ accumulations.

Small mammals respond physiologically to concentrations of CO₂ as low as 1 or 2% (Galantsev and Tumanov 1969) at temperatures about 20°C. and when stressed by lower temperatures they respond to amounts less than 1 or 2% (Dawson 1955, Baudinette 1974). The values I measured in the subnivean space (up to 0.12%/volume) while not as great as these were much greater than the ambient (0.02%/volume). In an enclosed subnivean area such as a nest, the CO₂ could accumulate

to even greater levels than those I measured in the subnivean space.

The response of small mammals to increased CO_2 , causing increased respiratory function, decreased heart rate and rectal temperatures may be an adaptive response to the stress caused by increased CO_2 concentrations. These altered body functions may lead to lessened activity, decreased winter metabolic rates and related lower calorie requirements. However, these adaptive responses may themselves cause problems for the animals. The lower reaction rate could lessen the small mammals' sensory preception to temperature differences, food and predators and thus decrease their survival rate. The changes in small mammal distribution that I observed were associated with increase in CO_2 . These physiological responses to CO_2 could lead to the aforementioned changes in distribution. It would be advantageous for the small mammal to frequent areas which elicit less physiological stress (i.e., those with less CO_2).

If food shortages caused the changes in distribution shown in the aspen upland and alder ridge ecotone one would also expect marked changes in the distribution of small mammals in the alder tamarack bog. This did not happen. The standard error to mean subnivean temperatures was low in all four plots. Higher small mammal numbers are not necessarily related to available food (Gorecki and Gebczynska 1962, Chitty et al. 1968) or supplementary food (Watts 1968, Flowerdew 1972, Fairbairn 1977) or warmer subnivean temperatures (Fuller 1977). Thus these factors do not explain all the observed variations in distribution. Given the facts that subnivean CO_2 does accumulate,

small mammals do react physiologically to CO₂, areas of homogeneous habitat that have increased concentrations of CO₂ have few small mammals, it is reasonable to conclude that small mammals avoid areas of greater CO₂ concentrations. CO₂ thus affects small mammal distribution. In the case of Clethrionomys gapperi, concentration of subnivean CO₂ must be considered as an additional dimension to its ecological niche.

SUMMARY

SECTION 1.

CO₂ ACCUMULATION

CO₂ accumulated specifically and consistently in some habitats while not in others. The aspen upland, black spruce bog, jackpine sandplain and part of the alder ridge ecotone exhibited CO₂ accumulations. However, the extent and amount of accumulation varied between these habitats, being greatest in the aspen upland. CO₂ did not accumulate in the alder tamarack bog or the jackpine ridge. CO₂ accumulation was found to be similar in the same habitats in both winters.

SECTION 2.

FACTORS INFLUENCING CO₂ ACCUMULATION

The accumulation of CO₂ was affected primarily by snow characteristics. Subnivean temperature did not show a strong correlation with the predictive value of CO₂ accumulation. Neither microtopography nor amount of green vegetation prior to snowfall were significant predictors of CO₂ accumulation. The effect of weather (ambient air temperature and barometric pressure) was ambiguous. In a few cases significant correlations between either

temperature or barometric pressure and CO_2 levels were found. One would expect weather factors to have similar effects in all areas, but this was not the case.

SECTION 3.

EFFECT OF CO_2 ON SMALL MAMMAL DISTRIBUTION

The reduction in small mammal numbers was associated with the fall critical period. The absence of accumulation of CO_2 was significantly associated with the absence of change in small mammal distribution especially in the alder tamarack bog. The accumulation of CO_2 was significantly associated with the change in distribution of small mammals especially in the aspen upland.

Given the facts that subnivean CO_2 does accumulate, small mammals do react physiologically to CO_2 , areas of homogeneous habitat that have increased concentrations of CO_2 have few small mammals, it is reasonable to conclude that small mammals avoid areas of greater CO_2 concentrations and thus subnivean CO_2 affects small mammal distributions.

LITERATURE CITED

- Anderson, J.M. 1973. CO₂ evolution from two temperate, deciduous woodland soils. *J. Appl. Ecol.* 10: 361-378.
- Andrzejewski, R. and M. Mazurkiewicz. 1976. Abundance of food supply and size of the bank vole's (Clethrionomys glareolus) home range. *Acta Theriol.* 21(17): 237-253.
- Atmospheric and Environmental Services. 1970. Temperature and precipitation 1941-70: prairie provinces. Department of Environment, Canada.
- Bashenina, N.V. 1956. Influence of the quality of subnivean air on the winter hibernation of voles. *Zool. Zh.* 35(6): 940-942.
- Baudinette, R.V. 1974. Physiological correlations of burrow gas conditions in the California ground squirrel. *Comp. Biochem. Physiol.* 48A: 733-743.
- Beer, J.R. 1961. Winter home ranges of the red-backed mouse and white-footed mouse. *J. Mamm.* 42: 174-180.
- Benoit, R.E., W.B. Campbell and R.W. Harris. 1972. Decomposition of organic matter in the wet meadow tundra, Barrow, a revised word model. *Proceedings of 1972 Tundra Biome Symposium*: 111-115.
- Brown, A. and A Macfayden. 1969. Soil CO₂ output and small scale vegetation patterns in a Calluna heath. *Oikos* 20: 8-15.
- Buckner, C.H. 1966. Populations and ecological relationships of shrews in tamarack bogs of southeastern Manitoba. *J. Mamm.* 47:181-194.
- Chitty, D., D. Pimentel and C.J. Krebs. 1968. Food supply of overwintered voles. *J. Anim. Eco.* 37: 113-120.
- Coyne, P.I. and J.J. Kelley. 1971. Release of CO₂ from frozen soil to the arctic atmosphere. *Nature* 234(5329): 407-408.
- Darden, T.R. 1971. Respiratory adaptations of a fossorial mammal, the pocket gopher (Thomomys bottae). *Diss. Abst. Int.* 33B:1284.
- Dawson, W.R. 1955. The relation of oxygen consumption to temperature in desert rodents. *J. Mamm.* 36(4): 543-553.
- Evernden, L.L.N. 1966. Light alteration caused by snow and its importance to subnivean rodents. MSc. Thesis. Dept. of Zoology, Edmonton, Alberta.


- Fairbairn, D.J. 1977. The spring decline in deer mice: death or dispersal? *Can. J. Zool.* 55(1): 84-92.
- Fienberg, S.E. 1970. An analysis of multidimensional contingency tables. *Ecology* 51: 419-433.
- Flanagan, P.W., and A. Scarborough. 1972. Laboratory and field studies on decomposition of plant material in Alaska tundra areas. *Proceedings of the 1972 Tundra Biome Symposium*:110-111.
- Flowerdew, J.R. 1972. The effect of supplementary food on a population of wood mice (Apodemus sylvaticus). *J. Anim. Ecol.* 41: 553-566.
- Fuller, W.A. 1977. Demography of a subarctic population of Clethrionomys gapperi: numbers and survival. *Can. J. Zool.* 55(1): 42-51.
- Fuller, W.A. and J.G. Holmes. 1972. *The life of the far north.* McGraw-Hill. New York. 232p.
- Fuller, W.A., L.L. Stebbins and G.R. Dyke. 1969. Overwintering of small mammals near Great Slave Lake Northern Canada. *Arctic* 22(1): 34-55.
- Galantsev, V.P. and I.L. Tumanov. 1969. Ecological-physical characteristics of the cardiac activity of rodents. *Zool. Zh.* 48: 1066-1073.
- Geiger, R. 1965. *The climate near the ground.* Harvard Univ. Press. Cambridge, Mass. 494p.
- Gentry, J.B. and E.P. Odum. 1957. The effect of weather on the winter activity of old field rodents. *J. Mamm.* 38: 72-77.
- Gorecki, A. and Z. Gebczynska. 1962. Food conditions for small rodents in a deciduous forest. *Acta Theriol.* 10: 275-295.
- Grodzinski, W. 1963. Can food control the number of small rodents in the deciduous forest? *Proceedings of the XVI International Congress of Zoology.* vol. 1: 257.
- Hamilton, W.J.Jr. 1942. Winter reduction of small mammal populations and its probable significance. *Amer. Nat.* 76: 216-218.
- Hayward, J.S. 1965. Microclimate temperature and its adaptive significance in six geographic races of Peromyscus. *Can. J. Zool.* 43: 341-350.
- Havas, P. and E. Mäenpää, 1972. Evolution of carbon dioxide at the floor of a Hylocomium - Myrtillus type spruce forest. *Aquilo.* Ser. Bot. 11: 4-22.
- Iverson, S.L. and B.N. Turner. 1972. Winter coexistence of Clethrionomys gapperi and Microtus pennsylvanicus in a grassland habitat. *Amer. Midl. Nat.* 88(2): 440-445.

- Kalela, O. 1962. On the fluctuations in the numbers of arctic and boreal small rodents as a problem of production biology. Ann. Acad. Sci. Fenn. Ser. A, IV Biologica, 66. 38p.
- Kelley, J.J., D.F. Weaver and B.P. Smith. 1968. The variation of CO₂ under the snow in the arctic. Ecology 49(2): 358-361.
- Kelley, J.J. and D.F. Weaver. 1969. Physical processes at the surface of the arctic tundra. Arctic 22(4): 425-437.
- Kennerly, T.E. Jr. 1964. Microenvironmental conditions of the pocket gopher burrow. Tex. J. Sci. 16(4): 395-441.
- Klein, G.L., D.C. Pearce and L.W. Gold. 1950. Method of measuring the significant characteristics of a snow cover. NRC Associate Committee on Soil and Snow Mechanics. Technical Memorandum #18.
- Kosonen, M. 1969. CO₂ production in relation to temperature and plant mass. Oikos 20: 335-343.
- Kucera, C.L. and D.R. Kirkham. 1971. Soil respiration studies in tall grass prairie in Missouri. Ecology 52: 912-915.
- McNab, B.K. 1963. Bioenergetics and determination of home range size. Amer. Nat. XCVII: 133-140.
- Morris, R.D. 1969. Competitive exclusion between Microtus and Clethrionomys in the aspen parkland of Saskatchewan. J. Mamm. 50: 291-301.
- Payne, L. 1974. The comparative ecology of introduced Clethrionomys gapperi proteus (Bangs) and native Microtus pennsylvanicus terranovae (Bangs) on Camel Island Notre Dame Bay, Newfoundland. Master's Thesis. Dept. of Zoology, Univ. of Man. Winnipeg, Manitoba.
- Pernetta, J.C. 1976. Diets of the shrews Sorex araneus L. and S. minutus L. in Wytham grassland. J. Anim. Ecol. 45: 899-912.
- Polyakova, O.Y. 1970. Contents of O₂ in soil H₂O and CO₂ in soil air in forest bogs of central Ukrainian Polesie. Visn. Sil' Skohospod Nauk translated from Ref. Zh. Otd. Vyp. Pochvoved Agrokhim 1970 No. 10.57.80.
- Pruitt, W.O. Jr. 1957. Observations on the bioclimate of some taiga mammals. Arctic 10: 131-139.
- 1959a. Microclimates and local distributions of small mammals on the George Reserve, Michigan, Misc. Publ. Mus. Zoo. Univ. Mich. #109. 27p.
- 1959b. A method of live trapping small taiga mammals in winter. J. Mamm. 40(1): 139-143.

- 1960. Animals in the snow. *Sci. Amer.* 202(1): 60-68.
- Pruitt, W.O. Jr., C.V. Lucier and L.L. Huffman. 1961. Small mammal bioclimate studies in northern regions. Alaskan Air Command Arctic Aeromedical Lab. Ft. Wainwright Alaska Technical Report 60-34 Project 8237.
- Reiners, W.A. 1968. CO₂ evolution from the floor of three Minnesota forests. *Ecology* 49: 471-483.
- Riewe, R. 1973. Food habits of insular meadow voles, Microtus pennsylvanicus terranovae, (Rodentia: Cricetidae) in Notre Dame Bay, Newfoundland. *Can Field Nat.* 87: 5-13.
- Smith, R.L. 1966. Ecology and field biology. Harper and Row, New York. 686p.
- Soholt, L.F., M.K. Yousef and D.B. Dill. 1973. Responses of Merriam's kangaroo rat Dipodomys merriami to various levels of CO₂ concentrations. *Comp. Biochem. Physiol.* 45A: 455-462.
- Stebbins, L.L. 1976. Overwintering of the red-backed vole at Edmonton, Alberta. *J. Mamm.* 57(13): 554-561.
- Stickel, L.F. 1960. Peromyscus ranges at high and low population densities. *J. Mamm.* 41: 433-441.
- Studier, E.H. and J.W. Proctor. 1971. Respiratory gases in burrows of Spermophilus tridecemlineatus. *J. Mamm.* 52:631-633.
- Watts, C.H.S. 1968. The foods eaten by wood mice (Apodemus sylvaticus) and bank voles (Clethrionomys glareolus) in Wytham Woods Berkshire. *J. Anim. Ecol.* 37: 25-41.
- West, S.D. 1977. Midwinter aggregations in the northern red-backed vole, Clethrionomys rutilus. *Can. J. Zool.* 55(9): 1404-1409.
- Williams, D.D. and R.L. Rausch. 1973. Seasonal CO₂ and O₂ concentrations in the dens of hibernating mammals (Sciuridae). *Comp. Biochem. Physiol.* 44A: 1227-1235.
- Withers, P.C. 1975. A comparison of respiratory adaptations of a semi-fossorial and surface dwelling Australian rodent. *J. Comp. Physiol.* 98: 193-203.
- Witcamp, M. 1963. Microbial populations of leaf litter in relation to environmental conditions and decomposition. *Ecology* 44:370-377.
- 1966. Rates of CO₂ evolution from the forest floor. *Ecology* 47: 492-494.
- Witcamp, M. and M.L. Frank. 1969. Evolution of CO₂ from litter, humus, and subsoil of a pine stand. *Pedobiologica* 9:358-365.

APPENDIX A

Calibration of a Dräger multigas analyzer
against a gas chromatograph on April 3, 1974
with a r value of 0.9956



APPENDIX B

Procedure for Field Operation of the Dräger Multigas Analyzer

1. Keep the analyser and measuring tubes warm (with pocket warmer in paper bags or next to the body).
2. Take cork out of tygon and remove 10 pumps (approximately 5000 cc.) of air from the tube. This draws subnivean air into the tube.
3. Take meter and tube out of warm place.
4. Break ends of tube off and insert blue arrowed end into the meter, and the white end into the tygon.
5. Hold tube and tygon in bare hand to maintain as much warmth as possible. Tubes work best at about 15° C. and this was the quickest most efficient way of approximating this temperature. Different people measured the same reading, so this method can be used by different people for the same result.
6. Once analyzer has been pumped 10 times, remove CO_2 measuring tube from pump and record reading from 0 point to point when color begins to fade. Often the reading is vague, and so the reading is read from where the color begins to fade to where it has disappeared altogether (ex. .02 - .025%/volume CO_2). The color change was much harder to read during bright sunny light than during cloudy days.