

THE EFFECTS OF LANDSCAPE RESTORATION ON GREENHOUSE GAS
EMISSIONS AND PLANT SPECIES AND ABUNDANCE

BY

MICHELLE M. ERB

A Thesis
Submitted to the Faculty of Graduate Studies
In Partial Fulfillment of the Requirements
For the Degree of

MASTER OF SCIENCE

Department of Soil Science
University of Manitoba
Winnipeg, MB

© October, 2005



Library and
Archives Canada

Bibliothèque et
Archives Canada

0-494-08846-X

Published Heritage
Branch

Direction du
Patrimoine de l'édition

395 Wellington Street
Ottawa ON K1A 0N4
Canada

395, rue Wellington
Ottawa ON K1A 0N4
Canada

Your file *Votre référence*

ISBN:

Our file *Notre référence*

ISBN:

NOTICE:

The author has granted a non-exclusive license allowing Library and Archives Canada to reproduce, publish, archive, preserve, conserve, communicate to the public by telecommunication or on the Internet, loan, distribute and sell theses worldwide, for commercial or non-commercial purposes, in microform, paper, electronic and/or any other formats.

The author retains copyright ownership and moral rights in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

AVIS:

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque et Archives Canada de reproduire, publier, archiver, sauvegarder, conserver, transmettre au public par télécommunication ou par l'Internet, prêter, distribuer et vendre des thèses partout dans le monde, à des fins commerciales ou autres, sur support microforme, papier, électronique et/ou autres formats.

L'auteur conserve la propriété du droit d'auteur et des droits moraux qui protègent cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

In compliance with the Canadian Privacy Act some supporting forms may have been removed from this thesis.

Conformément à la loi canadienne sur la protection de la vie privée, quelques formulaires secondaires ont été enlevés de cette thèse.

While these forms may be included in the document page count, their removal does not represent any loss of content from the thesis.

Bien que ces formulaires aient inclus dans la pagination, il n'y aura aucun contenu manquant.


Canada

THE UNIVERSITY OF MANITOBA
FACULTY OF GRADUATE STUDIES

COPYRIGHT PERMISSION

**The Effects of Landscape Resotoration on Greenhouse Gas Emissions and Plant
Species and Abundance**

By

Michelle M. Erb

A Thesis/Practicum submitted to the Faculty of Graduate Studies of The University of
Manitoba in partial fulfillment of the requirement of the degree
of
Master of Science

Michelle M. Erb © 2005

Permission has been granted to the Library of the University of Manitoba to lend or sell copies of this thesis/practicum, to the National Library of Canada to microfilm this thesis and to lend or sell copies of the film, and to University Microfilms Inc. to publish an abstract of this thesis/practicum.

This reproduction or copy of this thesis has been made available by authority of the copyright owner solely for the purpose of private study and research, and may only be reproduced and copied as permitted by copyright laws or with express written authorization from the copyright owner.

ABSTRACT

Erb, Michelle M. M.Sc., The University of Manitoba, October, 2005. The Effect of Landscape Restoration on Greenhouse Gas Emissions and Plant Species and Abundance. Major Professor: Dr. David Lobb.

The objective of this study was to investigate the effects of landscape restoration on two main parameters: greenhouse gas emissions (GHG; carbon dioxide, methane and nitrous oxide) and plants (species - including crop and weed species - and abundance). Two field experiments and two growth chamber experiments were completed over a two-year period to fulfill this objective. The two field experiments were similar in nature and examined the impacts of landscape restoration on both parameters under field-scale conditions. Greenhouse gases and plants were studied in separate growth chamber experiments. The growth chamber experiments complemented the field experiments but focused on one of the two parameters.

In the field, soil was removed from the lower slope riparian area of the landscape where soil had accumulated due to past tillage erosion, and was added to the eroded upper slope area. Nitrous oxide (N_2O), and methane (CH_4) emissions were not influenced by the removal of soil; however, carbon dioxide (CO_2) emissions were reduced in the first year following soil removal. In the upper slope area where soil was added, greenhouse gas emissions were not impacted. The growth chamber experiments assessed varying depths of soil removal and addition and the effects on GHG emissions. The depth of soil removal did not influence cumulative gas flux but in general, and similar to field results, soil removal reduced CO_2 emissions. Reductions in N_2O

emissions following soil removal were also found in the growth chamber. Soil addition did not impact GHG emissions by depth.

To address the question of landscape restoration effects on plants, plant emergence, abundance, and species composition was monitored in the field. In the first year following soil addition, weed emergence increased where soil was added. In one of the two study areas, crop yield was greater where soil was added despite the increase in weed emergence. The number of weed species present following soil addition remained the same as the controls. In year two following the addition of soil, weed emergence numbers where soil was added were similar to the control. The number of weed species present did increase. Because the area from which soil was removed was not part of the cropland, the type (weedy vs. native or wetland) of species revegetating the lower slope removal area was the primary interest. The species observed in the lower slope area were predominantly weeds and similar to those found in the adjacent cropland. In the growth chamber, the soil seed bank was examined to assess the viability of the seed bank within the soil profile and assess the species present; this information may be useful for predicting potential impacts of landscape restoration on weed populations in the restored cropland. It was found that the most viable seeds exist near the soil surface (within the top 5 cm of soil) and that the species were predominantly weeds. The species found in the seed bank correspond with those found in the field experiments in both the upper and lower slope areas.

In summary, this study demonstrated that in the medium-term, landscape restoration does not increase greenhouse gas emissions from soil. In fact, the removal of soil will benefit atmospheric CO₂ levels by reducing CO₂ emissions from the lower slope

removal areas. Weed emergence will likely increase in the first year following the addition of soil; which may or may not adversely affect crop yield depending on the crop type. The overall impact of increased weed pressures can be reduced by planting a competitive crop, and using the appropriate herbicide in correct rotation.

ACKNOWLEDGEMENTS

I would like to thank Ducks Unlimited Canada, Wildlife Habitat Canada and Manitoba Climate Change Action Fund for the financial support of this project.

I want to acknowledge Dr. David Lobb for being an advisor and friend who challenged and encouraged me throughout the course of this project. Your knowledge and support helped make this experience interesting, fun, and rewarding. Thank you!

I would like to thank my advisory committee: Dr. David Burton, Dr. Alan Moulin and Dr. Rene Van Acker for their input and guidance in developing this project.

Thank you to the technical support staff in the lab and in the field (Rob Ellis, Val Ward, Diane Smith, Clay Sawka, Marla Riekman, Myron Kroeker, Brad Sparling, Cedric MacLeod, Ruifang Wang, Sheng Li, Tim Stem, Christine Cantello) and to the administrative support of Barb Finkelman and Terri Ramm; and to fellow grad students for their friendship and encouragement. I would also like to acknowledge the Manitoba Zero Tillage Research Association (Bryce Wood and Lindsay Coulthard) for helping with plot establishment.

A very special thanks to my friends and family for standing by me and telling me to just get it done! Mom, Dad and Nikki, I appreciate your emotional support and encouragement throughout the past three years. Michael, you have been extremely kind, patient and supportive; I hope you know how much I truly appreciate you and how I could not have done this without you.

FOREWORD

This thesis has been prepared in the manuscript format in adherence with the guidelines established by the Department of Soil Science at the University of Manitoba. The Canadian Journal of Soil Science was the reference style used in this document. Chapters 3 and 4 will be submitted to Journal of Soil and Water Conservation. For all papers, I will be the lead author and co-authorship will be designated accordingly.

TABLE OF CONTENTS

ABSTRACT.....	ii
ACKNOWLEDGEMENTS.....	v
FOREWORD.....	vi
TABLE OF CONTENTS.....	vi
LIST OF TABLES.....	x
LIST OF FIGURES.....	xi
1. INTRODUCTION.....	13
2. LITERATURE REVIEW.....	15
2.1 Soil Erosion in Topographically Complex Landscapes.....	15
2.1.1 Water and Wind Erosion.....	15
2.1.2 Tillage Erosion.....	16
2.2 Effects of Soil Loss on Crop Productivity.....	17
2.3 Current Methods for Improving Soil Quality and Productivity.....	19
2.3.1 Fertilizer Use.....	19
2.3.2 Use of Organic Amendments.....	20
2.3.3 Cropping System.....	22
2.3.4 Tillage System.....	23
2.4 Soil Addition/Landscape Restoration.....	23
2.4.1 Soil Addition Effects on Greenhouse Gas Emissions.....	26
2.4.2 The Effects of Topsoil Addition on Weeds.....	27
3. IMPACTS OF LANDSCAPE RESTORATION ON GREENHOUSE GAS EMISSIONS.....	28
3.1 Abstract.....	28
3.2 Introduction.....	28
3.3 Materials and Methods.....	30
3.3.1 Site Selection and Description.....	31
3.3.2 Field Experiments.....	34
3.3.2.1 Field Experiment 1 (FE1 2003-2004).....	34
3.3.2.2 Field Experiment 2 (FE2).....	38
3.3.2.5 Gas Flux Measurement and Analysis.....	39
3.3.2.6 Soil Sampling and Analysis.....	40
3.3.2.7 Rainfall.....	41
3.3.3 Soil Column Experiment.....	41
3.3.3.1 Part 1 – Addition Experiment.....	41

3.3.3.2 Part 2 – Removal Experiment	42
3.3.4 Data Analyses	43
3.3.4.1 Field Data	43
3.3.4.2 Column Experiment Data	44
3.4 Results	44
3.4.1. Soil Environment	44
3.4.1.1 Soil Temperature and Moisture	45
3.4.1.2 Soil Nutrients	50
3.4.1.3 Microbial Biomass Carbon	55
3.4.2 Gas Flux	58
3.4.2.1 Carbon Dioxide	58
3.4.2.2 Methane	60
3.4.2.3 Nitrous Oxide	61
3.4.3 Relationship between Flux and Depth of Added Soil and Depth of Removed Soil	71
3.4.3.1 Column Experiment Part 1	71
3.4.3.2 Column Experiment Part 2	72
3.5 Discussion	76
3.5.1 The Effects of Landscape Restoration on the Soil Environment	76
3.5.1.1 Soil Temperature and Moisture	76
3.5.1.2 Soil Nutrients	78
3.5.1.3 Microbial Biomass Carbon	79
3.5.2 The Effects of Landscape Restoration on Soil Gas Flux	80
3.5.2.1 Carbon Dioxide	80
3.5.2.2 Methane	82
3.5.2.3 Nitrous Oxide	83
3.5.3 Relationship Between Flux and Depth of Added Soil/Depth of Removed Soil	86
3.5.3.1 Column Experiment Part 1: Soil Addition	86
3.5.3.2 Column Experiment Part 2: Soil Removal	86
3.5.4 A Comparison of the Field Experiment and Column Experiment Findings	87
3.6 Summary and Conclusions	88
4.0 IMPACTS OF LANDSCAPE RESTORATION ON PLANT SPECIES AND ABUNDANCE	91
4.1 Abstract	91
4.2 Introduction	91
4.3 Materials and Methods	95
4.3.1 Field Experiment	94
4.3.1.1 Plant Measurements	95
4.3.2 Weed Seed Bank Experiment (Growth Chamber)	95
4.3.2.1 Soil Collection and Preparation	96
4.3.2.2 Plant Measurements	98
4.3.3 Data Analysis	99
4.3.3.1 Field Data	99

4.3.3.2 Weed Seed Bank Study.....	99
4.4 Results.....	99
4.4.1 Plant Parameters.....	99
4.4.1.1 Crop Emergence and Yield.....	99
4.4.1.2 Plant and Species Abundance.....	100
4.4.2 Weed Seed Bank in the Soil Profile (Seed Bank Experiment).....	107
4.5 Discussion.....	110
4.5.1 Restoration Effects on Plant Parameters.....	110
4.5.2 Implications of the Soil Seed Bank on Landscape Restoration.....	113
4.6 Summary and Conclusions.....	114
5.0 GENERAL DISCUSSION.....	116
6.0 SUMMARY AND CONCLUSIONS.....	120
7.0 CONTRIBUTIONS TO KNOWLEDGE.....	123
8.0 REFERENCES.....	125
9.0 APPENDICES.....	133
A - Soil Addition and Removal Depth Measurements, Field Plans and Topographical Survey Maps.....	133
B – Gas Chromatograph Specifications.....	159
C – Soil Microbial Biomass Extraction Methodology.....	159
D – Soil Test Analysis Methodologies.....	159
E – Soil Temperature Data.....	159
F – Soil Gravimetric Moisture Content Data.....	159
G – Greenhouse Gas Flux Data.....	159
H - Calculation of the Expected Percent Increase in Organic Carbon in the Addition Soil (FE1).....	159
I – Crop and Weed Emergence Data for FE2.....	159
J - Soil Seed Bank Experiment Data.....	161
K - Core Method for Bulk Density Determination.....	161
L - Nitrous Oxide, Carbon Dioxide, and Methane Flux Equations.....	161
M - Bulk Density Measurement Data.....	161
N - Cumulative Greenhouse Gas Flux Data.....	161
O - Column Experiment Part 1 (Addition Experiment and Column Experiment Part 2 (Removal Experiment) Cumulative Flux Data.....	161

LIST OF TABLES

Table	Page
3.1 A summary of the field and growth chamber experiments.....	31
3.2 Treatment descriptions for FE1 in (a) the upper and (b) the lower slope areas.....	36
3.3 Fall soil fertility results in the upper slope area of FE1 in 2003.....	53
3.4 Fall soil fertility results for the lower slope area of FE1 in 2003	54
3.5 Spring soil fertility results for the upper slope area of FE2 in 2004.....	55
3.6 Microbial biomass carbon results in the upper slope area of FE1	56
3.7 Microbial biomass carbon results in the lower slope area of FE1	57
3.8 Microbial biomass carbon results ($\mu\text{g g soil}^{-1}$) for two sampling dates in the upper slope area of FE2 in 2004	58
3.9 Soil gas flux for the soil addition experiment (Column Experiment 1).....	74
3.10 Soil gas flux for the soil removal experiment (Column Experiment 2).....	75
4.1 Crop emergence, weed emergence, weed species numbers and crop yield for the upper slope area of FE1	102
4.2 Crop emergence, weed emergence, weed species numbers and crop yield for FE2, 2004.....	103
4.3 Weed species (number of plants m^{-2}) in the upper slope area of FE1	105
4. 4 Plant emergence, plant species numbers and above ground plant biomass in the lower slope area for FE1	106
4.5 Plant species (number of plants m^{-2}) in the lower slope area of FE1	107
4.6 Weed species (plants m^{-2}) in the upper slope area in FE2	107
4.7 Weed species identified in the weed seed bank experiment.....	1099

LIST OF FIGURES

Figure	Page
3.1 Soil disking in the soil addition treatments of the upper slope area.	35
3.2 Soil gravimetric moisture (bars) and soil temperature (symbols) (5 cm below the soil surface) in the upper slope area for (a) Block A and (b) Block B in 2003 and 2004	47
3.3 Soil gravimetric moisture (bars) and soil temperature (symbols) (5 cm below the soil surface) in the lower slope area for (a) Block A and (b) Block B in 2003 and 2004	48
3.4 Precipitation at the MZTRA research farm over the (a) 2003 and (b) 2004 sampling periods.....	49
3.5 Soil gravimetric moisture (bars) and soil temperature (symbols) (5 cm below the soil surface) in FE2, 2004. a-b Mean bar values followed by the same letter (within sampling dates) are not significantly different. LSM means groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$; values indicate the means of 3 replicates	50
3.6 Soil respiration (CO_2 flux) in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data but non-transformed data are presented.	63
3.7 Soil respiration (CO_2 flux) in the lower slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data but non-transformed data are presented.	64
3.8 Soil respiration (CO_2 flux) in the upper slope area for FE2 in 2004. a-b Mean values followed by the same letter (within sampling dates) are not significantly different. LSM means groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$; values indicate the means of 3 replicates.	65
3.9 Soil methane flux in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004.*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data non-transformed but data are presented.	66

3.10 Soil methane flux in the lower slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at P<0.10; ** Significant at P<0.05; ***Significant at P<0.01; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data but non-transformed data are presented.	67
3.11 Soil methane flux in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at P<0.10; ** Significant at P<0.05; ***Significant at P<0.01; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data non-transformed but data are presented,	69
3.12 Soil nitrous oxide flux in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at P<0.10; ** Significant at P<0.05; ***Significant at P<0.01; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data but non-transformed data are presented	69
3.13 Soil nitrous oxide flux in the lower slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at P<0.10; ** Significant at P<0.05; ***Significant at P<0.01; Values indicate the means of 3 replicates; significance symbols following differences are based on log-transformed data but non-transformed data are presented.	70
3.14 Soil nitrous oxide flux in the upper slope area for FE2 in 2004.....	71
4.1 The excavation area for one replicate of each depth increment.....	97
4.2 The soil profile within the excavation area.....	97
4.3 Mean total plant emergence by soil depth. Values represent the mean of three reps combined across three sites; error bars represent standard deviations. Letters indicate significant (P<0.05) treatment differences based on LSMeans groupings using log ₁₀ (X+1)-transformed data but non-transformed data are presented.	108
4.4 Mean total plant species by soil depth. Values represent the mean of three reps combined across three sites; error bars represent standard deviations. Letters indicate significant (P<0.05) treatment differences based on LSMeans groupings.	109

1. INTRODUCTION

Soil erosion by wind, water and tillage results in the redistribution of soil within the landscape. In topographically complex landscapes where conventional tillage is practiced, organic-rich topsoil becomes redistributed from hilltops (or convexities) to lower slope and depressional areas (or concavities). When hilltops become eroded they lose productivity while the lower slope and depressional areas often become more productive. The yield variability within the landscape resulting from soil erosion makes agricultural land management difficult and reduces the economic viability of the cropping system.

There are several ways hilltops can be managed to improve their productivity. These include increasing the fertility of the hilltop via chemical or organic fertilizers, increasing the organic matter content of the hilltop by using manure, green manures or altering the crop rotation, and by implementing conservation tillage practices. Adding fertilizer or manure may improve the short-term productivity of the eroded hilltops, but may have limited economic return. Using crop rotation or conservation tillage requires many years of implementation to improve soil productivity. An innovative approach to improving eroded hilltops quickly and effectively is landscape restoration.

Landscape restoration is a land management practice whereby soil that has accumulated in lower slopes and depressions is returned to the adjacent eroded hilltops. This technique has been practiced for many years by farmers in China and some parts of the U.S. and Canada, but has not been documented. Some researchers have looked at crop response to the application of soil and have found optimistic results, however,

landscape restoration as a general management practice has not been studied. General agronomic, economic and environmental factors of landscape restoration should be considered. For example, there are many potential environmental impacts that may be associated with the movement of organic-rich soil within a landscape that have not been considered. These include pesticide fate, nutrient cycling as it relates to runoff, leaching, and greenhouse gas emissions, as well as impacts on plant species and abundance including crops, weeds, and native plants. The latter two topics are the focus of this study.

The purpose of this thesis project was to determine the medium-term effects of landscape restoration on greenhouse gas emissions and on plant species and abundance.

The objectives of this study were:

- 1) To study greenhouse gas emissions (carbon dioxide, methane and nitrous oxide) following landscape restoration comparing the restored areas to the eroded area in the upper slope area, and comparing removal areas to the accumulated areas in the lower slope area.
- 2) To study plant emergence (including crop and weeds) and examine weed composition in the area that is restored, and monitor plant species composition where soil was removed.

Several experiments were conducted to fulfill each of these objectives. Two field experiments were initiated to address both objectives, while one growth chamber experiment used intact soil cores to address objective 1, and a second growth chamber experiment looked at the seed bank within the soil profile to address objective 2. The findings from the growth chamber studies were compared with field experiment findings.

2. LITERATURE REVIEW

Erosion processes and erosion effects on crop productivity are outlined in this chapter. A review of the current methods for improving soil quality on eroded hilltops is included. Past research on landscape restoration and its effects on plant species and emergence, including crop, weeds and native plants, and on greenhouse gas emissions is summarized.

2.1 Soil Erosion in Topographically Complex Landscapes

The three predominant forms of soil erosion reported in the literature include water, wind and tillage erosion. Each form of erosion acts on the soil in different ways; each are predominant in different types of landscapes and may cause the redistribution of soil from and to different parts of the landscape. This section briefly describes the three types of soil erosion including soil and landscape factors affecting erosion.

2.1.1 Water and Wind Erosion

Water erosion occurs in moderately to strongly sloping landscapes and is most severe on soils of silty texture and with poor aggregate stability (Troeh et al.1980). When considering landform, soil tends to be lost from concave curvatures or backslopes (Lobb and Kachanoski 1999).

Wind erosion is most common on level landscapes. Fine sands and silts are prone to wind erosion, especially when soil moisture is low. When the landscape is topographically complex, wind erosion is most severe on exposed hilltops within the landscape (Lobb 1999). Although wind erosion is responsible for the loss of soil from

hilltops, it is not the predominant erosion acting on hilltops of the prairie pothole region of Canada. The losses from hilltops are equally severe on a range of soil types on both sheltered and unsheltered hilltops indicating wind erosion cannot be the only cause (Lobb 1999).

Brady and Weil (2002) report 4 billion Mg soil per year is moved by water and wind erosion in the U.S. Both wind and water erosion are thought to be the predominant forces degrading agricultural soils, including the loss of soil from hilltops and ridges. However, there is little experimental or theoretical evidence that can adequately explain the degree of soil erosion on hilltops (Lobb et al.1995). Recent research has found that tillage erosion is in fact the cause (Govers et al. 1999).

2.1.2 Tillage Erosion

Tillage erosion is responsible for the loss of large quantities of soil from crest and shoulder slope landscape positions (Lobb et al.1995; Govers et al.1999;). Soil lost from these convexities accumulates in the concave footslopes and hollows (or depressions) downslope (Govers et al.1999). Slope gradient strongly affects tillage erosion (Van Muysen and Govers 2002) while other factors such as tillage depth, tillage speed, tillage direction (upslope vs. downslope tillage), and soil condition also play an important role (Lobb et al. 1995; Lobb et al. 1999; Govers et al. 1999; Van Muysen et al. 2002).

Tillage erosion has been reported to account for soil losses of $54 \text{ t ha}^{-1} \text{ year}^{-1}$ from upland shoulder slope positions in Southern Ontario (Lobb et al. 1995). This level far exceeds Canada's Agri-Environmental Indicator - Risk of Tillage Erosion tolerable level of $6 \text{ t ha}^{-1} \text{ yr}^{-1}$ (King et al. 2000). In Manitoba it was estimated that 56% of cropland fell under unsustainable soil conditions in terms of tillage erosion risk (King et al. 2000). It is

clear that tillage erosion is an important process affecting soil quality and crop productivity in agricultural landscapes (Li and Lindstrom 2001).

2.2 Effects of Soil Loss on Crop Productivity

‘The effects of erosion depend largely on the thickness and the quality of the topsoil (being lost) and the nature of the subsoil’ being exposed (Frye et al.1985). The loss of topsoil from hilltops can dramatically reduce the productivity of those hilltops for various reasons.

When topsoil is lost, the total amount of soil organic matter¹ is reduced on the eroded areas (Lal 2000; Frye 1985; Lobb1995; Tanaka 1989). The loss of organic matter brings with it altered chemical and physical properties, such as reduced fertility and lower soil water holding capacity (Lal 2000; Langdale 1982).

Soil fertility is reduced in eroded soils (Frye et al. 1985). With each Mg ha⁻¹ of lost organic matter, approximately 60 kg of nitrogen (N) is also lost (Frye et al. 1985). Phosphorus concentrations of soil may also be lowered. Larney et al. (2000a) found the extractable phosphorus (P) concentration decreased with increasing depth of soil removal at two of three sites while Tanaka and Aase (1989) also found extractable P significantly decreased when soil was removed.

In addition to influencing soil fertility, soil organic matter content influences the soil’s water holding capacity by retaining water that is much more plant available than that held in the mineral fraction (Brady and Weil 2002). When organic matter is lost, so is

¹ The term organic matter refers to the organic fraction of soil consisting of all organic elements including carbon, nitrogen, phosphorus, sulphur, hydrogen and oxygen; the terms organic matter and organic carbon are sometimes used interchangeably in the literature but here, when the term organic matter is used, the organic compounds in general are being reported; when the term organic carbon is used, the measured organic carbon fraction is being reported.

the soil's ability to retain moisture for plants. Organic matter indirectly affects plant available water by stabilizing soil structure and increasing pore space (Brady and Weil 2002; Lal 2000). Soils with higher organic content, therefore, show better water infiltration.

Finally, loss of topsoil may result in soil profile mixing. Any remaining topsoil becomes mixed with subsoil material through further tillage thereby diluting the topsoil's organic matter content (Frye et al. 1985; Tanaka and Aase 1989). When subsoil becomes mixed with topsoil, the inorganic carbon content will increase (Larney et al. 2000b) when subsoils are rich in carbonates. When calcium carbonates are present in the subsoil, this profile mixing may in turn cause problems with soil fertility since small amounts of P can be adsorbed by and/or precipitated with calcium especially when P concentrations in the soil are low (Havlin et al. 1999). A co-precipitation reaction can also occur with sulphate ions and CaCO_3 (Havlin et al. 1999). The availability of P and S is therefore reduced. Subsoils are often stony, lending to undesirable seedbeds and problems with seedling emergence. The subsoil is often less friable and less permeable to air, water and roots (Troeh et al. 1980).

These changes in soil physical and chemical properties ultimately reduce productivity in eroded areas. There is a significant relationship between grain yield and organic C content of soil (Larney et al. 2000b) and grain quality may also be affected (Tanaka and Aase 1989). The negative effect of topsoil loss on soil quality and productivity has been well documented over the past two decades and started as early as 1944 (Horner et al. 1944; Lyles 1977; Langdale and Shrader 1982; Sadler 1984; Burnett et al. 1985; Meyer et al. 1985; Carter et al. 1985; Battiston et al. 1987; Larney et al. 1998;

Schumacher et al.1999;Larney et al 2000 a and b; Lal et al. 2000). The effects of erosion are not always recognized, however, due to the use of fertilizers and other restorative amendments.

2.3 Current Methods for Improving Soil Quality and Productivity

Producers often deal with poor soil quality on hilltops by increasing fertilizer use, adding organic amendments, and changing cropping and soil management practices. There are several studies that looked at the effects of one of, or combinations of, these practices on previously eroded or artificially eroded soils.

2.3.1 Fertilizer Use

Since nutrient deficiency is a common problem of eroded soils, it makes sense to increase fertilizer inputs in eroded areas. The introduction of precision farming has made variable rate application of fertilizer possible (Beckie et al. 1997). Adding increased amounts of fertilizer, however, may not be sustainable in the long-term both agriculturally and economically. In fact, it has been well documented that fertilizer addition alone will not return crop yields to pre-erosions states (Masse and Waggoner, 1985; Mielke and Schepers, 1986; Dormaar et al. 1988; Robbins et al. 1997; El-Swaify 2000). Masse and Waggoner (1985) found that large amounts of N fertilizer were unsuccessful at achieving high yields in artificial erosion plots where topsoil was lost or absent in an intermountain dryland region of Idaho. This study concluded that “adding N fertilizer each crop year was only a partial solution to inadequate topsoil.” In contrast, there is evidence that using both N and P in different combinations was effective at

increasing yield on eroded soil to equal to or greater than the level of the non-eroded check (Tanaka and Aase 1989; Dormaar et al. 1997)

There is evidence supporting and contradicting the benefits of adding fertilizer to eroded areas to improve yield. Even if fertilizer does successfully restore crop yield to non-eroded soil yield levels, there may not be an economic benefit to increasing fertilizer use (Smith et al. 2000). For example, Smith et al. (2000) evaluated the economics of N and P fertilization to restore wheat yields in the Brown and Dark Brown soil zones of the Canadian Prairies. They found that when economic optimum fertilizer levels were used crop yields declined as depth of erosion increased and, therefore, the net return over fertilizer cost also declined with increased erosion. They concluded that there was little to no economic benefit of applying additional inorganic fertilizer to eroded soils.

2.3.2 Use of Organic Amendments

Manure can be a good source of macro- and micronutrients while also providing long-term improved physical characteristics (Dormaar et al. 1988). Most research shows multiple benefits from adding manure to artificially eroded soils including increased yields, increased organic matter content and improved soil physical properties (Larney et al. 2000 a and b; Dormaar et al. 1988; Dormaar et al. 1997; Robbins et al. 1997; Izaurrealde et al. 1997). Larney et al. (2000a) found manure (applied at a rate of 75 Mg ha⁻¹ wet weight or 22 g kg⁻¹ total N and 190 g kg⁻¹ total C) was effective at restoring eroded soils over the two-year study period (or medium-term), however, the duration of these effects were uncertain and needed further study. Larney et al. (2000b) also reported manure addition significantly increased the organic C content of the soil (by 1.0%: from 10.7 in check to 11.9 in manure treatment) such that water holding capacity and yield was

increased. This enhancement was linearly and inversely related to organic C concentrations of the recipient soil. Dormaar et al. (1988 and 1997) showed manure significantly increased soil organic matter (OM); however, the increase was not substantial. They (Dormaar et al. 1988) concluded that adding manure annually for several years may be required to rebuild the OM of eroded soil. Manure also added greater biological activity as indicated by measuring dehydrogenase activity (Dormaar et al. 1997). Robbins et al. (1997) showed adding manure benefited soil quality by increasing the organic carbon of the subsoil to the level of the topsoil (increased by 50% i.e. doubled the amount) after four study years. Izaurralde et al. (1997) also found a significant increase in OM (1.73 to 3.47%) with manure when soil was severely eroded (20 cm cut).

There are additional sources of organic matter that may also be used as ammendments; these include whey (cottage cheese by-product), sugar by products, crop biomass and green manures. Organic ammendments in the form of whey and wheat straw did not benefit yield or OM of eroded soil (Robbins et al. 1997; Dormaar et al. 1997). Robbins et al. (1997) added two rates of whey and found it did not affect bean seed yields or OM content. Dormaar et al. (1997) repeated the study reported on in 1988 to include the addition of wheat straw plus commercial fertilizer. They found this treatment did not increase OM or total N of the soil over time and had little beneficial effect on wheat yield. Green manures may positively affect soil quality attributes essential to improving the productivity to eroded soils. Biederbeck et al. (1998) found an annual legume green manure had a strong influence on soil quality in the short-term including improved C and N mineralization, wet aggregate stability and light fraction organic matter.

Based on the literature reviewed here, manure appears to be a somewhat effective organic amendment for restoring crop yields; however, there are other important factors associated with manure application that must be considered. These include a) the availability of manure (not all producers cropping severely eroded topographically complex landscapes are also livestock producers); b) the cost of manure application increases as the distance from the facility increases therefore, manure application may not be a good option for all of the eroded areas of a producer's entire land base; c) eroded areas may be environmentally sensitive areas (e.g., nearby wetlands) that are not suitable for manure application; d) manure application to the same field year after year is not a common practice. For these reasons, alternatives to manure for improving eroded areas are still needed.

2.3.3 Cropping System

Crop rotation is known to play an important role in cropping systems by optimizing water and nutrient use, by breaking disease, insect, and weed growth cycles and thereby improving yields (Saskatchewan Agriculture and Food 1997; Stevenson et al. 1998; Stevenson and van Kessel, 1996). For these reasons, there is the potential for crop rotation to improve eroded soils. The literature regarding specific rotation effects on eroded soil is minimal. Robbins et al. (1997) included rotations in their study of improving exposed subsoils and concluded that the bean yield increase resulting from crop rotation was only minor when compared with the manure addition treatment. Although crop rotation is a beneficial tool in crop management, it may not be the ideal tool for restoring eroded soils.

2.3.4 Tillage System

The adoption of conservation tillage, specifically zero-till is increasing across the Canadian Prairies (Campbell et al. 2001). By reducing tillage practices, tillage erosion is reduced and the degraded soil is allowed to rebuild because a greater amount of organic matter from crop residues remains on and in the soil profile (Lal and Kimble 1997). However, conservation tillage takes years to rebuild topsoil to pre-erosion levels, and an alternative/additional management option that provides rapid results for improving topsoil levels, and ultimately soil productivity, is needed.

2.4 Soil Addition/Landscape Restoration

Landscape restoration as a soil management practice has not been studied, but there is some research documenting the effects of topsoil addition as a restorative amendment and in determining the impacts of soil erosion.

In 1986 in Nebraska, Mielke and Schepers (1986) monitored crop response to the replacement of lost topsoil (0, 10 and 20 cm additions). They found improved crop emergence as indicated by earlier emergence and greater total emergence, and overall greater corn yields were achieved. Interestingly, soil water content did not differ between added soil and control treatments. However, when below-normal precipitation occurred, the 20 cm topsoil addition treatment achieved 5 and 22% higher yields compared to the 10 cm and 0 cm soil addition treatments, respectively. The topsoil treatment consistently increased the dry-matter production of oats and the greater amount of added topsoil positively influenced total N uptake (however, grain N concentration did not differ

among treatments). One important conclusion was that crop response to topsoil addition was more favorable in years where there was greater plant stress.

There were several studies that included a topsoil amendment treatment in their analysis of amending eroded soils using fertilizer and/or manure. The depth of the topsoil treatment was 5 cm for most studies. Dormaar et al. (1997) repeated the study reported on in 1988 this time including a topsoil addition treatment. Topsoil addition immediately increased the OM and total N content of the soil, however, these effects declined over time. This may have been the result of soil profile mixing because 5 cm of topsoil was not thick enough and easily became incorporated with the subsoil. Larney et al. (2000 a) explain that topsoil plays an important role beyond fertility and water storage as indicated by the fact that irrigation and fertilizer use did not offset the effects of lost topsoil. They also found greater organic carbon content resulted from adding topsoil compared with adding fertilizer, however, manure addition proved to add the greatest amount of OC (0.4 and 1.1 g kg⁻¹ higher than the checks at the Lethbridge Dryland and Lethbridge Irrigated sites, respectively) (Larney et al. 2000b).

Massee and Waggoner (1985) studied the effect of topsoil depth on soil moisture regimes and on fertilizer response. Among the artificial erosion treatments, they included a topsoil addition treatment where 15 cm of topsoil was placed over the original profile. Compared with the untreated soil plot, the topsoil addition treatment showed a 40% yield increase when no nitrogen was added. Also, water-use efficiency increased with added soil. When adding 15 cm of topsoil to the original soil surface (15 cm Ap layer), Massee (1990) found the additional topsoil had a beneficial yield effect. Providing twice the

active topsoil to maintain soil moisture favored nitrification of the added ammonium based fertilizer. They concluded that topsoil provides benefits to yield and water storage.

Verity and Anderson (1990) characterized soil erosion effects on soil quality and yield by adding incremental depths of topsoil to eroded knolls (5, 10, and 15 cm). Wheat grain yields resulting from the restored soil exceeded the control by 45 and 58% in the first year and 42 to 88% in the second. In this comparison, the 5 cm addition treatment resulted in yields that were similar to the 10 and 15 cm treatments. The straw yields from these treatments were also higher, ranging from 62 to 88% over the control. The 10 and 15 cm soil addition treatments yielded significantly greater straw yields in year two.

There are benefits to topsoil beyond nutrients alone (Tanaka and Aase 1989). It has been demonstrated that adding soil can improve some indicators of soil quality including productivity (in terms of yield), organic matter content (to varying degrees) and soil water holding capacity. Topsoil that is eroded by tillage moves down slope and accumulates in depressions but is not degraded, it can be a good source of organic matter. Using this accumulated soil and adding it to eroded hilltops to reverse the effects of tillage erosion is possible; however, landscape restoration as a soil management practice has not been researched. There are many potential environmental and agronomic factors that may be affected by moving soil within the landscape. Research is needed in the areas of environmental impacts, such as impacts on greenhouse gas emissions, and agronomic impacts beyond crop yield, such as impacts on weeds. Furthermore, research on soil removal from hill bottoms/depressions does not exist.

2.4.1 Soil Addition Effects on Greenhouse Gas Emissions

Currently there is pressure on the agriculture sector to reduce net GHG (mainly carbon dioxide, methane and nitrous oxide) emissions through the implementation of beneficial management practices, including those aimed at increasing carbon storage in soil (Janzen et al. 1998). The potential impact of landscape restoration on GHG is, therefore, important because it will affect soil carbon within the landscape. Changes in soil carbon may in turn induce changes in greenhouse gas emissions including carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O). The literature regarding landscape restoration/application of topsoil effects on greenhouse gas (GHG) emissions does not exist. There is, however, a lot of research on GHG dynamics in agroecosystems as influenced by management system and/or soil parameters (Burton and Beauchamp 1985; Mahli et al. 1990; Lal and Kimble 1997; Pol-van Dasselaar et al 1998; Janzen et al. 1998). Soil erosion will likely influence C flux (and other GHG emissions) because of its effect on a number of processes including: 1) removal of soil organic carbon-rich soil, 2) burial of soil through deposition, 3) textural changes with subsoil exposure, 4) microbial activity due to soil moisture and temperature changes, and 5) plant growth and residue amounts (Bajracharya et al. 2000). Some of these processes are also influenced by landscape position (Pennock et al 1992; Corre et al.1996). Therefore, it is expected that landscape restoration will effect GHG emissions by influencing the same process affected by soil erosion and landscape factors. More research is needed to assess whether landscape restoration will positively or negatively affect GHG emissions from the landscape.

2.4.2 The Effects of Topsoil Addition on Weeds

The literature regarding landscape restoration/addition of topsoil effects on weeds does not exist; however there is some research that is applicable. One relevant topic in weed research is the weed seed bank. Much of the research looks at trying to characterize the seed bank on a horizontal plane (across level landscapes) with correlations to existing weed populations (Benoit et al. 1992) and few have looked at the relationship between seed bank and soil depth. It has been documented that most viable weed seeds exist within the top 15 cm of soil, depending on the type of tillage system, crop rotation and weed management practices (Barberi and Lo Cascio 2001; Felix and Owen 2001; Buhler et al. 1997; Cavers 1995). This has implications for landscape restoration because it is the surface layer of soil that will be removed and applied to eroded hilltops. The composition of the seed bank within the soil profile, and the proportion of viable seed banks below the soil surface of accumulated soil are important for assessing the potential weed populations that might arise following landscape restoration in areas where soil is removed and where soil is applied.

While the weed seed bank relates to weed population variability in time, a second relevant topic is the variation of weed populations in space. Weed populations will vary within the landscape due to landscape variations in soil physical and chemical properties such as soil type, moisture, texture, and fertility (Dieleman et al. 2000). Therefore, it is expected that weed populations may be influenced by changes in soil properties within the landscape caused by landscape restoration.

3. IMPACTS OF LANDSCAPE RESTORATION ON GREENHOUSE GAS EMISSIONS

3.1 Abstract

Soil management can lead to increased or decreased greenhouse gas emissions; however, the variability of soil properties, and thus gas emissions, in space make it difficult to predict these effects. Landscape restoration is one soil management practice that can reduce the spatial variability of some soil properties in topographically complex landscapes, including soil organic matter. This study examined the effect of landscape restoration on greenhouse gas emissions (CO₂, CH₄, and N₂O) from the soil. Landscape restoration did not affect greenhouse gas emissions in the upper slope area where soil was applied. Where soil was removed, soil CO₂ flux was decreased in the first year following landscape restoration. Changes in the soil environment including soil nitrate and soil microbial biomass carbon were also observed.

3.2 Introduction

Increase in the atmospheric concentration of greenhouse trace gases (e.g., CO₂, CH₄, N₂O) since pre-industrial times has led to climate change (Mosier 1998). Globally, agriculture contributes greenhouse gas emissions representing approximately one-fifth of the annual increase in global radiative forcing of climate change (Cole et al. 1997). In Canada, agriculture contributes 10% of anthropogenic greenhouse gas emissions (Environment Canada 2005). Soils act as a major source and potential sink for greenhouse gases (Janzen et al. 1998) and, therefore, managing the soil environment can

lead to increased or decreased emissions. The influence of soil management on net emissions is difficult to predict due to the inherent variability of the soil environment in space and time (Mosier 1998; Corre et al. 1996). Soil properties and, as a result, greenhouse gas emissions, vary greatly within the landscape (Meixner and Eugster 1999).

Tillage can affect soil properties and thus greenhouse gas emissions from soil. Conservation tillage, specifically no-till, is a beneficial management practice (BMP) designed to reduce erosion processes, improve the overall quality of the soil environment and reduce landscape variability. Soil properties including soil organic matter content, soil moisture and soil nutrients all influence greenhouse gas emissions and, therefore, conservation tillage will also affect greenhouse gas emissions from the landscape. In fact, conservation tillage is known to increase soil organic carbon in the surface layer (Lal and Kimble 1997) resulting in reduced CO₂ emissions. Also, there is evidence of a strong positive correlation between organic carbon in the soil and nitrous oxide emissions (Malhi et al. 1990; Luo et al. 1998; Lemke et al. 1998).

Soil erosion can influence the spatial variability of soil properties in topographically complex landscapes. Soil erosion by wind, water and tillage results in the redistribution of soil within and often beyond the landscape. Tillage erosion is largely responsible for soil variability within the landscape by displacing organic rich topsoil from hilltops and moving it to lower slopes and depressions (Lobb 1999). Hilltops low in organic matter consequently have lower water and nutrient holding capacity and show low agricultural productivity; lower slopes and depressions tend to have an accumulation of organic rich topsoil with higher moisture and nutrient holding capacity and also greater productivity (Lobb 1999).

Landscape restoration is an innovative soil management practice that can be used in combination with conservation tillage to further reduce the variability of soil properties within the landscape. Landscape restoration is a practice whereby soil accumulated in lower slope and depressional areas is removed and added to eroded hilltops or upper slope areas. Adding soil to hilltops will likely increase soil organic matter of the surface soil; removing soil from lower slopes and depressions will also influence soil organic matter therein. By altering the organic matter within the landscape and soil properties influenced by soil organic matter, such as soil moisture and nutrients, there is the potential for landscape restoration to affect greenhouse gas emissions from the landscape. Landscape restoration is a practice with the potential for widespread adoption in areas where tillage erosion is prevalent and therefore, the impact of landscape restoration on greenhouse gas emissions from the landscape should be understood.

The objective of this study was to measure greenhouse gas emissions (carbon dioxide, methane and nitrous oxide) from the soil following landscape restoration, focusing on the hilltops where soil was added and on the lower slope and depressional areas from where soil was removed. A complementary growth chamber study was carried out using intact soil cores to determine if the depth of added and the depth of removed topsoil will influence gas flux. The findings from the growth chamber study were compared to the field experiment findings. The medium-term impacts of landscape restoration on greenhouse gas emissions are discussed.

3.3 Materials and Methods

Field and growth chamber experiments were conducted to address the thesis objectives which include the study of the impacts of landscape restoration on greenhouse

gas (GHG) emissions and on plant species and abundance (Table 3.1). This section provides a description of the study site where the field experiments were carried out and from where soil was collected for the growth chamber experiments. The complete methodology for the field and growth chamber experiments designed to study greenhouse gas emissions is presented. The methodology for the study of plant species and abundance in the field and the complementary growth chamber experiment is included in Section 4.3.

Table 3.1 A summary of the field and growth chamber experiments

	Study A – Greenhouse gas emissions	Study B – Plant species and abundance
Field Experiment 1 – FE1	Sites identified fall 2002/experiment initiated spring 2003 and carried out through to fall 2004	
Field Experiment 2 – FE2	Sites identified fall 2003/experiment initiated fall 2003 and carried out through to fall 2004	
Growth Chamber Experiment 1 – GC1	Soil Column Experiment – Part 1 and Part 2 carried out winter/spring 2004	
Growth Chamber Experiment 2 – GC2	Weed Seed Bank Experiment carried out fall 2003	

3.3.1 Site Selection and Description

The study area exists within the Aspen Parkland of the Prairies Eoregion of South-western Manitoba and is also considered to be a part of the Prairie Pothole Region of the province. Within this area, the Manitoba Zero Tillage Research Association (MZTRA) farm is located (Section 31-12-18) and was chosen as the study site. The landscape is topographically complex and contains numerous wetlands (or potholes) that

are ephemeral to permanent in nature. The soils are predominantly Black Chernozems formed over calcareous glacial tills. Mean annual temperature for the area is 1.4°C and total precipitation is 460 mm with 340 mm mean annual rainfall (Podolsky and Schindler 1993). The land-use is largely agricultural with a focus on cereal, oilseed and livestock production.

The landscape of the MZTRA farm is characterized as undulating to hummocky with gently sloping (2-5%) topography (Podolsky and Schindler 1993). Soils are of the Newdale Association and are representative of the Newdale soils in the Parkland Region (Podolsky and Schindler 1993). The study site consists predominantly of Newdale Series soils with a small portion belonging to the Rufford, Cordova, Drokan and Angusville Series. Newdale soils (Orthic Black Chernozem) generally appear in the mid to upper slope positions while the Rufford (Rego Black Chernozem) and Cordova (Calcareous Black Chernozem) soils are found in upper slope positions and knolls. Drokan soils are typically found in depressional positions while the Angusville soils are found in the middle and lower slope positions of the undulating topography. In the lower slope areas, drainage ranges from poor in the depressions to well in the upper slopes. Twenty-two percent of the soils on the farm are considered weakly saline (Podolsky and Schindler 1993). Higher salinity levels occur in saturated soils or soils found adjacent two permanent sloughs and water bodies. The Drokan series found at the study site was characterized by Podolsky and Schindler (1993) as being weakly saline (4-8 mS cm⁻¹). Bulk density ranged from 0.99 and 1.08 g cm⁻³ in the top 12 cm of the Newdale and Rufford soils, respectively, and 1.24 g cm⁻³ in the top 17 cm of the Angusville soils. Bulk density measurements were not available for the Drokan and Cordova soil series.

The MZTRA farm has been under zero-till crop production since 1993 and was intensively tilled for decades prior to that. The farm is currently managed under a 5-year crop rotation under two different cropping systems: an annual rotation consisting of cereals and oilseeds as well as a livestock rotation consisting of cereals and oilseeds and alfalfa for hay production and for grazing cattle. The study site exists within the annual cropping system. The farm manager is responsible for all seeding, harvesting and herbicide application operations.

There is visual evidence of the tillage erosion that occurred prior to 1993 still present on many of the hilltops and upper slope areas of the MZTRA farm. This includes stoniness, high CaCO_3 content, low soil organic matter (SOM) at the surface, and a thin A horizon in soils occurring in upper slope areas; and a thick, organic-rich A horizon in soils in lower slope and depressional areas. The severity of past erosion on the hilltops was determined to be $20\text{-}40 \text{ t ha}^{-1} \text{ yr}^{-1}$ during the period of conventional tillage between 1960 and 1993 using the ^{137}Cs method described by Lobb and Kachanoski (1999).

Several severely eroded ridges and knolls were identified in the center of the west half of the MZTRA research farm in the fall of 2002 for the purpose of studying greenhouse gases and plants following landscape restoration. The first field experiment was established in the spring of 2003, while the second field experiment was established in the fall of 2003 (Table 3.1).

Two growth chamber experiments were conducted using soil collected at the MZTRA farm. A soil column experiment (Section 3.3.3) used intact soil cores sampled from a landscape on the NE quarter of the farm (GC 1). The landscape was similar in form, soil composition, and severity of erosion to that used in the field experiments. A

weed seed bank experiment (Section 4.3.2) used soil collected from the same site as the field experiments (GC 2).

3.3.2 Field Experiments

Two field experiments were conducted with the same objectives but with slightly different experimental designs; one initiated in spring of 2003 and the other initiated in the fall of 2004.

3.3.2.1 Field Experiment 1 (FE1 2003-2004). A restoration experiment was initiated in May 2003 at the MZTRA research farm in a cropped landscape adjacent to a large permanent wetland and was conducted over two growing seasons (2003 and 2004). The edge of the wetland consisted of two distinct rings of riparian vegetation, the first containing hydrophilic plants (cattails and sedges) and the second, outermost ring, containing perennial grass (brome) and some perennial broadleaf plants (thistle; meadow arnica). For this experiment, soil was removed from the riparian area (the second, outermost riparian area) where soil had accumulated due to past erosion, and was added to the upper eroded ridge from where the soil was originally lost.

Approximately 20 cm of soil was removed from 3.2 m² plots in the riparian area and applied to 7.3 m² plots on the severely eroded upper slope area to give a final added depth of approximately 10 cm on May 5th, 2003. The soil was allowed to dry for several days and then disked with three passes to break up the sod and any large clods of soil (Figure 3.1 and 3.2). The depth of topsoil added was measured using a meter stick in nine places throughout the plots and averaged to characterize the final depth of addition (Appendix A.1). The depth of topsoil removed was also measured (Appendix A.2). The

plots from where soil was removed contained large wheel tracks and soil debris. The wheel tracks were filled and the soil debris was removed in order to smooth and clean the surface of these plots. There are, therefore, two areas of interest in this experiment – the upper slope area to which soil was added (addition area) and the depression area where soil that had accumulated from past erosion was removed (removal area).



Figure 3.1 Soil disking in the soil addition treatments of the upper slope area.

In the addition area, the treatments included 10 cm topsoil added plus disking (addition), 0 cm topsoil added plus disking (disturbed), and a control where 0 cm topsoil was added (control). The treatments were arranged as a paired treatment comparison with one pair consisting of an addition and disturbed treatment and the other pair consisting of an addition and control treatment (Table 3.2). The plot layout was such that each pair alternated across a ridge to give three replicates each. In the removal area, the treatments were: removal of 20 cm of soil plus removal of standing plant biomass (removal),

removal of standing plant biomass only (disturbed) and the control where nothing was removed (control). Standing plant biomass was removed from the plots by cutting the plants at the soil surface and taking them off the plots. The treatments were arranged in the same paired comparison as the addition area treatments. No attempt was made to pair plots of addition and removal. Both areas of interest were replicated in an adjacent field seeded to a different crop to give two blocks. Greenhouse gas emissions, soil temperature and moisture, soil chemistry and soil organic carbon (SOC), crop emergence and yield, and weed emergence were monitored over two growing seasons.

Table 3.2 Treatment descriptions for FE1 in (a) the upper and (b) the lower slope areas

Treatment	Pair	Treatment Description
<i>(a) Addition Area – Upper slope</i>		
Addition (U1A) Disturbed (U1D)	Pair 1	Addition of 10 cm of soil followed by disking (6 passes) Disking only (6 passes, no soil addition)
Addition (U2A) Control (U2C)	Pair 2	Addition of 10 cm of soil followed by disking (6 passes) No soil addition or disking
<i>(b) Removal Area – Lower slope</i>		
Removal (L1R) Disturbed (L1D)	Pair 1	Removal of standing plant biomass followed by 20 cm of soil removal Removal of standing plant biomass only
Removal (L2R) Control (L2C)	Pair 2	Removal of standing plant biomass followed by 20 cm of soil removal No plant biomass or soil removal

The restoration area plots were seeded by the MZTRA farm manager as a part of normal seeding operations for the farm in both years of the experiment. In 2003, Block A was seeded to flax (cv. “Bethune”) on May 21 at 39 kg ha⁻¹ and 2 cm deep with a Morris Maxim air drill seeder equipped with 2.5 cm knife-type openers on 25 cm spacing and

metal packers. Nitrogen fertilizer (urea-ammonium nitrate 28-0-0) (67 kg N ha^{-1}) was side dribble banded at the time of seeding. Peas (cv. "Mozart") were seeded in Block B on May 15 at 202 kg ha^{-1} and 3 cm deep after being inoculated with a self-stick peat based inoculant. Weed control in both blocks followed normal weed control practices for the MZTRA farm. Both fields were treated with glyphosate (Block A and B received 648 and $432 \text{ g a.i. ha}^{-1}$, respectively) as a pre-seed burn off of winter annuals and volunteer cereals. The flax was sprayed with Flaxmax ($660 \text{ g a.i. ha}^{-1}$) and Poast ($211.5 \text{ g a.i. ha}^{-1}$) and the peas were sprayed with an Odyssey-Poast mix ($31.5 \text{ g a.i. ha}^{-1}$ and $90 \text{ g a.i. L ha}^{-1}$, respectively) for control of grassy and broadleaf weeds. Both crops also received a pre-harvest burn off of glyphosate (864 and $605 \text{ g a.i. ha}^{-1}$) in Blocks A and B, respectively) to help dry down the crop as well as control the Canada thistle (*Cirsium arvense*) in the pea field (Block B).

In 2004, Block A was seeded to canola (cv. "46A76") on May 28 at a rate of 5.6 kg ha^{-1} and 2 cm deep. Nitrogen (67 kg N ha^{-1} equivalent) and sulphur (22.5 kg ha^{-1}) was side dribble banded at the time of seeding. Block B was also seeded to canola (cv. "822 Nexera") on May 21 at the same rate. Nitrogen (74 kg N ha^{-1} equivalent) and sulphur ($22.5 \text{ kg SO}_4\text{-S ha}^{-1}$) were side dribbled banded at the time of seeding and nitrogen (5.6 kg N ha^{-1} equivalent) and phosphorus ($28 \text{ kg H}_2\text{PO}_5 \text{ ha}^{-1}$) were placed with the seed. Both blocks received a pre-seed burn off with glyphosate at the same rates as the preceding year. Block A was treated with Odyssey (42 g ha^{-1}) on June 21 and Block B was first treated with Sevin XLR Plus (0.5 L ha^{-1}) on June 10 and 16 to control flea beetle, and on June 21 with Odyssey (42 g ha^{-1}).

3.3.2.2 Field Experiment 2 (FE2). The second field experiment also consisted of moving soil from the lower slope area to the severely eroded upper slope area, however, cropland depressions were the source of topsoil. The two areas of interest in this experiment are similar as in FE1 – the upper slope (addition area) and the depressional area where eroded soil has accumulated (removal area). This experiment was established in the late-fall of 2003 and only monitored over the growing season of 2004. Three adjacent cropland depressions occurring in the same field as Block B were selected as removal areas for this field experiment. The depressions are all similar in size and are surrounded by eroded ridges. Topsoil thickness in each depression was greater than 0.5 m. The depression was split in half with one side used as the removal area and the other as the control. Each depression was intended to represent one replication (rep) to give a total of three reps each of removal plots and controls. However, due to excess moisture (i.e. flooded conditions) in the spring and summer of 2004, the depression areas were not monitored.

The placement of the treatments was decided by the degree of erosion on the adjacent upper slopes and hilltops as indicated by the amount of and depth to carbonates at the surface as well as the degree of stoniness and the lack of soil organic matter. The upper slope treatments were the same as in FE1 (addition, disturbed, control) and applied in a randomized complete block design. One replicate, 12 m² in size, of each treatment was established in 9 different places along the eroded knolls to give 9 blocks and a total of 27 plots. The treatments were randomly assigned to three plots in each block. The treatments were prepared and applied using the same method as FE1. The depth of topsoil added was measured using a meter stick placed randomly in nine places

throughout the plots and averaged to characterize the final depth of added soil (Appendix A.3). The same parameters were monitored as in FE1.

The addition area plots were seeded as a part of normal seeding operations for the MZTRA farm using the seeding equipment described for FE1; these operations were the same as for Block B in 2004. A problem occurred with the seeder in this field. It appeared that half of the seeder was plugged during seeding operations causing irregular emergence patterns throughout the FE2 plots. FE1 Block B plots did not appear to be affected in this way.

3.3.2.5 Gas Flux Measurement and Analysis. Greenhouse gas emissions (CO_2 , CH_4 and N_2O) were measured throughout the growing season in the upper and lower slope areas to determine the effect of moving soil on soil gas flux. Greenhouse gases were sampled using vented static chambers (Hutchinson and Livingston 2002). The chambers consisted of a 20.3-cm-diameter x 10-cm-high collar and a 20.3-cm-diameter lid. The lid contained a vent tube (0.4-cm-internal-diameter, 7.5-cm-length) and second port fitted with a serum stopper to allow sampling (MacLeod et al. in preparation). Greenhouse gas chambers were installed 5 cm into the soil in each plot in both the removal and addition areas. After securing the lid in place, from the headspace at 30 and 60 minutes following closure. Samples consisted of 15 mL of atmosphere taken from the headspace of the chamber using a 20 mL disposable syringe (Becton-Dickinson), injected into a 10 mL exetainer (Labco, UK) and sealed with silicone sealant. The exetainers had been flushed with helium and evacuated to < 0.5 Torr prior to use. Five replications of fifteen mL samples of air were taken immediately after securing the lids to represent the headspace atmosphere at the start of the flux measurement. Three replications of fifteen mL samples

of two standard gas mixtures were also injected into exetainers and handled in a similar manner as other gas samples. Exetainers were returned to the lab for analysis and gas was analyzed using a Varian Star Gas Chromatograph (Appendix B). The equations used for calculating gas flux from the sampling interval can be found in Appendix L.

Gases were sampled on nine dates between May 27 and September 17 in 2003 and on seven dates between April 21 and Aug 16 in 2004 for FE1; and on seven dates between April 27 and Aug 16 in 2004 for FE2. At each sampling date, soil temperature, and ambient air temperature were measured using a digital thermometer. Soil temperature was measured at two depths by inserting the thermometer into the surface soil and 5 cm below the soil surface. Ambient air temperature was measured by holding the thermometer 30 cm above the soil surface in the shade. Soil gravimetric moisture was also monitored for most sampling dates. Soil samples were collected to evaluate the size of the soil microbial biomass using the chloroform fumigation-direct extraction method (Vance et al. 1987; see Appendix C for a description of the method used). Microbial biomass C and N numbers were expressed simply as the amount of C and N released during fumigation and not corrected using published extraction coefficients (k_c , k_n). These parameters were measured to monitor change in soil biophysical characteristics following restoration as well as to help with the interpretation soil gas flux results.

3.3.2.6 Soil Sampling and Analysis. Soil was sampled from the 0-15 and 15-60 cm depth increments in the fall of 2003 in FE1 and in the spring of 2004 in FE2 for soil nutrient status. Soil samples from FE1 were sent to Agvise Laboratories and analyzed for nutrient status (N, P, K, and S), total carbon including organic carbon and carbonate content and salinity (Appendix D.1). Soil samples from FE2 were sent to South Dakota

State University and analyzed for the same parameters using the same methodologies. The soil nutrient status was monitored to reflect any changes in nutrients, total carbon, carbonates and salinity following landscape restoration. These parameters have important implications for the overall productivity of the soil (although this is not the focus of this study) and soil gas flux.

Soil bulk density was measured once during the duration of the study on July 15, 2004 in both FE1 and FE2. The core method was used to sample bulk density (Appendix K); however, the data for the upper slope area are not consistent with findings by Podolsky and Schindler (1993) and the lower slope area data are very low (Appendix M.2) implying the data is suspect and cannot be referred to with confidence.

3.3.2.7 Rainfall. In year 2003, rainfall was monitored on site using a tipping bucket rain gauge. In 2004, the data was obtained from MZTRA farm records.

3.3.3 Soil Column Experiment

This experiment consisted of a two-part landscape restoration experiment using intact soil columns taken from an eroded upper slope landscape position (Part 1) and from the accumulation area of a cropland depression (Part 2). Greenhouse gas flux from the columns was measured in a growth chamber after treatments were applied.

3.3.3.1 Part 1 – Addition Experiment. In the fall of 2003, intact soil columns were taken from an eroded ridge in the NE quarter section of the MZTRA farm using PVC pipe 20.3 cm in diameter (1.25 cm wall thickness) and equipped with apparatus to secure a gas chamber lid as described in Section 3.3.2.5. The columns were taken to a depth of 30 cm with enough length remaining to add the restoration treatments and allow for the headspace necessary for greenhouse gas flux measurement. This depth was chosen based

on the assumption that gases produced in the upper 30 cm provide the major contribution to surface emissions. The soil columns were frozen, thawed and incubated at 25°C for one week prior to adding the treatments. Incubation allowed the microbial populations to re-stabilize after being frozen and thawed. At the time of taking the soil columns from the field, bulk density samples were taken at four places along the ridge at three depths (0-10 cm, 10-20 cm and 20-30 cm) to provide an estimate of the bulk density. The bulk density method used was the core method (Appendix K). This information was used to determine the volumetric water content of the columns at the start of the experiment.

Soil addition treatments were 0, 10 and 20 cm of added soil using soil that had accumulated in the depression (D) to give treatments 0, 10D and 20D, respectively. A fourth treatment was included where 10 cm of soil was added using soil from the upper (U) eroded ridge named treatment 10U. The 0 cm increment served as a control. Each treatment was replicated three times. Soil was added to the columns at a bulk density of 1.0 g cm^{-3} . Soil columns were stored in a growth chamber at 25 °C, 80% RH. Soil moisture was maintained at 70% of water filled pore space by adding 600 mL of water to the surface of the column at the end of each gas sampling. Columns were open ended and excess water was allowed to drain out of the column. The surface flux was taken on 12 days during a one-month period using the same apparatus and method as described in Section 3.3.2.5. The interval between measurements varied as the experiment proceeded with samples taken frequently following the treatment additions and less frequently towards the end of the one-month period.

3.3.3.2 Part 2 – Removal Experiment. In the spring of 2004, intact soil columns were taken from the depositional area of a depression associated with the eroded ridge used in

Part 1 of the experiment. A depth of 30 cm of intact soil was taken using the same PVC conduit as in the previous experiment. The treatments consisted of 0, 20, and 40 cm of removal and were each replicated three times. For this experiment, the treatments were applied in the field before taking the columns. That is, 20 or 40 cm of soil was removed and then the 30-cm-deep intact column was taken from the new surface and the same assumption was made that only gases produced in the upper 30 cm contribute to surface emissions. The control columns were taken from the original soil surface. Therefore, each treatment consisted of soil from a different part of the soil profile. After removing the columns, bulk density samples were taken from each column sample location at two depths (5 and 20 cm from the surface) to provide an estimate of the bulk density of the soil columns. This information was used to determine the volumetric water content of the columns at the start of the experiment.

Soil columns were stored in a growth chamber at 25 °C, 80% RH. Soil moisture was maintained at 70% of water filled pore space based on soil volumetric moisture prior to starting the experiment by adding 600 mL of water to the surface of the column at the end of each gas sampling. Columns were open ended and excess water was allowed to drain out of the column. The surface flux was sampled on 12 days over a one-month period using the same apparatus and method as described in Section 3.3.3.

3.3.4 Data Analyses

All statistical analyses were performed using SAS version 8 software (SAS Institute Inc. 2000).

3.3.4.1 Field Data. A paired t-test was used to determine treatment differences for FE1. FE2 data were analyzed using a mixed linear model in SAS to test for treatment and rep

effects. The rep*treatment interaction was specified as the random statement. The level of significance used for these experiments was $\alpha = 0.10$ based on the high degree of variability characteristic of uncontrolled, field-based experiments (Corre et al. 1996). In both experiments, the upper slope data was treated and analyzed separately from the lower slope data. The gas flux was calculated as the change in the amount of gas contained in the headspace as a function of time and expressed per unit area based on the area enclosed by the chamber (MacLeod et al. in preparation). Rates of CO₂, CH₄ and N₂O flux were compared on an individual date basis after log(X+1) transformation of the data was made to improve the normality of the distribution of the error variances. LSMEANS were also calculated to compare mean significance groupings of treatments.

3.3.4.2 Column Experiment Data. Gas flux was calculated as the change in the amount of gas contained in the headspace as a function of time and expressed per unit area based on the area enclosed by the chamber (Appendix I). A general linear model analysis of variance tested for treatment effects using SAS version 8. Daily greenhouse gas emission data was log-transformation based on the Kolmogorov-Smirnov test for homogeneity of error variance. On dates where significant treatment effects were observed, Tukey's test determined the minimum significant difference groupings at $\alpha = 0.05$.

3.4 Results

3.4.1. Soil Environment

Soil temperature, gravimetric moisture content, soil nutrient status, and soil microbial biomass were all monitored throughout the sampling periods to document changes in the soil environment following landscape restoration. Soil temperature and

gravimetric moisture were monitored on each greenhouse gas sampling date while soil microbial biomass was sampled once per month in year one and in the spring of year two; soil nutrients were measured in the fall of 2003 for FE1 and in the spring of 2004 for FE2 prior to seeding and fertilization. These characteristics directly affect greenhouse gas production and emission (Corre et al. 1996; Smith et al. 2003) and therefore, understanding the change in the soil environment may also help with the interpretation of gas flux results.

3.4.1.1 Soil Temperature and Moisture. In FE1, soil temperature measurements followed a seasonal trend in the upper slope position in both blocks with lowest temperatures occurring in the spring and fall in 2003 and in the spring and late-summer in 2004 (Figure 3.2). In general, the control treatment (U2C) showed consistently lower temperatures compared with all other treatments. In the lower slope, similar seasonal trends were observed for soil temperature. The removal treatments (U1A and U2A) showed consistently significantly ($P < 0.10$) higher temperatures compared with the control and disturbed treatments for most dates in both years except in Block B in 2004 where the removal treatments were consistently cooler.

Mean soil gravimetric moisture was 24 % in 2003 and 27% in 2004 across Blocks A and B. The pattern in soil moisture is consistent with that of rainfall for both years (Figure 3.3). Significant ($P < 0.10$) moisture differences were most common in Block B where the addition treatment maintained higher soil moisture than the disturbed and control treatments on three dates in both 2003 and 2004 (Figure 3.4). In the lower slope, mean gravimetric moisture was 36% and 40% across both blocks in 2003 and 2004, respectively. In Block B, soil moisture was significantly ($P < 0.10$) higher in the removal

treatments in 2003 (Appendix F.2). However, the removal treatments showed consistently lower soil moisture in all other comparisons.

In FE2, soil temperatures followed a similar seasonal trend as in FE1. The addition and disturbed treatment soils were warmer than the control soil on the first three sampling dates, but all soils maintained similar temperatures throughout the rest of the sampling period.

The range in soil gravimetric moisture was similar to that found in FE1 in 2004. Significant treatment effects were found on April 21 and June 9 where the addition treatment soils showed the highest soil moisture (Figure 3.5).




Figure 3.2 Soil gravimetric moisture (bars) and soil temperature (symbols) (5 cm below the soil surface) in the upper slope area for (a) Block A and (b) Block B in 2003 and 2004 (see Appendix E.2 and F.1 for numeric values and statistical significance).

¹U1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2




Figure 3.3 Soil gravimetric moisture (bars) and soil temperature (symbols) (5 cm below the soil surface) in the lower slope area for (a) Block A and (b) Block B in 2003 and 2004 (see Appendix E.4 and F.2 for numeric values and statistical significance).

¹U1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2




Figure 3.4 Precipitation at the MZTRA research farm over the (a) 2003 and (b) 2004 sampling periods.




Figure 3.5 Soil gravimetric moisture (bars) and soil temperature (symbols) (5 cm below the soil surface) in FE2, 2004. a-b Mean bar values followed by the same letter (within sampling dates) are not significantly different. LSMeans groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$; values indicate the means of 3 replicates (see Appendix E.5 and F.3 numeric values and statistical significance).

3.4.1.2 Soil Nutrients

In the upper slope area of FE1 soil addition significantly increased soil nitrate-nitrogen (NO_3^- -N) in the 0-15 cm depth by 80% over the control in Block A and by 50% over the control in Block B (Table 3.3). Although phosphorus appears to have increased as well, this difference was only significant in Pair 2 of Block A. Sulphate-sulphur (SO_4 -S) increases in the addition treatment followed a similar pattern as nitrogen. Soil calcium carbonates in the surface soil were reduced by adding soil, with a significant reduction of 50% occurring in Block B. Soil addition did not significantly affect the percent total carbon (TC), although a small but statistically significant ($p < 0.10$) increase in total organic carbon (TOC) occurred in both comparisons of Block B. Soil addition also

increased the salt concentration of the soil at the 0-15 cm depth from 0.5 in the control to 0.8 mS cm⁻¹ in the U2A treatment in Block A, and from 0.6 to 1.1 mS cm⁻¹, and 0.6 to 1.4 (mS/cm) in U1A and U2A, respectively, in Block B. At depth (15-60 cm) the percent TC and TOC were found to be significantly higher in U1A of Block B compared with the disturbed treatment, however the difference is small.

In the lower slope area, removal of soil had the greatest effect on soil NO₃-N, P, K, TC and TOC at both sampling depths, with the effects being more pronounced in Block A (Table 3.4). Significant increases in SO₄-S were found in Block B in the 0-15 cm sampling depth. TC and TOC contents were each significantly reduced by an average of 65% in Block A. In Block B, the reduction was small with significant reductions occurring only for TOC contents. At the 15-60 cm sampling depth, similar reductions were found; however, the magnitude of the decrease was slightly less.

In FE2, the addition of soil significantly increased soil nutrients in the surface layer (0-15 cm) (Table 3.5a). Soil nitrate, phosphorus, potassium and sulphate levels were 40, 50, 30 and 90% higher, respectively, in the addition treatment compared with the disturbed and control treatments. No significant treatment effect was observed for total carbon; however, inorganic carbon was significantly lower in the addition treatment compared with both the disturbed and control treatments. Organic carbon can be calculated by subtracting inorganic carbon (%) from total carbon (%). Based on this calculation and the spring soil fertility results, organic carbon was 3.9% in the addition treatment compared with 2.7% in the disturbed treatment and 2.8% in the control. The addition of soil did not alter the pH to any great degree; however salts were increased following soil addition from 0.5 mS cm⁻¹ in the control to 0.7 mS cm⁻¹.

It should be noted that the removal area soils of FE1 (Drokan series) were characterized as being being weakly saline (Podolsky and Schindler 1993) and commonly showed gypsum crystals (CaSO_4) in the surface and subsurface soil horizons. The sulphate measurements provided here represent total sulphate-sulphur (including sulphate associated with gypsum) and therefore high levels of gypsum in the removal area soil may falsely inflate the sulphate-sulphur results of the addition treatments.

Table 3.3 Fall soil fertility results in the upper slope area of FE1 in 2003

Nutrient	Block A ^b						Block B					
	U1A ^c	UID	Δ	U2A	U2C	Δ	U1A	UID	Δ	U2A	U2C	Δ
<i>a) 0-15 cm Upper Slope Area</i>												
NO ₃ ⁻ -N (kg ha ⁻¹)	102	56	46	134	29	105**	64	31	25*	66	27	39*
P-Olsen (kg ha ⁻¹)	36	20	16	40	36	16**	27	18	9	25	20	5
K (kg ha ⁻¹)	1104	782	322	1185	853	332	820	652	168	889	681	208
SO ₄ -S (kg ha ⁻¹)	84	60	24	122	21	100**	134	46	87***	124	69	65**
CaCO ₃ (%)	0.4	4.4	-4.0	1.0	2.7	-1.6	4.4	9.7	-5.3***	4.3	8.7	-4.4*
Total Carbon (%)	5.1	5.2	0.0	5.2	4.1	1.1**	4.1	3.8	0.3	4.3	3.9	0.4
Total Organic Carbon (%)	5.1	4.6	0.7	5.1	3.8	1.3	3.5	2.6	0.9*	3.8	2.9	0.9*
EC ^a (mS cm ⁻¹)	0.7	0.6	0.1	0.9	0.5	0.3*	1.1	0.6	0.5*	1.4	0.6	0.8*
<i>b) 15-60 cm Upper Slope Area</i>												
NO ₃ ⁻ -N (kg ha ⁻¹)	84	56	29	71	48	23	38	28	9	38	22	16
SO ₄ -S (kg ha ⁻¹)	68	89	-21	78	36	43*	203	202	1	182	252	-69
CaCO ₃ (%)	10.9	13.2	-2.3	13.4	10.8	2.6	12.1	12.7	-0.6	12.2	12.2	0.0
Total Carbon (%)	4.0	3.9	0.1	3.9	4.0	-0.1	3.4	3.1	0.3**	3.5	3.1	0.4
Total Organic Carbon (%)	2.7	2.4	0.3	2.3	2.7	-0.4	2.0	1.5	0.5*	2.1	1.6	0.5

^a Electrical Conductivity based on a 1:1 soil:water extraction

^b Flax in Block A and Peas in Block B

^c U1A = Addition treatment pair 1; UID = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table 3.4 Fall soil fertility results for the lower slope area of FE1 in 2003

Nutrient	Block A						Block B					
	L1R ^b	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 0-15 cm Lower Slope Area												
NO ₃ ⁻ -N (kg ha ⁻¹)	2	6	-4***	3	8	-5	12	19	-7*	28	15	13*
P-Olsen (kg ha ⁻¹)	11	49	-38***	9	34	-25**	11	31	-20	20	38	18
K (kg ha ⁻¹)	585	1510	-925***	564	1149	-584***	587	1192	-605	703	1174	-471
SO ₄ -S (kg ha ⁻¹)	134	134	0	134	134	0	134	134	0	134	134	0
CaCO ₃ (%)	0.2	0.2	0.0	1.4	0.2	1.2	8.6	2.8	5.8***	5.6	3.2	2.4*
Total Carbon (%)	2.4	8.3	-5.9**	2.6	6.9	-4.3**	5.3	5.4	-0.1	6.4	6.0	0.4
Total Organic Carbon (%)	2.3	8.2	-5.9**	2.4	6.9	-4.5**	4.2	5.1	-0.9*	5.8	5.6	0.1*
EC ^a (mS/cm)	1.5	1.1	0.34	1.4	1.1	0.2	2.4	2.3	0.1**	3.4	2.4	1.03*
b) 15-60 cm Lower Slope Area												
NO ₃ ⁻ -N (kg ha ⁻¹)	8	16	-8**	8	23	-15	15	36	-22	35	37	-2
P-Olsen (kg ha ⁻¹)	11	31	-20**	13	25	-12**	16	27	-11	25	31	-6
K (kg ha ⁻¹)	1042	1369	-327**	883	1236	-353**	1205	1427	-222	1263	1243	-20
SO ₄ -S (kg ha ⁻¹)	403	403	0	403	403	0	403	403	0	403	403	0
CaCO ₃ (%)	11.2	5.9	5.3*	13.0	5.2	7.8*	12.8	6.3	6.5	12.8	8.9	3.9
Total Carbon (%)	2.9	4.1	-1.2**	3.3	3.6	-0.3	3.9	4.9	-1.0***	4.7	4.7	0
Total Organic Carbon (%)	1.6	3.4	-1.8***	1.8	3.0	-1.2	2.4	4.2	-1.8*	3.1	3.7	-0.6

^a Electrical Conductivity based on a 1:1 soil:water extraction

^bU1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table 3.5 Spring soil fertility results for the upper slope area of FE2 in 2004

	NO ₃ ⁻ -N ² (kg ha ⁻¹)	P-Olsen (kg ha ⁻¹)	K (kg ha ⁻¹)	SO ₄ -S (kg ha ⁻¹)	Total Carbon (%)	Inorganic Carbon (%)	Total Nitrogen (%)	pH	EC ¹ (mS cm ⁻¹)
<i>a) 0-15 cm Upper</i>									
<i>Slope Area</i>									
Addition	34a	54a	965a	43a	4.8	0.9b	0.3a	7.7c	0.7a
Control	19b	25b	701b	4b	4.2	1.4a	0.3b	7.8b	0.5b
Disturbed	21b	25b	668b	4b	4.5	1.8a	0.3b	7.8a	0.5b
<i>b) 15-30 cm Upper</i>									
<i>Slope Area</i>									
Addition	24a	-	-	6.9	4.5a	0.2	-	-	-
Control	12b	-	-	10.5	4.0b	0.1	-	-	-
Disturbed	11b	-	-	14.6	4.5a	0.1	-	-	-

¹Electrical Conductivity based on a 1:1 soil:water extraction

a-c Mean values followed by the same letter (within columns) are not significantly different based on an LSMean comparison at $P < 0.01$

Note: Values indicate the means of 9 replicates

3.4.1.3 Microbial Biomass Carbon (MBC). The soil microbial biomass carbon data is referring to the organic carbon released upon chloroform fumigation and not the corrected biomass (Jenkinson et al. 2003). These numbers provide an estimation of the microbial pool indicative of changes in soil organic matter and they might be useful for interpreting soil gas flux results.

In FE1, the addition treatment soil contained 60% more MBC on average than the disturbed or control treatments (Table 3.6). In year two, only two dates were sampled early in the sampling period. No clear trend was observed for microbial biomass C.

In the lower slope position in 2003 of FE1, significant reductions in microbial biomass C in the removal treatment occurred on all dates in Block A (Table 3.7). In 2004, a similar pattern was observed.

Microbial biomass was analyzed from two sampling dates for FE2. Soil addition or soil disturbance did not significantly affect soil MBC (Table 3.8).

Table 3.6 Microbial biomass carbon results in the upper slope area of FE1

Date	Block A						Block B					
	U1A	UID	Δ	U2A	U2C	Δ	U1A	UID	Δ	U2A	U2C	Δ
	$\mu\text{g g soil}^{-1}$			$\mu\text{g g soil}^{-1}$			$\mu\text{g g soil}^{-1}$			$\mu\text{g g soil}^{-1}$		
a) 2003												
27-May	338.6	180.2	158.5	357.9	193.2	164.7**	323.0	183.9	139.0**	297.3	196.6	100.7**
17-Jun	242.7	212.8	29.9	342.6	216.2	126.4**	309.8	174.0	135.9***	255.9	165.1	90.9*
3-Jul	178.3	174.2	4.1	230.5	169.0	61.5*	266.2	199.1	67.1**	344.2	197.0	147.2**
31-Jul	236.4	205.6	30.8	223.6	209.6	14.0	281.4	207.2	74.2	260.4	301.3	-40.9
17-Sep	237.4	247.3	-9.9	270.2	240.0	30.2	272.4	232.3	40.1	272.9	192.2	80.7**
CV %	32	21		25	20		16	21		16	24	
b) 2004												
27-May	311.2	286.5	24.7	283.9	296.4	-12.6	337.6	232.6	105.0	357.6	204.6	153.0*
9-Jun	236.7	272.4	-35.7	247.0	240.3	6.7	271.9	204.4	67.4**	297.5	289.0	8.4
CV %	20	10		24	23		18	14		15	33	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table 3.7 Microbial biomass carbon results in the lower slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
			$\mu\text{g g soil}^{-1}$						$\mu\text{g g soil}^{-1}$			
a) 2003												
27-May	92.4	578.4	-485.9*	217.9	329.9	-112.0**	360.7	371.8	-11.1	343.9	504.7	-160.8
17-Jun	127.1	314.1	-187.0***	182.8	307.2	-124.4**	222.7	336.5	-113.8	275.4	314.1	-38.7
3-Jul	121.5	492.1	-370.6**	177.8	454.6	-276.8**	266.9	362.1	-95.3	298.9	327.1	-28.2
31-Jul	130.4	460.9	-330.5**	161.5	432.3	-270.9**	278.4	342.0	-63.6	319.7	452.5	-132.8
17-Sep	72.4	436.3	-363.8**	162.5	536.9	-374.4**	227.3	414.4	-187.1*	256.4	517.5	-261.2
CV %	30	28		22	25		24	23		20	35	
b) 2004												
27-May	165.4	581.3	-415.9**	344.4	603.6	-259.2	426.1	438.0	-11.9	235.9	515.4	-279.4
9-Jun	126.9	594.3	-467.4**	164.8	491.9	-327.1	333.7	370.4	-36.7	278.4	465.7	-187.3
CV %	28	16			74	27	39	52		66	46	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table 3.8 Microbial biomass carbon results ($\mu\text{g g soil}^{-1}$) for two sampling dates in the upper slope area of FE2 in 2004

Treatment	27-May	9-Jun	CV %
	Carbon		
Addition	243.7	241.0	27
Control	562.4	274.3	195
Disturbed	294.5	259.8	39

CV% for each treatment was calculated by dividing the standard deviation units by the mean using all reps and all dates

3.4.2 Gas Flux

Soil addition did not have a consistent positive or negative effect on gas flux in the field. The gas flux results were variable and few significant treatment differences occurred. This may be the result of having limited data for individual dates in FE1 because only a small number of replications of each pair were used. Also, because gas emissions are highly variable in space and time (Corre et al. 1996; Bremner 1997; Smith et al. 2003), sampling a small number of replications made it difficult to capture any treatment effects. Coefficients of variation (calculated for each treatment by dividing the standard deviation units by the mean using all reps and dates) were on average 100%, 250%, and 500% for CO₂, CH₄ and N₂O, respectively.

3.4.2.1 Carbon Dioxide. Soil respiration rates (CO₂ flux) increased as the growing season progressed and soil temperatures rose; all treatments followed the same general pattern. In the upper slope area in 2003 of FE1, significant differences in soil respiration rates occurred on four of the nine sampling dates where the control treatment (U2C) emitted higher levels of CO₂ than the addition treatment (U2A) (Figure 3.6). In 2004, the opposite was found; significantly higher respiration rates were observed from the

addition treatment on two sampling dates in both blocks. A sharp peak in CO₂ flux was observed on July 15 (Day 195) for all treatments which corresponds with a rainfall event and an increase in soil moisture and temperature. When the soil CO₂ flux was calculated as a cumulative rate over each sampling period, treatment effects were not found to be statistically significant in either Block A or B; however, the data suggests the control treatment soil may have emitted greater cumulative soil CO₂ flux than the addition treatments (Appendix N.1).

In the lower slope area, soil respiration rates varied greatly between dates in 2003 and in Block A in 2004. Respiration generally increased as the growing season progressed with a peak also occurring on July 15 (Day 195) as was observed in the upper slope. In Block A, the disturbed (L1D) and control (L2C) treatments showed higher soil CO₂ flux throughout both sampling periods with significantly higher rates occurring on three dates in 2003 and on five dates in 2004 (Figure 3.7). In Block B, similar flux rates were observed for all treatments in both years except for on two dates in 2003 where the removal (L1R) treatment showed significantly greater soil CO₂ flux than the disturbed treatment. Cumulative CO₂ flux results indicate the removal treatment soil emitted significantly lower emissions than the control soil in 2003 and the disturbed soil in 2004 in Block A. In Block B the opposite was found where the data suggests that the removal treatment emitted greater cumulative soil CO₂ flux than the disturbed and control treatments; however, the comparison was only statistically significant when compared to the control soil in 2004 (Appendix N.2).

On June 25 (Day 176), 2003, the mean rate of soil respiration for the removal treatment (L2R) (399 mg CO₂-C m⁻² hr⁻¹) was three times higher than the average rate

across all treatments. This value appears to be anomalous, and in reviewing the literature pertaining to CO₂ evolution from soils, this number does exceed the expected range of emission (Bajracharya et al. 2000). This high value is attributed to the 60 min sample vial showing CO₂ concentration higher by a factor of 10.

In FE2, soil respiration remained below 20 mg CO₂-C m⁻² hr⁻¹ for the early part of the growing season with a large rise in respiration rates on July 15 (Day 195) and August 16 (Day 227) (Figure 3.8). The addition and disturbed treatment soils of FE2 emitted greater rates of CO₂ compared with the control on April 21 (Day 111) and May 17 (Day 137). On July 15 (Day 195), soil respiration in the control significantly exceeded the other treatments by a factor of three. The control soil emitted significantly higher cumulative flux over the 118-day sampling period in 2004 compared with both the addition and disturbed treatments (Appendix N.3).

3.4.2.2 Methane. When both years and both blocks are considered, methane emissions generally remained below or near zero for the majority of sampling dates in 2003 and 2004 in the upper slope area. In 2003 a positive emission occurred for most treatments early in the growing season on May 27 (Julian Day 147) and in the fall (September 17 – Day 259). Significant treatment effects were observed on three sampling dates in both 2003 and 2004 (Figure 3.9). In most cases, methane consumption was lower in the addition treatment compared with the disturbed or control treatments. When the methane flux was calculated as a cumulative rate over the each sampling period, significant treatment effects were found in Block A in 2003 and 2004 and in Block B in 2003; however, the treatment effect is variable (Appendix N.1).

In the lower slope area, methane emissions remained near zero for most sampling dates. The removal treatments showed lower methane consumption rates compared with the disturbed and control treatments with significant differences occurring on two sampling dates in 2003 and one sampling date in 2004 (Figure 3.12).

In FE2, methane emissions also remained near or below zero for most dates. Early in the growing season, the addition and disturbed treatment soils consumed higher rates of methane compared with the control on April 21 (Day 111) and May 17 (Day 137) (Figure 3.11). However, on Aug 16 (Day 227), the addition treatment emitted $3.59 \mu\text{g CH}_4\text{-C m}^{-2} \text{ hr}^{-1}$ while the disturbed and control treatment soils consumed methane.

3.4.2.3 Nitrous Oxide. When looking at nitrous oxide flux across both sampling periods, the flux for all treatments generally remained below $20 \mu\text{g N}_2\text{O-N m}^{-2} \text{ hr}^{-1}$ in year one but greatly exceeded this level in year two. In the upper slope area, no clear and consistent significant treatment effects were observed for soil N_2O emissions in either 2003 or 2004. However, when looking at individual dates, significant differences were found on two dates in 2003 and on three dates in 2004 (Figure 3.12). Although some significant difference was found, there was no consistent treatment effect. A slight burst in soil N_2O flux was observed on June 9, 2004 in both blocks. This corresponded with a rainfall event during that time (Figure 3.4) as well as canola seeding and fertilization which occurred on May 21 and May 28, 2004 in Block B and Block A, respectively. Cumulative soil N_2O emission rates were significantly greater in the addition treatment compared with the disturbed treatment in 2003 (Appendix N.1).

The effect of removing soil on soil N_2O emissions was also variable (Figure 3.13). No significant treatment differences were found in Block B. On April 21, 2004

the disturbed treatment emitted a slightly higher rate of soil N₂O in Block A. In Block B, the removal treatment showed significantly higher rates of soil N₂O emission on two of the seven sampling dates. When cumulative soil N₂O emission rates are considered, the removal treatment showed significantly higher emission rates than the disturbed treatment in 2003 in Block A and in 2004 in Block B. Cumulative soil N₂O emission rates were higher in the removal treatment compared with the control treatment in Block B in 2004 (Appendix N.2).

Soil nitrous oxide emissions were quite low in the upper slope area of FE2 in the early part of the sampling period. A burst of activity on June 9 (Figure 3.14) was also observed for this experiment where the addition and control treatments showed much higher emissions than the disturbed treatment; however, no significant differences between treatments were found for this or any other date.




Figure 3.6 Soil respiration (CO₂ flux) in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at P<0.10; ** Significant at P<0.05; *Significant at P<0.01; Values indicate the means of 3 replicates; significance symbols following differences are based on an analysis of log-transformed data but non-transformed means are presented (see Appendix G.1 for numeric values and statistical significance).**

¹U1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2




Figure 3.7 Soil respiration (CO₂ flux) in the lower slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at P<0.10; ** Significant at P<0.05; *Significant at P<0.01; Values indicate the means of 3 replicates; significance symbols following differences are based on an analysis of log-transformed data but non-transformed means are presented (see Appendix G.2 for numeric values and statistical significance).**

¹U1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

²This value exceeds the scale of the CO₂ flux for all other treatments and dates.




Figure 3.8 Soil respiration (CO₂ flux) in the upper slope area for FE2 in 2004. a-b Mean values followed by the same letter (within sampling dates) are not significantly different. LSMeans groupings are associated with significant treatment effects found using linear mixed ANOVA at P < 0.10; values indicate the means of 3 replicates (see Appendix G.7 for numeric values and statistical significance).




Figure 3.9 Soil methane flux in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004.*Significant at $P < 0.10$; ** Significant at $P < 0.05$; *Significant at $P < 0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on an analysis of log-transformed data but non-transformed means are presented (see Appendix G.3 for numeric values and statistical significance).**

¹U1A = Addition treatment pair 1; UID = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2




Figure 3.10 Soil methane flux in the lower slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at $P<0.10$; ** Significant at $P<0.05$; *Significant at $P<0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on an analysis of log-transformed data but non-transformed means are presented (see Appendix G.4 for numeric values and statistical significance).**

¹U1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2




Figure 3.11 Soil methane flux in the upper slope area for FE2 in 2004. a-b Mean values followed by the same letter (within sampling dates) are not significantly different. LSMeans groupings are associated with significant treatment effects found using linear mixed ANOVA at $P < 0.10$; values indicate the means of 3 replicates (see Appendix G.7 for numeric values and statistical significance).




Figure 3.12 Soil nitrous oxide flux in the upper slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at $P < 0.10$; ** Significant at $P < 0.05$; *Significant at $P < 0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on an analysis of log-transformed data but non-transformed means are presented (see Appendix G.5 for numeric values and statistical significance).**

¹U1A = Addition treatment pair 1; UID = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

²This value exceeds the scale of the N₂O flux for all other treatments and dates.




Figure 3.13 Soil nitrous oxide flux in the lower slope area for (a) Block A and (b) Block B in FE1 in 2003 and 2004. *Significant at $P < 0.10$; ** Significant at $P < 0.05$; *Significant at $P < 0.01$; Values indicate the means of 3 replicates; significance symbols following differences are based on an analysis of log-transformed data but non-transformed means are presented (see Appendix G.6 for numeric values and statistical significance).**

¹U1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

²This value exceeds the scale of the N_2O flux for all other treatments and dates.




Figure 3.14 Soil nitrous oxide flux in the upper slope area for FE2 in 2004. No significant treatment effects were observed based on LSmeans comparison using a linear mixed ANOVA at $P < 0.10$; values indicate the means of 3 replicates (see Appendix G.7 for numeric values and statistical significance).

3.4.3 Relationship between Flux and Depth of Added Soil/Depth of Removed Soil

3.4.3.1 Column Experiment Part 1. On average, the daily CO₂ flux results indicate the depth of soil addition does not affect soil respiration (Table 3.9 a). Significant treatment effects were observed on three of the twelve sampling days; however, no trend in treatment effect was found. Mean respiration rates for the 12 sampling days were 22.2, 30.5, 27.3 and 21.4 mg CO₂-C m⁻² hr⁻² for the control, 10U, 10D and 20D treatments, respectively.

The data suggests that soil CH₄ flux was consistently higher for the control treatment compared to the 20 cm addition treatment; however, this was only statistically

significant on one of the twelve sampling days (Table 3.9 b). The addition treatments showed significantly higher soil N₂O flux on three of the twelve sampling days; the depth of soil added did not consistently affect soil N₂O flux (Table 3.9 c). The quality of the soil did not influence soil greenhouse gas flux where the 10U and 10D treatments acted the same on all dates where significant treatment effects were found. Data was also analyzed on a flux per kg of soil basis and the same results were found. When the daily soil greenhouse gas flux measurements were calculated as a cumulative rate over the 30-day sampling period, significant treatment effects were found for cumulative soil CO₂ flux where the 10U treatment emitted the greatest soil CO₂ flux compared with all other treatments. The 10D, 20D and control treatments emitted similar rates based on Tukey's Least Significant Difference comparison (P<0.05) (Appendix O.1a).

3.4.3.2 Column Experiment Part 2. Mean daily soil CO₂ flux was five times higher in the surface treatment (control) than in either of the removal treatments (Table 3.10 a). The control treatment showed significantly higher (P<0.5) soil CO₂ flux compared with both removal treatments on the first nine sampling days but all treatments emitted similar rates thereafter. No significant differences between treatments were observed for cumulative soil CH₄ flux. The surface treatment (control treatment) also showed significantly higher (P<0.05) nitrous oxide flux compared with the removal treatments, with 33.1, 4.5, and 4.6 µg N₂O-N m⁻² h⁻¹ emitted from the surface on average for the control, -20 and -40 cm treatments, respectively (Table 3.10 c). When the daily soil greenhouse gas flux measurements were calculated as a cumulative rate over the 36-day sampling period, significant treatment effects were found for cumulative soil CO₂ and N₂O flux. The removal treatments emitted significantly lower soil CO₂ and N₂O flux

compared with the control treatment. (Appendix O.3). The depth of soil removal did not affect daily or cumulative gas flux.

Table 3.9 Soil gas flux for the soil addition growth chamber experiment (Column Experiment 1)

Treatment ¹	Sampling Day											
	1	2	3	4	5	6	7	8	9	10	11	12
a) CO₂ (mg CO₂-C m⁻² hr⁻¹)												
Control	40.3a	26.8	28.9	45.5	23.8b	30.5	24.3	4.3	9.4	14.5	16.9	19.3b
10U	9.5b	44.6	47.2	1.8	64.7a	41.9	35.5	15.1	19.9	24.8	25.4	25.9a
10D	12.2b	13.7	46.5	30.7	62.3a	31.2	31.2	12.7	20.4	28.1	27.9	27.6ab
20D	12.8b	11.1	25.7	7.2	22.0b	37.5	28.4	20.4	21.2	22.0	22.7	23.4ab
b) CH₄ (μg CH₄-C m⁻² hr⁻¹)												
Control	3.0	3.7	-28.8	4.6	8.6	-0.4	2.5	5.3	5.4a	5.5	1.5	-2.6
10U	4.3	3.8	-23.4	8.8	0.9	2.0	2.7	9.5	4.8ab	0.2	0.1	0.0
10D	2.5	2.4	-24.6	0.2	1.8	5.0	2.4	7.9	4.0ab	0.2	-3.0	-6.1
20D	1.8	0.5	-20.8	2.9	6.4	-0.9	1.3	1.6	0.2b	-1.2	-1.4	-1.5
c) N₂O (μg N₂O-N m⁻² hr⁻¹)												
Control	12.1	16.9b	-4.2	15.7	5.0	7.8	8.0	2.7	4.2	5.7b	5.8	5.8b
10U	21.4	65.5ab	24.4	1.3	34.6	23.1	17.5	12.4	36.3	60.2a	59.6	59.0a
10D	24.0	20.1b	30.3	69.9	32.7	28.4	26.7	17.5	31.7	46.0a	54.0	61.9a
20D	11.3	140.9a	72.5	2.6	30.3	41.9	28.8	16.2	36.0	55.9a	54.6	53.2a

a-b Mean values followed by the same letter (within columns) are not significantly different based on a Tukey's student minimum significant difference comparison at $P < 0.05$ based on log-transformed data but non-transformed means are presented

Note: values indicate the mean of three replicates

¹Control: upper slope soil only, no soil added; 10U: 10 cm upper slope soil added to the upper slope soil;

10D: 10 cm depression soil added to the upper slope soil; 20D: 20 cm depression soil added to the upper slope soil

Table 3.10 Soil gas flux for the soil removal growth chamber experiment (Column Experiment 2)

Treatment ¹	Sampling Day												
	1	2	3	4	5	6	7	8	9	10	11	12	13
a) CO₂ (mg CO₂-C m⁻² hr⁻¹)													
Control	28.6a	52.3a	54.3a	51.6a	63.8a	48.8a	56.3a	43.9a	42.4a	15.3	19.2	15.0	12.8
-20	8.8b	6.1b	6.4b	5.9b	8.1b	5.4b	14.1b	12.5b	12.8b	6.0	6.1	1.3	4.1
-40	3.5b	5.3b	8.9b	2.0b	4.3bb	6.8b	6.1b	3.9b	3.0b	13.9	17.1	11.6	20.5
b) CH₄ (μg CH₄-C m⁻² hr⁻¹)													
Control	-1.8	-0.4	3.3	11.8	10.0	7.0	20.2	6.7	6.4	1.1	2.1	-4.1	4.6
-20	-8.0	-1.9	0.5	-3.2	-0.5	-0.7	2.1	3.3	0.8	-4.0	3.0	-0.4	4.4
-40	-9.7	-4.0	-5.2	-3.4	1.1	-6.2	-1.0	-1.2	-3.6	1.8	16.2	11.4	7.8
c) N₂O (μg N₂O-N m⁻² hr⁻¹)													
Control	18.4a	71.5a	62.0a	38.8a	27.6a	42.4a	40.2a	39.5a	32.3	15.9	18.0	12.1	11.5
-20	8.4b	2.7b	4.0b	7.3b	5.0b	4.7b	9.3ab	8.6ab	2.3	0.5	1.6	0.7	3.3
-40	5.0b	2.0b	2.1b	2.4b	2.0b	1.5b	0.4b	1.2b	4.7	5.8	5.8	6.1	21.2

a-b Mean values followed by the same letter (within columns) are not significantly different based on a Tukey's student minimum significant difference comparison at $P < 0.05$ based on log-transformed data but non-transformed means are presented

Note: values indicate the mean of three replicates

¹Control: depression soil only, no soil removed; -20: 20 cm soil removed from the depression;
-40: 40 cm soil removed from the depression

3.5 Discussion

3.5.1 The Effects of Landscape Restoration on the Soil Environment

3.5.1.1 Soil Temperature and Moisture. In the upper slope area, landscape restoration was expected to increase soil temperature and moisture due to increased organic matter content and reduced plant residue cover resulting in decreased albedo and higher water holding capacity. Warm and moist soil conditions are favorable for microbial activity which in turn may affect gas flux from soil.

In the first year of the field experiments, a slight increase in soil temperature in the addition treatments compared with the control treatment was observed. The dark color of the added soil absorbed more solar radiation causing it to warm sooner in the season and reach warmer temperatures at depth. The eroded soils that comprised the disturbed and control treatments were much lighter in color and, therefore, would have reflected more radiation and stayed cooler longer. In addition to light colored surface soil, the control was a zero-till soil and the light colored plant residue on the soil surface increased the proportion of reflected radiation (Figure 3.1). These results are consistent with Malhi et al. (2001) who found that during the spring, soil temperatures at the same depths were lower for no-till versus conventional tillage systems. In year two, little differences in temperature were found between treatments in FE1. This is likely the result of all treatments being under zero-till cropping and, therefore, all treatments having a layer of standing stubble which increased the albedo in the addition and disturbed treatments.

The added topsoil increased the moisture holding potential of the soil in general. The results show that moisture content is greater in the spring following restoration

compared to the disturbed treatment; the magnitude of the difference is smaller when compared to the control treatment. This is especially important considering 2003 was a dry year at the MZTRA farm and any added moisture could be beneficial to crop growth. In year two when higher rainfall was observed over the growing season (Figure 3.4), there were more variable and less consistent moisture results when comparing the addition of soil to the control and disturbed treatments. In FE2, moisture was also greater on the upper slope after the addition of soil and this effect was most evident in the spring.

In the lower slope area, the effects of landscape restoration, specifically soil removal, on soil temperature were much more obvious. Lower slope temperatures were significantly higher for the removal treatments compared with the controls, most likely due to the dark soil surface absorbing more radiation. The grass residues on the controls may have reflected radiation and may act as insulation causing the soil to stay cooler longer. The same results are observed at depth. In the fall, the opposite is true; the removal treatments were cooler in September compared with the controls. Again, the layer of plant material on the soil surface may have acted like an insulating layer, causing the soil below to stay warmer longer.

When soil is removed from a riparian area, there are a few factors influencing the effect of soil removal on soil moisture. One factor is the physical effect of removing soil. Removing 20 cm of soil results in a new surface that is closer to the water table thereby increasing soil moisture (increase H₂O per volume of soil). Also, the area of removal might act as a catchment for runoff water and snow concentrating the total volume of water for a given area. A second factor includes the biologic effects of soil removal. In the control plots, drier soil conditions over summer and fall may be caused by greater

plant respiration and evapotranspiration. In the spring, there is little plant growth and little evapotranspiration resulting in little difference in soil moisture in the removal area compared with non-removal areas. As plants begin to take up water and transpire, the difference grows where the removal plots are expected to maintain higher moisture levels. Higher moisture was observed in the removal plots of Block B in 2003; however, there was no clear trend in Block A.

3.5.1.2 Soil Nutrients. Landscape restoration positively influenced soil fertility and CaCO_3 content in the surface soil based on fertility results from 2003. Increased fertility has important implications for crop and weed growth and also for soil gas flux.

It was hypothesized that restoration would increase the organic carbon of the topsoil; however, the increase that was found was only minor. Based on calculations using the measured TOC (%) in the upper slope and lower slope area treatments in FE1, and based on the assumption that 10 cm of soil was in fact added to the upper slope area addition plots, a 1.6% increase in TOC should have been observed (Appendix H). In FE1, the addition treatments showed 1.3% greater TOC in Block A compared with the control, and 0.8% greater TOC compared with the control in Block B. In FE2, the increase was also minor at 1.1%. The minor increase in TOC in Block B may be attributed by the fact that the actual depth of soil may not have been 10 cm once the soil had settled and been seeded. Also, some of the organic carbon may have been mineralized resulting from a “flush” of microbial CO_2 in response to heavy disking (Reicosky 1997b); however this was not observed statistically in the field or in the growth chamber. Finally, it is difficult to measure changes in soil organic matter over time and space due to the inherent variability of this soil property (Izaurrealde et al. 1997).

There was a significant increase in salt content in the addition treatments over the control in both field experiments; however, the increase was small and remained within tolerable levels (less than 3 mS cm^{-1}) for crop growth (Bower and Wilcox 1965). There was a concern with the potential for causing salinity in the upper slope area where soil was added because salinity had been indicated (Podolsky and Schindler 1993) in the areas from where soil was removed. The higher salinity level where soil was applied is likely only a short-term concern as salts are easily leached lower in the soil profile with the downward movement of water.

Landscape restoration reduced fertility in the removal areas. Since soil was removed from an area of accumulation of up to 0.75 m, it was expected that there would have been a large quantity of nutrients remaining after removing 20 cm of surface soil. The perennial vegetation and the adjacent wetland may influence the low reserve of nutrients at depth. An examination of soil fertility is necessary prior to removing the soil to ensure enough nutrients remain to achieve sustainable crop production (this is relevant to soil removal from cropland depressions).

3.5.1.3 Microbial Biomass Carbon. Adding organic rich topsoil was expected to increase the organic matter content, specifically organic carbon, of the soil and even though the fertility results show little increase in TOC, the soil microbial biomass carbon results better support this hypothesis. In the upper slope area, the addition treatments showed significantly greater soil microbial biomass carbon levels. The increase is somewhat short-lived, however, based on year two biomass numbers. There were too few sampling points from year two to make any strong conclusions.

In the lower slope area of FE1, it was expected that soil removal would remove much of the active organic fraction (i.e. the microbial population). The buried material exposed by soil removal would likely contain a smaller amount of active soil microbial biomass carbon because soil oxygen is reduced at depth and soil moisture and temperature are below optimum conditions for microbial activity (Lobb et al. 2002). This effect was observed in the removal treatments where the soil microbial biomass carbon levels were significantly lower. This helps explain the reduced gas flux from the removal treatment soil in the short-term but this result may only occur until microbial populations build up.

3.5.2 The Effects of Landscape Restoration on Soil Gas Flux

3.5.2.1 Carbon Dioxide. Soils with higher organic matter content generally show larger microbial populations that in turn can cause greater mineralization rates of organic nutrients such as carbon and nitrogen (Lobb et al. 2002; Pekrun et al. 2003). For this reason, adding organic rich topsoil to the previously eroded hilltop should have resulted in higher CO₂ flux. Tillage also affects the amount of carbon and nitrogen being cycled and stored in the soil system (Eghball et al. 1994; Reicosky 1997b) and therefore disking the addition plots should have caused a greater CO₂ flux to be observed. Tillage creates favorable conditions for microbial decomposition to take place. It improves aeration and temperature in cool, wet soils, or where soils are dry, it mixes drier surface soils with moister subsurface soils (Campbell et al. 1996). Increasing organic matter decomposition and microbial activity will increase the amount of carbon being mineralized and released as CO₂ (Pekrun et al. 2003; Reicosky 1997a). However, this was found only in FE2 and

only in the spring; in contrast, the control treatments showed greater flux in the later part of the growing season.

There is evidence that a large loss of soil CO₂ gas occurs immediately (less than 1 hour) following tillage with smaller differences observed 19 days after the tillage operation (Reicosky 1997b). Considering that gas flux was not sampled for four weeks following soil addition and disking in FE1 the treatment effect may have been missed. This implies that the effect of adding soil on soil respiration is possibly short-lived and may only occur in the first spring season following restoration or shortly after the restoration event. The objective of this study was to identify medium-term impacts of landscape restoration on greenhouse gas emissions and, therefore, CO₂ lost immediately following landscape restoration (i.e. in the short-term) may be overlooked.

The greater CO₂ flux exhibited in the control treatments of both experiments may be explained by the fact that zero-till soils can show higher respiration rates compared with conventional tillage soils (Reicosky 1997b); however, this is more common in regions experiencing higher annual precipitation and temperature. A second possible explanation is that the source of soil CO₂ gas emissions may be inorganic in nature resulting from the dissolution of CaCO₃ (Burton and Beauchamp 1994) in the control soils which contained higher CaCO₃ levels. A third possible explanation is the greater proportion of roots that exist below the surface of the zero-till control soils compared with the addition and disturbed treatments (that had been recently tilled) may be contributing to CO₂ flux because carbon dioxide is released by autotrophic root respiration (Smith et al. 2003).

The removal of soil does not appear to increase soil respiration but it either remains the same as non-removal treatments or is significantly reduced. Soil moisture and temperature are two key factors influencing soil respiration. The trend in these two parameters is generally opposite to that observed in the respiration data i.e. where soil moisture and temperature are higher for removal treatments, the respiration is lower. Therefore, there may be other factors influencing respiration such as substrate and/or root activity. The removal plots had lower soil organic C (this is true based on soil microbial biomass carbon results and soil fertility TOC results) and lacked the root system and organic residues found in the non-removal treatments. Carbon dioxide is released from soil organic matter both by heterotrophic respiration and by autotrophic root respiration (Smith et al. 2003). It has also been noted that 'spatial variations in N and C dynamics have been related to differences in vegetation' (Corre et al. 1996) where the vegetation influences N and C availability.

3.5.2.2 Methane. Soil methane emission and oxidation (or absorption) is influenced by soil redox potential, depth to water table, the vegetation present and soil texture (Smith et al. 2003) as well as the nitrogen dynamics in soil (Paul and Clark 1996). In the upper slope area where soils are generally drier, methane oxidation, carried out by methanotrophs, is more likely. This process requires the presence of NO_3^- and NO_2^- and it is inhibited by high levels of NH_4^+ (Paul and Clark 1996). In FE1 higher NO_3^- levels were observed, however, methane emissions were quite variable with the majority of the values lying near 0 (positive or negative 1). In FE2, significantly greater absorption of methane occurred in the addition and disturbed treatments on the first two sampling

dates. The lack of a consistent effect on methane indicates no major impact of landscape restoration on methane flux.

Methane emission (vs. absorption) is expected in the lower slope area where moisture may be higher favoring methanogens. When considering soil moisture data, Block B removal treatments had consistently higher soil moisture, which should translate into greater methane emissions or at least lower consumption rates. This was indeed the case. Because methane absorption by soil declines with soil removal and soil respiration declines or remains the same with soil removal, the reduction in soil carbon due to methane emission may be offset in the medium-term by reduced soil respiration. The amount of methane absorption by soils are likely small compared to total methane emissions from farms and a large increase in methane absorption by soil is needed to offset a small proportion of current emissions (Janzen et al. 1998). The magnitude of soil methane emission and absorption observed in this study were likely quite small compared with global agricultural emissions of methane.

3.5.2.3 Nitrous Oxide. The addition of soil did not cause any clear or consistent response in N_2O emissions. Nitrous oxide is released from the soil during the processes of denitrification and nitrification. Denitrification is the process whereby nitrate (NO_3^-) is sequentially reduced to nitrite (NO_2^-), nitric oxide (NO), nitrous oxide ($N_2O_{(g)}$) and finally dinitrogen gas ($N_{2(g)}$) (Havlin et al. 1998). Denitrification is a form of anaerobic bacterial respiration where NO_3^- is the electron acceptor in the absence of oxygen gas (O_2). Denitrification may occur in moist soils or in soils with anaerobic microsites (created where the 'respiratory consumption of O_2 in the soil by plant and soil microorganisms exceeds the rate of replenishment by diffusion from the atmosphere')

(Smith et al. 2003)). Incomplete denitrification results in nitrous oxide being emitted versus completing the sequence to emit dinitrogen. Nitrification is an aerobic process, however, when O₂ is limited, the nitrifying bacteria can use nitrite as an electron acceptor and reduce it to NO and N₂O as with denitrification (Bremner 1997; Smith et al.2003). Neither of these processes were measured in any of the experiments in this study. It is assumed that denitrification is the primary process contributing to the N₂O emissions that were measured in the field and growth chamber experiments.

The rate of denitrification depends largely on the concentration of oxygen and nitrate in the soil as well as the amount of available carbon. Denitrifying bacteria are heterotrophic resulting in a high demand of available carbon as an energy source in the process (Paul and Clark 1996). Further to the amount of carbon, the quality and spatial distribution of carbon in the soil system are also key factors regulating denitrification rates (Van Cleemput 1998).

Although the data was variable, some significant increases in N₂O flux did occur in the addition treatment of FE1. This correlates well with the conditions that favor denitrification since the addition of soil resulted in a general increase in soil moisture, in nitrate content and soil biomass carbon. (Corre et al.1996; Bremner 1997).

No statistically significant differences or major trends occurred in FE2; however, there is an interesting pattern of nitrous oxide emissions. On June 9, 2003 a slight peak in N₂O flux was observed, especially in Block A. This occurred after spring seeding and the application of ammonium-nitrate fertilizer with the seed (on May 21) and therefore the N₂O flux may be associated with the nitrogen fertilization practice. In 2004, little N₂O was emitted early in the spring. Some research has found a significant loss of soil

nitrogen through bursts of N_2O flux during spring thaw (Goodroad and Keeney 1984; Burton and Beauchamp 1994). In this study increasingly higher emissions were observed as the season progressed with a peak emission occurring in all treatments on June 9 (Day 159), 2004. Soil flux was measured on April 21 (Day 111) and May 17 (Day 137) but these dates may not have captured a spring thaw event. The higher emissions in June may be explained by the warming of the soil as the air temperature increased resulting in greater denitrifier activity and also by a rainfall event occurring on June 4.

Assuming that denitrification is the predominant process resulting in the N_2O emissions that were observed in this study, greater differences between treatments were expected from both field experiments based on the large increase in soil nitrate content following the addition of soil. The large increase in soil nitrate content in the addition treatments should have resulted in larger increases in soil N_2O emissions than those that were observed since soil nitrate content is one of limiting factors in denitrification (Drury et al. 1998).

In the lower slope area, higher denitrification rates were expected resulting from higher soil moisture contents; however, most treatments emitted similar rates. The low levels of nitrate in the removal treatments may partly explain the lack of high N_2O flux. The removal treatments also showed significantly lower soil microbial biomass carbon and showed a reduction in organic carbon which may also explain the lack of high N_2O flux. Carbon substrate and organic carbon availability are two other limiting factors affecting biological denitrification (Drury et al. 1998) (assuming this is the process responsible for the N_2O emissions observed in this study).

3.5.3 Relationship Between Flux and Depth of Added Soil/Depth of Removed Soil

3.5.3.1 Column Experiment Part 1: Soil Addition. In the controlled growth chamber experiment, the addition of soil did significantly increased soil CO₂ flux from the surface compared with the control on one of the twelve sampling dates and although the differences were not significant, there is some indication that greater cumulative nitrous oxide emissions occurred in the addition treatments compared with the control. The soil nitrate results from the field study showed that soil nitrate was significantly greater where soil was added compared with the control. If it is assumed that soil nitrate and soil microbial biomass carbon also increased in the addition treatments of the growth chamber addition experiment, then it is expected that N₂O emissions would be greater following soil addition especially if soil moisture and temperature conditions were favorable. The soil moisture of the column soil was maintained at or near field capacity and the growth chambers were maintained at 25°C; therefore, with favorable conditions existing in the addition treatments, the high N₂O emissions are warranted. The lack of statistical significance may be are result of the high variability found within treatments. The results provided by this experiment show that the depth of added soil will not influence the flux of gas from the upper slope restored soil, however soil addition may increase N₂O emissions in general if soil moisture conditions are favorable.

3.5.3.2 Column Experiment Part 2: Soil Removal. The depth of soil removal decreased the GHG emissions from the lower slope area. Significantly lower cumulative respiration rates were observed from both removal treatments compared with the control. This is consistent with field observations in 2004 resulting from lower organic carbon levels at depth, which in turn would show lower microbial activity and less substrate and

thus lower respiration (Lind and Eiland 1989; Lobb et al. 2002). The same reasoning explains the lower cumulative N₂O flux. Surface soil has a higher organic carbon content which likely supports a more active microbial population, including denitrifiers. The increased microbial activity is likely to increase the rate of O₂ consumption and therefore increase the likelihood and frequency of anaerobic (or microanaerobic) conditions. Substrate at the surface may also be of better quality and higher concentration.

3.5.4 A Comparison of Field Experiment and Growth Chamber Experiment Findings

The field and growth chamber experiments were complimentary and both focused on evaluating the effects of landscape restoration, including soil addition and soil removal, on greenhouse gas flux from soil. The time scale that was captured by each set of experiments, however, was different. The field experiments measured soil greenhouse gas emissions in the medium-term following the landscape restoration practice while the time interval captured by the growth chamber experiments was short-term. The time scale that was captured by each set of experiments has implications for interpreting and comparing the results from each. For example, in the short-term following a soil disturbance (such as the addition of soil), a de-gasing may occur resulting in higher gas emissions compared with the soils that did not experience the disturbance.

When the field experiments are considered, there was little significant impact of adding and removing soil on soil greenhouse gas flux. In most comparisons over each of the two sampling periods, most treatments acted the same, and when statistically significant differences were found between treatments, the treatment effect was often variable. These results imply there is little negative impact of landscape restoration on

greenhouse gas emissions from soil in the medium-term. The short-term effects of landscape restoration were not captured in the field because flux measurements did not begin until one month after the disturbance event. The soil column experiments were useful because they provided information about the short-term effects of landscape restoration. Based on the soil column experiment findings, the addition of soil does not effect soil respiration or methane flux but there is an increase in soil nitrous oxide flux approximately 30 days after the disturbance occurs. The removal experiment using columns clearly showed that soil respiration and soil nitrous oxide flux is significantly reduced following the removal of soil but that this effect is short-lived and microbial populations are able to rebound and begin emitting similar levels as the control after only three weeks following the disturbance.

The magnitude of daily gas emissions were similar in the field and growth chamber experiments even though the time scale was different. This is an important observation indicating that higher emissions are not necessarily observed immediately after the disturbance event associated with landscape restoration but that the treatment effect is simply more pronounced.

3.6 Summary and Conclusions

In general, no clear and consistent significant effects of soil addition or removal on greenhouse gas flux were observed in the field. This may be the result of a small number of sampling points in space and time, and because of the high variability of greenhouse gas emissions from soil. Soil removal did reduce CO₂ emissions from soil in the first year following soil removal. In the growth chamber, the depth of soil added or

removed did not affect gas flux although soil removal did decrease soil respiration and nitrous oxide emissions in the first three weeks following the disturbance.

Although the results presented here show that landscape restoration did not affect greenhouse gas emissions from soil, some changes in the soil environment were observed following landscape restoration. Increases in soil nitrate and soil microbial biomass carbon following landscape restoration were observed. Because these parameters play a significant role in regulating gas flux, it can be assumed that if landscape restoration will influence the soil nitrate and soil microbial biomass carbon, it will also influence soil gas flux.

This study focused primarily on measuring greenhouse gas emissions in the medium-term (over two growing seasons). Monitoring gas emissions immediately following restoration (especially CO₂) may provide a better indication of short-term effects of landscape restoration because it is at the time of moving soil that a major release of soil organic carbon is expected due to soil disturbance, aeration and organic carbon mineralization. Also, to better understand the long-term effects of restoration, it is necessary to continue monitoring the effects of soil movement for years after restoration took place with a focus primarily on the early spring, and rainfall events when soil moisture is high. This is often difficult due to the unpredictable nature of weather, such as temperature fluctuations in the spring season, sporadic rainfall events throughout the growing season, and excessively high or low moisture conditions.

Finally, because this study was the first of its kind and it measured a broad range of variables, it was difficult to compare the results found here with those from published work investigating the effects of soil management on similar parameters. Future research

focusing in greater detail on landscape restoration effects on only one or two of the variables measured here is needed.

4.0 IMPACTS OF LANDSCAPE RESTORATION ON PLANT SPECIES AND ABUNDANCE

4.1 Abstract

The objective of this study was to look at the effects of landscape restoration on plant species and abundance by removing soil from the lower slope area of the landscape and adding it to the eroded upper slope area. Two field experiments were carried out over two growing seasons where plant emergence, including crop and weed, was measured and the weed species present were observed. A growth chamber experiment characterized the viability of weed seeds within the soil profile and the species present. Where soil was added, pea crop yield improved following the addition of soil however flax yield was unaffected and canola yields declined. Weed emergence increased in the addition treatments in the first year following restoration and the number of species present only increased slightly. The weed seed bank findings indicate the most viable weed seeds exist in the surface soil (top 5 cm) with similar species present as those found in the field site. Increased herbicide use may be required to control the increased weed pressures in the first year following the restoration practice. Using competitive crops is recommended to help out-compete weeds. Removing greater than 20 cm of soil from smaller areas to dilute the weed seed bank is also recommended.

4.2 Introduction

In agricultural systems, weeds are a common pest contributing to crop yield loss. The weed seed bank in soil is a persistent source of viable seeds available for successful

weed seedling recruitment (Benoit et al. 1992). Early work studying seed banks in soil suggests that enormous numbers of viable seeds exist in arable soils and wetlands, and that, although most buried seeds die within a few years, significant numbers of seeds from some species can survive for decades (Cavers 1995). The success of weed seedling recruitment will then depend on a number of factors including the depth of the seed within the soil profile, soil moisture and temperature conditions and biotic and abiotic variables that influence both of these factors including soil type and management (Boyd and Van Acker 2003).

Tillage is one land management practice that influences the soil seed bank and weed seedling recruitment. Tillage can affect seed placement within the soil (Bullied et al. 2003). It can bury seeds shed on the soil surface preventing germination and can also bring seeds closer to the surface where germination is more likely. Tillage erosion is common in topographically complex landscapes and can influence the spatial variability of the soil seed bank both vertically and laterally. Tillage erosion redistributes soil and thus weed seeds within the landscape resulting in a loss of soil from hilltops and an accumulation of soil in lower slope areas and depressions; the accumulation of soil may occur within the cropland or at the field edge adjacent to wetlands. The seed bank in soils where tillage erosion has occurred may, therefore, show a greater proportion of buried seeds deeper in the profile. Although Cavers (1995) concluded that most buried seeds die within a few years, he also states that seeds buried at greater depths tend to remain dormant longer. These dormant seeds generally don't germinate successfully and emerge as seedlings, however, due to their proximity to the soil surface.

Another land management practice that may influence the soil seed bank and weed seedling recruitment is landscape restoration. Landscape restoration is an innovative practice aimed to improve landscapes degraded by soil erosion. Landscape restoration consists of removing soil accumulated in lower slope areas and depressions and applying it to hilltops. There are two main implications of landscape restoration on the soil weed seed bank. First, seeds within the soil profile will be brought to the soil surface and be allowed to germinate and emerge on the restored hilltops. Second, seeds at depth may be exposed to the appropriate microsite conditions once soil is removed also allowing germination and emergence in the lower slope areas and depressions. Knowledge of the weed seed bank can, therefore, help identify future weed problems (Wiles and Schweizer 2002) in eroded croplands where landscape restoration is practiced.

Some work has looked at estimating the maximum depth of emergence and found that generally, emergence decreases with depth of burial but this does vary with weed species (du Croix Sissons et al. 2000; Benvenuti et al. 2001; Boyd and Van Acker 2003). du Croix Sissons et al. (2000) found that the mean minimum and maximum emergence depths were 1.2 and 4.2 cm, respectively, for four common weed species found in the northern Great Plains. Benvenuti et al. (2001) looked at the effect of seed burial depth on seedling emergence rate of 20 weed species and found that 10 cm was the maximum depth from which some weed species could emerge. The information obtained from these studies is useful; however, it does not provide information about the viability of weed seeds with depth. Information about seed bank viability within the soil profile would help identify the potential impacts of landscape restoration on weed emergence.

The objective of this study was to identify the impact of landscape restoration on weed emergence and weed species in the restored upper slope area and in the lower slope area where soil was removed. A growth chamber experiment looking at the weed seed bank within the soil profile was completed to complement the field study. The objectives of this complementary experiment were to determine the viability of the seed bank within the soil profile in the zones of accumulated soil, including identifying the species present, and to evaluate the potential for plant establishment in the removal and addition zones based on seed bank findings. The findings from this seed bank experiment were compared with the field experiment findings.

4.3 Materials and Methods

4.3.1 Field Experiment

Two field experiments, named Field Experiment 1 (FE1) and Field Experiment 2 (FE2), were established at the Manitoba Zero Tillage Research Association (MZTRA) farm in the spring of 2003 and the fall of 2003, respectively. This site has been under high-disturbance zero-till crop production since 1993. Prior to 1993, the soil had been intensively tilled resulting in the loss of topsoil from upper slope areas and the accumulation of soil in lower slope areas; the variation in topsoil depth within the landscape is still visible. The field experiments consisted of three main treatments in both the upper and lower slope areas. In the upper slope, the treatments included an addition treatment, disturbed treatment and control; in the lower slope the treatments included a removal treatment, disturbed treatment and a control (Table 3.2). FE1 was replicated in two adjacent fields under different crop rotations (Block A and Block B)

while FE2 was established in Block B using a slightly different experimental design. In 2003, FE1 was seeded to flax in Block A and peas in Block B. In 2004, both experiments were seeded to canola. In 2004, a problem occurred with the seeder causing large portions of the field to show little to no emergence; FE2 was greatly impacted by this. A more complete description of the study site, field experimental design and treatments is provided in Section 3.3.2.

4.3.1.1 Plant Measurements. Ten to fourteen days after crop emergence, plant counts were taken from three one-meter lengths along the seed row. Weed species plant counts were taken following herbicide application within 1 m² quadrats in each plot. Herbicide use only occurred in the upper slope area in both field experiments. Herbicide application followed general land management practices typical at the MZTRA farm (Section 3.3.2.3). Plant emergence levels and plant species were also measured in 0.25 m² quadrats in each of the removal plots of the lower slope area.

Above ground plant samples were harvested at crop maturity from 1 m² quadrats from each plot in the upper slope area in FE1 in 2003. A one-meter by four-row crop and weed sample was harvested from each plot in both FE1 and FE2 in 2004. Following harvest, samples were air dried at 25 °C, threshed and weighed for grain yield and total biomass yields (kg ha⁻¹). A 0.25 m² sample of above ground biomass was harvested from the lower slope area treatments in both years. Samples were dried and weighed to determine total above ground biomass (g m⁻²).

4.3.2 Weed Seed Bank Experiment (Growth Chamber)

A weed seed bank experiment was carried out to characterize the size and species composition of the seed bank within the profile of the accumulated soil and to evaluate

the potential for plant establishment in the removal and restoration areas based on seed bank findings.

4.3.2.1 Soil Collection and Preparation. Soil was collected from the MZTRA farm within the same area as the soil removal treatments for FE1 described in Section 3.3.2. This soil was situated within a vegetative buffer strip between the cropland and a deep permanent wetland. Soil was taken in November of 2002 in 75-cm by 10-cm layers in 5-cm depth increments to a depth of 25 cm (Figures 4.1 and 4.2). This depth was chosen based on the maximum depth of soil removed being 20 cm and, therefore, allowing the underlying 20-25 cm soil depth to be studied also. This was replicated three times at each of three locations spaced approximately 100 m apart within the vegetative buffer strip. The experimental design was a randomized complete block design where site was the block factor and was random; depth was considered as the treatment. Site was specified as random because it was considered as a location and each site was predicted to act differently due to a potential moisture and salinity gradient existing along the edge of the wetland. Soil moisture and salinity was not measured prior to and following the sampling of soil for this experiment.




Figure 4.1 The excavation area for one replicate of each depth increment.



Figure 4.2 The soil profile within the excavation area.

The collected soil was taken to the lab where it was frozen to help break seed dormancy. Following thawing, each depth-increment sample was sieved, thoroughly mixed and sub-sampled. Gravimetric moisture content of each depth increment was measured. The following method was adapted from Cardina and Sparrow (1996). Soil (1.6-kg oven dry equivalent) was placed in a round mesh-lined trays 20.3 cm diameter lined with fine mesh and tamped to a bulk density of 1.2 g cm^{-3} and a depth of 4 cm. The trays were placed in larger plastic containers lined with silica sand to allow for bottom-fed watering. This was necessary to prevent surface crusting of the soil (Cardina and Sparrow 1996). The moisture content of the soil was maintained at field capacity to encourage germination of the entire seed bank. The trays were kept in the growth chamber set to 16/8 and 25/15 °C day/night hours and temperatures, respectively, and a relative humidity of 80%.

4.3.2.2 Plant Measurements. Plant emergence was monitored daily until emergence ceased, representing the first cycle of the study. Plant species in each tray were identified. Representative plants of each species were grown to full maturity to ensure correct species identification.

Following the first cycle, the trays were placed in the freezer at -20 °C for one month to further break dormancy of any remaining seeds, the soil from each tray was mixed and re-tamped and plant growth was monitored a second time. Because emergence during the second cycle was so low in comparison to the first cycle, only two cycles were completed.

4.3.3 Data Analysis

All statistical analyses were performed using SAS version 8 software (SAS Institute Inc. 2000). Where appropriate generalized linear and linear mixed models were used.

4.3.3.1 Field Data. A paired t-test was used to test for treatment differences for FE1. For field experiment 2 data were analyzed using a linear mixed model in SAS to test for treatment and replication effects. Least Square Means (LSMeans) were also calculated to compare mean significance groupings of treatments. The replication*treatment interaction was specified as the random statement. A probability level of $\alpha = 0.10$ was used for both field experiments.

4.3.3.2 Weed Seed Bank Study. A mixed linear model ANOVA tested for depth and site effects using a probability level of $\alpha = 0.05$. Plant emergence counts were $\log_{10}(X+1)$ transformed based on a significant test for homogeneity of error variances. LSMeans were calculated to compare mean significance groupings among depth treatments.

4.4 Results

4.4.1 Plant Parameters

4.4.1.1 Crop Emergence and Yield. In 2003 in FE1, no significant differences occurred between treatments for crop emergence in either blocks in the upper slope area (Table 4.1). The flax crop yield was significantly higher in the addition plots when compared to the disturbed treatment in Block A; however, no difference in yield occurred between the addition and control treatments where flax yields were 1771 and 1670 kg ha⁻¹,

respectively. In Block B, the addition treatment yielded significantly higher pea yields than the disturbed and control treatments.

In the upper slope area in 2004 the addition treatment showed significantly lower crop emergence compared with the disturbed treatment in Block A. In Block B, the emergence numbers were relatively uniform across treatments (Table 4.1). Canola yields were significantly lower in the addition treatment compared with the disturbed treatment in Block A. It should be noted that the quality of the canola was very poor following threshing. The canola was harvested prematurely resulting in seed that could not be threshed out completely. Also, sclerotia bodies were present in the threshed seed resulting in inaccurate estimates of seed weight and thus yield.

In FE2, irregular emergence patterns occurred throughout plots 1 to 12 due to a problem with the seeder that went undetected until long after the crop had emerged. As a result, there were noticeable differences in crop biomass and yield (Table 4.2; Appendix I). The mean crop emergence and yield numbers are, therefore, questionable. Also, the irregular emergence may have affected weed emergence and persistence since weeds tend to be more prolific in areas with low plant competition (Boyd and Van Acker 2004).

4.4.1.2 Plant and Species Abundance. In 2003 in the upper slope area, weed emergence was significantly higher in pair-two of Block B and consistently higher in all addition plots (Table 4.1). On average, five weed species were present in all treatments. In 2004, no significant differences between treatments occurred for weed emergence levels; however, the addition plots consistently showed two more species compared with the other treatments. The number and type of plant species were consistent over both years

(Table 4.3). Weed emergence levels in all treatments declined in 2004 compared to 2003.

Table 4.1 Crop emergence, weed emergence, weed species numbers and crop yield for the upper slope area of FE1

	Block A						Block B					
	U1A ^b	UID	Δ	U2A	U2C	Δ	U1A	UID	Δ	U2A	U2C	Δ
a) 2003 Upper Slope Area												
Crop Emergence ^c (plants m ⁻²)	187	54	133	138	149	-11.1	36	18	18	43	25	18.4
Weed Emergence (plants m ⁻²)	96	60	36	88	32	55.7	73	63	10	97	43	54*
Weed Species (species m ⁻²)	5	5	0	5	4	2	5	5	0	5	4	2
Yield (kg ha ⁻¹) ^a	1407	1195	211**	1771	1670	101.7	3398	2186	1212**	2723	1480	1243***
b) 2004 Upper Slope Area												
Crop Emergence ^c (plants m ⁻²)	50	100	-50*	49	110	-61	41	38	3	59	61	-2
Weed Emergence (plants m ⁻²)	21	27	-6	37	18	19	34	42	-8	35	27	8
Weed Species (species m ⁻²)	4	2	2	4	2	2	6	4	2	6	4	2**
Yield (kg ha ⁻¹) ^a	1977	2860	-883**	2063	2517	-453.3	1677	1550	127	2217	2437	-220

^a *Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

^b U1A = Addition treatment, pair 1; UID = Disturbed treatment, pair 1; U2A = Addition treatment, pair 2; U2C = Control treatment, pair 2

^c Flax in Block A and Peas in Block B in 2003; Canola in Block A and B in 2004

Table 4.2 Crop emergence, weed density, weed species numbers and crop yield for FE2, 2004

Treatment	Crop emergence (plants m ⁻²)	Weed density (plants m ⁻²)	Weed Species (plants m ⁻²)	Yield (kg ha ⁻¹)
Addition	25	17	4	904b
Control	38	13	4	1180ab
Disturbed	23	12	4	1451a

Values indicate the means of 3 replicates;
a-b Mean values followed by the same letter (within columns) are not significantly different. LSMeans groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$.

Grassy (*Poa* spp.) species were the most abundant species observed in Block A and B in 2003 (Table 4.3). In Block A, 42 stinkweed (*Thlaspi arvense*) plants m⁻² were observed in the addition treatment of pair two. Volunteer canola (*Brassica napus*) and Canada thistle (*Cirsium arvense*) showed high emergence levels in Block B. Plant species composition was slightly different in the second year following restoration where volunteer flax (*Linum usitatissimum*) was the most abundant species in Block A and perennial sow-thistle (*Sonchus arvensis*) showed the highest emergence levels in Block B.

In the lower slope, plant emergence was monitored in the removal plots only; plant biomass was measured in all treatments. Plant emergence levels were greater in year 2. Above ground plant biomass was 151 and 114 g m⁻² in the disturbed and control treatments, respectively, in Block A in 2003 (Table 4.4). The removal treatments showed significantly lower plant biomass production than the disturbed and control treatments. In Block B, significantly higher plant biomass was observed in the disturbed treatment

compared with the removal treatment. In 2004, treatment differences followed a similar trend as in 2003. All removal treatments showed higher above ground plant biomass than the preceding year; however, this was not tested statistically.

The plant species observed in the lower slope area of FE1 were predominantly upland weed species; wetland species were not observed. In 2003, stinkweed (*Thlapsi arvense*) was the most abundant species in both blocks followed by Canada thistle (*Cirsium arvense*) (Table 4.5). In Block A, wild buckwheat (*Polygonum convolvulus*) was abundant in treatment L2R.

In FE2, weed emergence levels were slightly higher in the addition treatment, but it is difficult to say whether this is because of adding soil or because of seeder malfunction causing more favorable conditions for weed seeds to germinate and emerge. The number of species was the same for all treatments. The most abundant weed species observed in all treatments was scentless chamomile (*Matricaria perforata*).

The depressions of FE2 were not monitored for plant emergence and species abundance because during the early months of spring of 2004, standing water was present in all three depressions. During this time, cattails (*Typha* spp.) were observed in the most westerly of the three depressions. Standing water remained in this westerly depression for the majority of the growing season. Water remained in the removal side of the central and easterly depressions throughout June and July.

Table 4.3 Weed species (number of plants m⁻²) in the upper slope area of FE1 after crop emergence and following in-crop herbicide application in a) 2003 and b) 2004

Scientific name	Common name	Block A ^b				Block B			
		U1A ^a	UID	U2A	U2C	U1A	UID	U2A	U2C
a) 2003 Upper Slope Area									
<i>Amaranthus retroflexus</i>	redroot pigweed	-	14	1	1	2	1	1	1
<i>Brassica kaber</i>	wild mustard	7	3	6	1	-	-	-	-
<i>Brassica napus</i>	volunteer canola	-	-	-	-	22	17	45	5
<i>Cirsium arvense</i>	Canada thistle	4	1	3	4	11	12	7	18
<i>Descurainia sophia</i>	flixweed	-	-	-	-	-	-	-	3
<i>Poa</i> spp.	grass	14	34	28	23	25	24	31	19
<i>Polygonum convolvulus</i>	wild buckwheat	4	2	9	-	8	3	5	-
<i>Taraxacum officinale</i>	dandelion	2	-	-	1	1	-	1	-
<i>Thalpsi arvense</i>	stinkweed	5	6	42	3	3	6	-	
b) 2004 Upper Slope Area									
<i>Agropyron repens</i>	quack grass	-	-	-	-	2	1	3	2
<i>Amaranthus retroflexus</i>	redroot pigweed	-	-	-	-	3	-	2	-
<i>Chenopodium album</i>	lamb's-quarters	-	-	-	-	1	-	1	-
<i>Cirsium arvense</i>	Canada thistle	1	-	3	1	4	11	7	5
<i>Echinochloa crusgalli</i>	barnyard grass	-	-	-	-	2	-	-	-
<i>Linum usitatissimum</i> ,	volunteer flax	6	25	10	16	-	-	-	-
<i>Matricaria perforata</i>	scentless chamomile	-	2	-	-	-	3	1	2
<i>Medicago sativa</i>	volunteer alfalfa	-	-	-	-	-	-	1	-
<i>Poa</i> spp.	grass	-	-	-	-	-	-	-	-
<i>Polygonum convolvulus</i>	wild buckwheat	12	-	20	-	2	-	-	-
<i>Setaria viridis</i>	green foxtail	-	-	-	-	1	2	-	3
<i>Sonchus arvensis</i>	perennial sow-thistle	1	-	1	1	19	24	20	16
<i>Thalpsi arvense</i>	stinkweed	-	-	-	-	1	-	1	-

^a U1A = Addition treatment, pair 1; UID = Disturbed treatment, pair 1; U2A = Addition treatment, pair 2; U2C = Control treatment, pair 2

^b Flax in Block A and Peas in Block B in 2003; Canola in Blocks A and B in 2004

Table 4.4 Plant emergence levels, plant species numbers and above ground plant biomass in the lower slope area for FE1 for a) 2003 and b) 2004

	Block A						Block B					
	LIR ^b	L1D	Δ	L2R	L2C	Δ	LIR	L1D	Δ	L2R	L2C	Δ
a) 2003 Lower Slope Area												
Weed Emergence (plants m ⁻²)	22	-	-	41	-	-	31	-	-	18	-	-
Weed Species (species m ⁻²)	5	-	-	4	-	-	3	-	-	3	-	-
Above Ground Plant Biomass (g m ⁻²) ^{a,c}	33	151	-119**	40	114	-73*	46	146	-99**	43	74	-31
b) 2004 Lower Slope Area												
Weed Emergence (plants m ⁻²)	59	-	-	63	-	-	54	-	-	44	-	-
Weed Species (species m ⁻²)	4	-	-	4	-	-	5	-	-	4	-	-
Above Ground Plant Biomass (g m ⁻²) ^a	54	145	-92*	83	140	-57**	96	158	-63	95	70	25

^a *Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

^b L1R = removal treatment, pair 1; L1D = disturbed treatment, pair 1; L2R = removal treatment, pair 2; L2C = control treatment, pair 2;

^c Above ground plant biomass was measured in the control treatments however weed emergence and species levels were not measured in the control treatments due to the abundance of perennial grass species present

Table 4.5 Plant species (number of plants m⁻²) in the lower slope area of FE1

Scientific name	Common name	Block A		Block B	
		L1R ^a	L2R	L1R	L2R
a) 2003 Lower Slope Area					
<i>Amaranthus retroflexus</i>	redroot pigweed	2	-	1	1
<i>Brassica kaber</i>	wild mustard	1	3	-	-
<i>Cirsium arvense</i>	Canada thistle	10	7	16	9
<i>Linum usitatissimum</i>	flax	1	-	-	-
<i>Poa</i> spp.	grass	1	2	4	12
<i>Polygonum convolvulus</i>	wild buckwheat	3	12	1	1
<i>Populus tremuloides</i>	trembling aspen	1	1	-	-
<i>Taraxacum officinale</i>	dandelion	-	-	-	5
<i>Thalpsi arvense</i>	stinkweed	6	19	18	9
b) 2004 Lower Slope Area					
<i>Cirsium arvense</i>	Canada thistle	15	21	11	10
<i>Descurainia sophia</i>	flixweed	31	28	2	1
<i>Echinochloa crusgalli</i>	brome grass	1	1	4	0
<i>Plantago major</i>	plantain	-	-	1	-
<i>Polygonum lapathifolium</i>	lady's-thumb	2	1	-	-
<i>Setaria viridis</i>	green foxtail	-	-	1	2
<i>Sonchus arvensis</i>	perennial sow-thistle	9	13	31	26
<i>Thalpsi arvense</i>	stinkweed	-	-	1	5
<i>Trifolium repens</i>	white clover	-	-	1	-

^a L1R = removal treatment, pair 1; L1D = disturbed treatment, pair 1; L2R = removal treatment, pair 2; L2C = control treatment, pair 2;

Table 4.6 Weed species (plants m⁻²) in the upper slope area in FE2 measured after crop emergence and following in-crop herbicide application

		Addition	Control	Disturbed
<i>Cirsium arvense</i>	Canada thistle	-	2	1
<i>Echinochloa crusgalli</i>	barnyard grass	6	1	1
<i>Matricaria perforata</i>	scentless chamomile	6	4	5
<i>Polygonum convolvulus</i>	wild buckwheat	1	1	3
<i>Sonchus arvensis</i>	perennial sow-thistle	3	4	2

Note: values indicate the mean of nine replicates per treatment

4.4.2 Weed Seed Bank in the Soil Profile (Seed Bank Experiment)

Weed emergence was predominant in the 0-5 cm increment where 31 plants per m⁻² emerged on average. Mean total plant emergence was significantly ($p < 0.05$) greater

in the 0-5 cm soil increment compared to all other depths (Figure 4.3) indicating the most viable seed bank exists near the surface of the soil. Species abundance was also greatest at this depth (Figure 4.4). Total plants and species decreased significantly for soil samples below 10 cm. The soil seed bank is variable in space making it difficult to measure and quantify. This variability results in the high standard deviations observed within each site (Appendix J) and within each depth increment.

The most abundant weed species observed in this experiment was stinkweed (*Thalpsi arvense*). Other weedy species included sow-thistle, lamb's-quarters, shepard's purse and grasses including foxtail barley, green foxtail and an unknown grass (Table 4.7). None of the plant species identified were wetland or aquatic species. Many rhizomes and tubers were removed from the soil samples upon sieving thus resulting in an incomplete analysis of all the plants that may arise from restoring the landscape.




Figure 4.3 Mean total plant emergence by soil depth. Values represent the mean of three reps combined across three sites; error bars represent standard deviations. Letters indicate significant ($P < 0.05$) treatment differences based on LSMeans groupings using $\log_{10}(X+1)$ -transformed data, but non-transformed means are presented.




Figure 4.4 Mean total plant species by soil depth. Values represent the mean of three reps combined across three sites; error bars represent standard deviations. Letters indicate significant ($P < 0.05$) treatment differences based on LSMeans groupings.

Table 4.7 Weed species identified in the weed seed bank experiment

Scientific name	Common name	Total Plants
<i>Thlapsi arvensis</i>	stinkweed	174
<i>Sonchus arvensis</i>	perennial sow-thistle	75
<i>Chenopodium album</i>	lamb's-quarters	63
<i>Capsella bursa-pastoris</i>	sheperd's purse	46
<i>Poaceae</i> spp.	grass species*	43
<i>Brassica kaber</i>	wild mustard	30
<i>Polygonum convolvulus</i>	wild buckwheat	28
<i>Setaria viridis</i>	green foxtail	23
<i>Cirsium arvensis</i>	Canada thistle	13
<i>Silene noctiflora</i>	night-flowering catchfly	13
<i>Artemisia biennis</i> Willd.	biennial wormwood	11
<i>Poygonum lapathifolium</i>	smartweed	8
<i>Polygonum erectum</i>	knotweed spp.	5
<i>Plantago major</i>	plantain	5
<i>Erysimum cheiranthoides</i>	wormseed mustard	3
<i>Potentilla norvegica</i>	rough cinquefoil	1
<i>Erigeron canadensis</i>	fleabane	1

*grass species includes foxtail barley and a brome species

4.5 Discussion

4.5.1 Restoration Effects on Plant Parameters

Crop emergence levels, weed emergence levels and yield were monitored in the upper slope area of both experiments to examine the impacts of landscape restoration on plant species and abundance (including weeds). The results from the first field experiment of this study indicate that landscape restoration does increase weed emergence in the first year following the addition of soil. The favorable soil conditions including increased moisture and temperature are likely a contributing factor to increased weed emergence. The disturbed treatment showed similar levels of weed emergence indicating the soil addition was not the only contributing factor and that tillage also plays a role. In Block B in the first year of the study (2003), where significant increases in weed emergence were observed, significant increases in crop yield were also observed (pea yield only). This implies that although landscape restoration may increase weed pressures, it also creates more favorable conditions for crop growth perhaps enabling the crop to out-compete the weeds. For example, soil nitrate, phosphate and sulphate content were higher in the addition treatments compared with the control, which should result in higher crop yield. On the other hand, since the added nutrients from soil addition are not necessarily seed placed, greater available nutrients may favor weed recruitment over crop growth. This may help explain why a significant increase in flax yield was not observed. The lack of response from the flax crop in Block A may also be attributed to the dry weather in 2003 and to the lack of competitive ability of flax. In the second year of the experiment, crop yield was lower in the addition treatment compared with the control. Again, because of the increase in fertility, soil moisture and temperature, a greater

response from the crop in terms of yield was expected. The 2004 growing season was not favorable for canola; the low ambient temperatures and higher moisture throughout the growing season resulted in late harvest and the problem with the seeder caused patchy emergence. These are contributing factors to the poor yield response despite improved nutrient and soil physical conditions. It is interesting to note that weed emergence was higher in all treatments in 2003 compared with 2004. This may be the result of poor weed control in year one due to limited in-crop herbicide options for peas and flax. It also indicates that increased weed pressures resulting from the restoration practice may be short-lived.

In terms of weed species present, the addition of soil did not greatly increase the number of species in the upper slope area. Weed species numbers were higher in the addition plots in 2004; the number and type of plant species was consistent over both years. The species present were also consistent with those present throughout the field in which the experiment took place. The farm manager of the MZTRA farm has identified volunteer wheat, stinkweed and sheperd's purse, night-flowering catchfly, dandelion and Canada thistle (in ascending order of abundance) as the most common weeds in the surrounding cropland. The species that were found in the restored landscape should not pose a major problem to the producer as there are a variety of herbicides that can be used to control their growth, and because the species were already present within the cropland.

It is difficult to comment on weed emergence levels and species data obtained from the second field experiment due to irregular crop emergence. Bare areas within the plots will favor weed emergence and survival. For this reason, the effect of landscape restoration on weed emergence cannot be distinguished. The weed species identified in

FE1 were also identified in FE2 indicating similar weed populations may appear regardless of where the soil was taken from.

When assessing the potential for increasing weed pressures following landscape restoration, it may be useful to look at near-by and historical weed populations of the farm. Seedling populations from the previous year have been reported to be the best predictor of future weed seedling populations (du Croix Sissions et al. 2000). Past weeds would have contributed to the seed bank for a given area of land which in turn will influence what weeds arise following this land management practice.

In the lower slope area, plant emergence in the removal treatments increased from the first to the second year. This is likely the result of seed output from the first year plants contributing to the seed bank and, therefore, to the second year population. The species observed were weeds; no wetland species were observed. The weed species observed in the removal treatments included those identified as a problem in the adjacent cropland. The weed species found in the lower slope removal areas and the upper slope addition areas were mostly the same.

Removing soil from the edge of a wetland did not cause wetland species to return in the medium-term, nor did the species that exist in the controls return to the removal areas in the medium-term. This is somewhat expected based on the population model/life cycle strategy described by the r-K selection theory and Grime's strategies of plants (Begon et al. 1996; Holliday and Burton 2001). These models describe the relationship between a species and its habitat where in situations where there is high disturbance or there is stress with low competition, ruderal species will thrive and persist (a ruderal species is defined as a plant that moves rapidly into recently disturbed environments that

are rich in resources, Holliday and Burton 2001). The weedy plants that were observed are examples of ruderal species. Wetland species or the grasses currently present in the riparian area of the lower slope area won't likely return until the effects of the disturbance are long past. The wetland species will only return if the soil remains moist enough to favor germination.

4.5.2 Implications of the Soil Seed Bank on Landscape Restoration

The weed seed bank experiment indicates that most viable weed seeds exist near the soil surface (top 5 cm) based on weed emergence levels; very few to no weeds emerged at depth. The species found in this experiment were predominantly typical weed species and were also the species identified in FE1 and FE2 experiments. These results imply that weed species may establish in the cropland if this soil is used for landscape restoration. As a result, higher rates of herbicides used may be required or a more effective herbicide for the given weed species present may be required in the first year following restoration to help control the increased weed populations. Since the viable seed bank occurs in the top 5 cm of soil, the producer should remove soil from deeper (up to 20 cm) and smaller areas. This will dilute the surface seed bank and reduce the potential threat of weed establishment in the cropland.

Since the viable seed bank occurs largely in the surface soil, removal of greater than 5 cm of soil would remove the majority of the undesirable species and make the revegetation of the riparian area by these undesirable species more difficult. The species identified in the seed bank study were not wetland species; however, with the removal of a relatively large quantity of soil, wetland hydrology may be impacted and may result in wetter conditions in the riparian area, potentially extending the area of wetland plant

species.

Finally, since much of the seed bank is removed with the restoration practice, which species will re-vegetate the area of soil removal remains a question. By removing the undesirable weed seeds, the re-vegetation of the riparian zone by weed species may be more difficult. However, since wetland species were not found below the 20 cm depth of soil removal, re-vegetation by wetland species may be equally as difficult.

4.6 Summary and Conclusions

Landscape restoration does increase weed emergence in the upper slope area in the first year following restoration. Depending on the crop species, the increase in weed emergence may or may not affect crop yield. Crop yield improved over the control where soil was added; however, this only occurred with the pea crop; flax and canola did not show a response. A greater response from the crop was expected due to increased soil nutrients and improved soil moisture and temperature conditions.

The weed species numbers were not greatly affected by landscape restoration; however, this practice tends to cause a minor increase in the number of species present. The weed species identified in the seed bank are likely to be present in the restored area following this practice. Higher rates of herbicides used or a more effective herbicide for the weed species present may be required following landscape restoration to control weed populations. Because the weed species found at the experimental site were similar to those found in the surrounding cropland, the increased weed pressures should be controlled with appropriate herbicide selection and rotation. Planting a competitive crop variety following the restoration practice will also help minimize the impact of weeds.

Since the viable seed bank exists in the surface soil, removing greater depths of soil from smaller areas will dilute the seed bank of the soil being used in the restoration practice and, therefore, reduce the number of weeds emerging in the restored upper slope area.

5.0 GENERAL DISCUSSION

Landscape restoration is an innovative soil management practice that has the potential to reduce the impacts of soil erosion. To date, there is no research on landscape restoration as a soil management practice. Some work has shown there is a positive relationship between topsoil thickness and crop productivity; this has been proven by removing topsoil from productive soils with thick surface layers or by adding topsoil in depth increments to previously eroded soils. The aim of past research was to characterize the relationship between topsoil depth or soil erosion and crop productivity. The feasibility of landscape restoration as a soil management practice has never been studied even though it is being carried out in various parts of the world including North America, Europe and Asia. There are important agronomic, economic and environmental factors that should be considered when looking at landscape restoration as a soil management practice. The aim of this study was to look at the feasibility of landscape restoration in terms of its impacts on an important environmental concern (greenhouse gas emissions) and an important agronomic concern (the potential for spreading weeds within the landscape and increasing weed pressures).

Greenhouse gas emissions either remained the same or were reduced following landscape restoration. The act of removing soil reduced CO₂ emissions in the first year following removal. This was a consistent finding in both the field and growth chamber experiments. This is an important finding since various beneficial management practices are being used in agriculture to reduce atmospheric CO₂ levels. The reduction in CO₂ emissions where soil is removed can help offset any increases in CO₂ emissions in the

upper slope area. In the growth chamber, soil N₂O emissions were also reduced with the removal of soil.

Soil addition did not affect greenhouse gas emissions from soil, and in fact, the control treatments showed greater soil respiration rates than did the addition treatments. Because the experiments took place on zero-till soils, greater soil respiration was expected to result from the act of disking the soil and incorporating surface residues. Furthermore, because higher soil nitrate levels were observed in the addition treatments compared to the control, it is expected that higher nitrous oxide emissions should have also been observed. This study did not show either of these effects in the field; however, N₂O emissions were observed from addition treatments compared with the eroded control in the growth chamber. The lack of findings in the field experiment were either the result of a small sampling size and high spatial variability or because, in the case of CO₂, the major flux event was missed. In either case, the effect of CO₂ may simply be insignificant experimentally and in terms of long-term (many years) effects of landscape restoration on CO₂ emissions. The overall impact of the removal and addition of soil was positive or at the very least neutral, with respect to greenhouse gas emissions.

The issue of landscape restoration effects on plant or weed species and abundance was important for two reasons. If weed problems were created or magnified following this practice, crop yield may be compromised and greater amounts of herbicide use may be required. Loss of crop yield and increased herbicide use has economic and environmental costs. It is also important that the practice doesn't encourage new or uncommon and difficult to control weeds to emerge and persist. Landscape restoration did impact weed emergence in the first year following the practice where higher weed

emergence was observed in addition versus control treatments. However, the disturbed treatment showed similar or higher emergence numbers than the addition treatment indicating the mixing of the original zero-till surface material may have influenced weed seed germination and that the addition of soil may not have been the only cause. The species observed in both areas of interest were common to the site. Furthermore, despite higher weed emergence, the pea crop in the addition treatment out-yielded the control. This is likely a result of higher nutrient content of the soil following restoration. In the second year, weed emergence was similar across all treatments. In general, the impacts of landscape restoration on weeds were negligible.

One of the drawbacks of landscape restoration of a zero-till landscape that will affect greenhouse gas emissions, weeds and general soil quality is that the applied soil must be incorporated to smooth the soil surface and prepare the seedbed and requires zero-till fields to be tilled. This may drive off CO₂ by incorporating surface materials and roots; it may increase weed pressures thereby resulting in increased herbicide use. Some tillage or harrowing of the soil following soil addition is required to improve the seedbed, but the amount of tillage may be minimized by using the appropriate equipment (i.e. an earth mover or scraper), carrying out the practice when conditions are optimal (in the fall when the soil is dry) and implementing zero tillage soil management following the practice. The impacts on weed populations may be minimized by using cultural practices such as planting winter wheat or competitive crop varieties.

In general, the issue of landscape restoration effects on greenhouse gas emissions may not be an important consideration for most producers when deciding for or against the adoption of this practice. The findings from this study are important, however,

because carbon storage in agricultural landscapes is currently being promoted to help achieve national greenhouse gas reductions in compliance with the Kyoto Protocol. Producers may become more interested in landscape restoration impacts on greenhouse gas emissions if, for example, carbon credit trading becomes reality or economic incentives are provided for increasing carbon storage/reducing greenhouse gas emissions in agricultural landscapes. Although the findings from this study are somewhat preliminary and general, they do provide information on the medium-term impacts of landscape restoration on greenhouse gas emissions and a starting point from where future research on this topic can evolve.

The factors that are likely to weigh more heavily with producers when considering adoption of this practice are landscape restoration effects on soil nutrients, crop and weed emergence and weed species because these are the factors that translate directly into economic benefits or costs. This study provided some information on all of these factors but more research is needed on landscape restoration effects on crop yield. Crop yield information will help determine the economic benefits and costs of the practice. Finally, another important factor for producers to consider but was not studied here is equipment (earthmover or scraper) availability and associated costs. Producers may be reluctant to attempt this practice if they do not already own an earthmover and have to rent the equipment because this will add to the cost of the practice.

6.0 SUMMARY AND CONCLUSIONS

The primary objectives of this study were two-fold: to measure greenhouse gas emissions from the landscape and to monitor plant emergence (including crops and weeds) and species present following landscape restoration. The two key areas of interest included in the study were the upper slope area to which soil was applied and the lower slope area from where soil was removed. The objectives were investigated with two field and two growth chamber experiments.

In the field, the effect of adding soil to eroded hilltops on greenhouse gas emissions was minimal. In the first year following the addition of soil, soil temperature and moisture increased slightly over the control; however, this effect was less pronounced in year two and showed little impact on greenhouse gases. Soil microbial biomass carbon and soil nitrate content were significantly higher where soil was added and therefore a greater impact of soil addition on greenhouse gas flux was expected. Where soil was removed, soil respiration was significantly reduced compared with non-removal treatments. Nitrous oxide and methane emissions were unaffected. This corresponds well with the lower soil nitrate and soil microbial biomass carbon levels found in the removal treatments.

In the growth chamber, the addition of soil did affect gas flux; mean carbon dioxide and nitrous oxide emissions were higher where soil was added compared with the eroded soil. This does not correspond with field results which may be attributed to the higher soil moisture content maintained in the growth chamber compared with the field experiment soils. The removal of soil resulted in significantly lower carbon dioxide and

nitrous oxide emissions compared with the control. The lower carbon dioxide emissions corresponds with field experiment findings and supports the conclusion that the overall impact of landscape restoration on greenhouse gas emissions is positive in the medium-term.

Plant measurements showed that landscape restoration will increase the emergence and yield of some crops in the upper slope area, and that it will increase weed emergence in the first following year. Smaller seeded crops such as flax and canola did not respond well to the restoration practice. This was attributed to such factors as poor competitive ability of flax, and poor seedling emergence and establishment of canola. Weed species present also increased slightly; however, the species were similar to those found at the site. As a result of the increased weed pressures, greater herbicide use may be required in the first year following the restoration and competitive crops should be used. Where soil was removed, the species present were predominantly weedy and were not wetland species or the native species observed in the control plots.

The weed seed bank study was useful in showing that the viable weed seeds exist in the surface soil within the soil profile and that the species present are similar to those found within the adjacent cropland. As a result, when using accumulated soil to restore hilltops, soil taken from deeper and smaller areas will likely dilute the surface seed bank and reduce the increase in weed emergence in the upper slope area.

Overall, landscape restoration showed a smaller impact on the soil environment (namely soil moisture and temperature, soil organic carbon) than was expected. Soil moisture was slightly higher in the addition treatment compared with the control; however, these results were variable and inconsistent. Soil moisture levels were higher in

the removal treatments compared with the controls likely due to the difference in vegetation present. The increased moisture in the removal treatment did not increase nitrous oxide emissions as would be expected; this can be explained by the low soil nitrate and soil microbial biomass carbon levels in the removal treatments compared with the controls. The organic carbon content of soil in the upper slope area did not increase with the addition of soil. Based on lower slope area organic carbon content of the surface soil, an increase of 4% should have been observed in the addition treatments. The reason this increase was not observed is unclear. Soil fertility was positively impacted.

The results from this study suggest that landscape restoration is a practice that can be used as a soil management technique for improving soil quality on eroded hilltops with minimal environmental impacts. The positive influence of restoration on soil fertility should result in increased yields of large-seed grain crops that have a good competitive ability. Increased herbicide use may be required in the first year following restoration.

7.0 CONTRIBUTIONS TO KNOWLEDGE

As the first of its kind, this study provides preliminary information regarding environmental and agronomic impacts of landscape restoration. Specifically, medium-term impacts of landscape restoration on the soil environment including soil temperature, moisture and nutrient status, greenhouse gas emissions, crop emergence, crop yield, weed emergence and weed species were determined. In general, this study showed that landscape restoration may be used a soil management practice to restore eroded upper slopes with minimal negative environmental and agronomic impacts if the appropriate conditions exist and if the appropriate equipment is used; that is if the soil is relatively dry, such as in the fall, and an earth mover or scraper is used.

Because this study was the first of its kind and relatively large in scope, it is not without weaknesses. The broad questions and the disconnected nature of the parameters measured resulted in a large volume of information being collected, but little correlation between greenhouse gases and weeds, for example, could be made. Because research looking at landscape restoration effects on the soil environment, greenhouse gases and weeds does not exist, it was impossible to compare the results found in this study with the literature. There are still questions that need to be addressed. Future research that focuses in greater detail on each of the parameters investigated in this study is required. Also, a more thorough investigation into the impacts of landscape restoration on crop yield is necessary from an agronomic perspective, but also to calculate the economic cost-benefit of the practice. The issue of soil salinity was touched upon in this study, however, a

more thorough analysis of using soils rich in salts to restore eroded hilltops should be carried out.

If a similar study is carried out in the future, the appropriate equipment should be used to establish removal and addition treatments, such as a scraper, and the plot size should be larger to facilitate the use of field scale equipment. Ideally, the treatments should be monitored for a longer period of time to evaluate the long-term effects of landscape restoration on the parameter in question.

8.0 REFERENCES

- Bajracharya, R. M., Lal, R. and Kimble, J.M. 2000.** Diurnal and season CO₂-C flux from soil as related to erosion phases in Central Ohio. *Soil Sci. Soc. Am. J.* **64**: 286-293.
- Barberi, P. and Lo Cascio, B. 2001.** Long-term tillage and crop rotation effects on weed seedbank size and composition. *European Weed Research Society Weed Research.* 325-340.
- Battiston, I. A., Miller, M. H. and Shelton, I. J. 1987.** Soil erosion and corn yield in Ontario, I. Field evaluation. *Can. J. Soil Sci.* **67**: 731-745.
- Beckie, H.J., Moulin, A.P. and Pennock, D.J. 1997.** Strategies for variable rate nitrogen fertilization in hummocky terrain. *Can. J. Soil Sci.* **77**: 589-595.
- Begon, M., Harper, J.L. and Townsend, C.R. 1996.** Ch 14: Life-History Variation. Pages *in Ecology - Individuals, Populations and Communities.* Blackwell Sciences Ltd.
- Benoit, D. L., Derksen, D.A. and Panneton, B. 1992.** Innovative approaches to seedbank studies. *Weed Science.* **40**: 660-669.
- Benvenuti, S., Macchia, M. and Miele, S. 2001.** Quantative analysis of emergence of seedlings from buried weed seeds with increasing soil depth. *Weed Science.* **49**: 528-535.
- Biederbeck, V. O., Campbell, C.A., Rasiyah, V., Zentner, R.P. and Wen, G. 1998.** Soil quality attributes as influenced by annual legumes used as green manure. *Soil Biol. Biochem.* **30**: 1177-1185.
- Bower, C.A. and Wilcox, L.V. 1965.** Ch 62: Soluble salts *in* Black, C.A., Evans, D.D., White, J.L., Ensminger, C.E. and Clark, F.E. *Methods of Soil Analysis Part 2: Chemical and Microbiological Properties.*
- Boyd, N. and Van Acker, R.C. 2003.** The effects of depth and fluctuating soil moisture on the emergence of eight annual and six perennial plant species. *Weed Science.* **51**: 725-730.
- Boyd, N. and Van Acker, R.C. 2004.** Seed and microsite limitations to emergence of four annual weed species. *Weed Science.* **52**: 571-577.
- Brady, N. C. and Weil, R. R. 2002.** *The Nature and Properties of Soil.* Prentice Hall. Upper Saddle River, NJ.

- Bremner, J. M. 1997.** Source of nitrous oxide in soils. *Nutrient Cycling in Agroecosystems*. **49**: 7-16.
- Buhler, D. D., Hartzler, R.G. and Forcella, F. 1997.** Implications of weed seedbank dynamics to weed management. *Weed Science*. **45**: 329-336.
- Bullied, W. J., Marginet, A.M. and Van Acker, R.C. 2003.** Conventional- and conservation-tillage systems influence emergence periodicity of annual weed species in canola. *Weed Science*. **51**: 886-897.
- Burnett, E., Stewart, B. A. and Black, A. L. 1985.** Regional effects of soil erosion on crop productivity - Great Plains. Pages 285-304 *in* Follet, R. F. and Stewart, B. A., eds. *Soil Erosion and Crop Productivity*. ASA-CSSA-SSSA.
- Burton, D. L., and Beauchamp, E.G. 1994.** Profile nitrous oxide and carbon dioxide concentrations in a soil subject to freezing. *Soil Sci. Soc. Am. J.* **58**: 115-122.
- Campbell, C. A., McConkey, B.G., Zentner, R.P., Selles, F., and Curtin, D. 1996.** Long-term effects of tillage and crop rotation on soil organic C and total N in a clay soil in southwestern Saskatchewan. *Can. J. Soil Sci.* **76**: 395-401.
- Campbell, C. A., Selles, F., Lafond, G.P. and Zentner, R.P. 2001.** Adopting zero tillage management: Impact on soil C and N under long-term crop rotations in a thin Black Chernozem. *Can. J. Soil Sci.* **81**: 139-148.
- Cardina, J. and Sparrow, D.H. 1996.** A comparison of methods to predict weed seedling populations from the soil seedbank. *Weed Science*. **44**: 46-51.
- Carter, D. L., Berg, R. D. and Sander, B. J. 1985.** The effect of furrow irrigation erosion on crop productivity. *Soil Sci. Soc. Am. J.* **49**: 207-211.
- Cavers, P. B. 1995.** Seed banks: Memory in soil. *Can. J. Soil Sci.* **75**: 11-13.
- Cole, C. V., Duxbury, J., Freney, J., Heinemeyer, O., Minami, K., Mosier, Paustian, K., Rosenber, N., Sampson, N., Sauerberck, D. and Zhao, Q. 1997.** Global estimates of potential mitigation of greenhouse gas emissions by agriculture. *Nutrient Cycling in Agroecosystems*. **49**: 221-228.
- Corre, M. D., van Kessel, C. and Pennock, D.J. 1996.** Landscape and season patterns of nitrous oxide emissions in a semiarid region. *Soil Sci. Soc. Am. J.* **60**: 1806-1815.
- Dieleman, J. A., Mortensen, D.A., Buhler, D.D. and Ferguson, R.B. 2000.** Identifying associations among site properties and weed species abundance. II. Hypothesis generation. *Weed Science*. **48**:576-587.

- Dormarr, J. F., Lindwall, C. W. and Kozub, G. C. 1988.** Effectiveness of manure and commercial fertilizer in restoring productivity of an artificially eroded Dark Brown Chernozemic soil under dryland conditions. *Can. J. Soil Sci.* **68**: 669-679.
- Dormarr, J. F., Lindwall, C. W. and Kozub, G. C. 1997.** Role of continuous wheat and ammendments in ameliorating an artificially eroded Dark Brown Chernozemic soil under dryland conditions. *Can. J. Soil Sci.* **77**: 271-279.
- Drury, C.F., Oloya, T.O., McKenney, D.J., Grerorich, E.G., Tan, C.S., and vanLulyk, C.L. 1998.** Long-term effects of fertilization and rotation on denitrification and soil carbon. *Soil Sci. Soc. Am. J.* **62**:1572-1579.
- du Croix Sissons, M. J., Van Acker, R.C., Derksen, D.A. and Thomas, A.G. 2000.** Depth of seedling recruitment of five weed species measured in situ in conventional- and zero-tillage fields. *Weed Science.* **48**: 327-332.
- Eghball, B., Mielke, L.N., McCallister, D.L. and Doran, J.W. 1994.** Distribution of organic carbon and inorganic nitrogen in a soil under various tillage and crop sequences. *J. of Soil and Water Cons.* **49**: 201-205.
- El-Swaify, S. A. 2001.** Impact of erosion and restoration on water and nutrient use efficiency in a Hawaii Oxisol. *Commun. Soil Sci. Plant Anal.* **32**: 1187-1201.
- Felix, J. and Owen, M. D. K. 2001.** Weed seedbank dynamics in post conservation reserve program land. *Weed Science.* **49**: 780-787.
- Frye, W. W., Bennett, O. L. and Buntley, G. J. 1985.** Restoration of crop productivity on eroded or degraded soils. Pages 335-356 *in* Follet, R. F. and Stewart, B. A., eds. *Soil Erosion and Crop Productivity.* ASA-CSSA-SSSA.
- Goodroad, L. L., and Keeney, D.R. 1985.** Site of nitrous oxide production in field soils. *Biol Fertil Soils.* **1**: 3-7.
- Govers, G., Lobb, D. A. and Quine, T. A. 1999.** Tillage erosion and tillage translocation. *Soil and Tillage Res.* **51**: 167-174.
- Hartwig, N. L., and Ulrich Ammon, H. 2002.** 50th Anniversary-Invited Article: Cover crops and living mulches. *Weed Science.* **50**: 688-699.
- Havlin, J. L., Beaton, J.D., Tisdale, S.L. and Nelson, W.L. 1999.** *Soil Fertility and Fertilizers.* Prentice-Hall, Inc. Upper Saddle River, New Jersey. 499 pp.
- Holliday, N. J. and Burton, D.L. 2001.** 65.351 Agroecology Class Notes. University of Manitoba, Faculty of Agricultural and Food Sciences. Winnipeg MB.

- Horner, G. M., McCall, A. G. and Bell, F. G. 1944.** Investigation in erosion control and the reclamation of eroded land at the Palouse Conservation Experiment Station, Pullman Wash. 1931-1942. United States Department of Agriculture. Washington. 83 pp.
- Hutchinson, G. L., and Livingston, G.P. 2002.** Gas Flux. Pages pp 1159-1182 *in* Dane, J. H., and Topp, G.C., ed. Methods of Soil Analysis, Part 4, Physical Methods. Soil Sci. Soc. Am.
- Izaurrealde, R. C., Nyborg, M., Solber, E.D., Janzen, H.H., Arshad, M.A., Malhi, S.S. and Molina-Ayala, M. 1997.** Carbon storage in eroded soils after five years of reclamation techniques. Pages *in* Lal, R., Kimble, J.M., Follet, R.F. and Stewart, B.A., ed. Soil Processes and the Carbon Cycle. CRC Press.
- Janzen, H. H., Desjardins, R. L., Asselin, J.M.R. and Grace, B. 1998.** The Health of Our Air: Toward sustainable agriculture in Canada. Agriculture and Agri-Food Canada. Ottawa, ON.
- Jenkinson, D. S., Brookes, P.C., and Powlson, D.S. 2004.** Measuring soil microbial biomass. *Soil Biol. Biochem.* **36**: 5-7.
- King, D. J., Cossette, J.M., Eilers, R.G., Grant, B.A., Lobb, D.A., Padbury, G.A., Rees, H.W., Shelton, I.J., Tajek, J., Wall, G.J. and van Vliet, L.P.J. 2000.** Risk of Tillage Erosion. Government of Canada. Ottawa. pp. 8, 77-83.
- Lal, R., Ahmadi, M. and Bajracharya, R. M. 2000.** Erosional impacts on soil properties and corn yield on alfisols in Central Ohio. *Land Degrad. Develop.* **11**: 575-585.
- Lal, R., and Kimble, J.M. 1997.** Conservation tillage for carbon sequestration. *Nutrient Cycling in Agroecosystems.* **49**: 243-253.
- Langdale, G. W. and Shrader, W. D. 1982.** Soil erosion effects on soil productivity of cultivated cropland. Pages 41-51 *in* Kral, D. M. and Hawkins, S., eds. Determinants of Soil Loss Tolerance. ASA.
- Larney, F. J., Bullock, M. S., Janzen, H. H., Ellert, B. H. and Oson, E. C. S. 1998.** Wind erosion effects on nutrient redistribution and soil productivity. *J. Soil and Water Cons.* **53**: 133-140.
- Larney, F. J., Janzen, H. H., Oslon, B. M. and Lindwall, C. W. 2000a.** Soil quality and productivity responses to simulated erosion and restorative amendments. *Can. J. Soil Sci.* **80**: 515-522.

- Larney, F. J., Olson, B. M., Janzen, H. H. and Lindwall, C. W. 2000b.** Early impact of topsoil removal and soil amendments on crop productivity. *Agron. J.* **92**: 948-956.
- Lemke, R. L., Izaurralde, R.C., Nyborg, M. and Solberg, E.D. 1998.** Tillage and N source influence soil-emitted nitrous oxide in the Alberta Parkland region. *Can. J. Soil Sci.* **79**:15-24.
- Li, Y. and Lindstrom, M. J. 2001.** Evaluating soil quality - soil redistribution relationship on terraces and steep hillslope. *Soil Sci. Soc. Am. J.* **65**: 1500-1508.
- Lind, A. M., and Eiland, F. 1989.** Microbial characterization and nitrate reduction in subsurface soils. *Biol Fertil Soils.* **8**: 197-203.
- Lobb, D. A., Kachanoski, R. G. and Miller, M. H. 1995.** Tillage translocation and tillage erosion on shoulder slope landscape positions measured using ¹³⁷Cs as a tracer. *Can. J. Soil Sci.* **75**: 211-218.
- Lobb, D. A., and Kachanoski, R.G. 1999.** Modelling tillage translocation on the topographically complex landscapes of southwestern Ontario. *Soil Tillage Research.* **51**: 261-277.
- Lobb, D. A. 1999.** Understanding and managing the causes of soil-landscape variability. *Success in Precision Agriculture, 4th Conference, Brandon Manitoba.*
- Lobb, D. A., Burton, D.L, Lindstrom, M.J., and Reicosky, D.C. 2002.** Impacts of soil erosion on the production and emission of greenhouse gases and carbon sequestration in the Canadian Prairies. *OECD Workshop on Soil Organic Carbon Indicators for Agricultural Land. Ottawa, Canada.*
- Luo, J., Tillman, R.W., White, R.E. and Ball, P.R. 1998.** Variation in denitrification activity with soil depth under pasture. *Soil Biol. Biochem.* **30**: 897-903.
- Lyles, L. 1977.** Wind Erosion: Processes and Effect on Soil Productivity. *Transactions of the ASAE.* 880-884.
- MacLeod, C., Burton, D.L., Lobb, D.A. and Erb, M. 2005 (Submitted).** Influence of method of manure application and tillage system on soil respiration and nitrous oxide production in a heavy clay soil. *Can. J. Soil Sci.*
- Malhi, S. S., McGill, W.B. and Nyborg, M. 1990.** Nitrate losses in soils: Effect of temperature, moisture and substrate concentration. *Soil Biol. Biochem.* **22**: 733-737.

- Malhi, S. S., Grant, C.A., Johnston, A.M., and Gill, K.S. 2001.** Nitrogen fertilization management for no-till cereal production in the Canadian Great Plains: a review. *Soil and Tillage Res.* **60**: 101-122.
- Masse, T. W. 1990.** Simulated erosion and fertilizer effects on winter wheat cropping Intermountain Dryland Area. *Soil Sci. Soc. Am. J.* **54**: 1720-1725.
- Masse, T. W. and Waggoner, H. O. 1985.** Productivity losses form soil erosion on dry cropland in the intermountain area. *J. of Soil and Water Cons.* **September-October**: 447-450.
- McCarty, G. W., Lyssenko, N.N. and Starr, J.L. 1998.** Short-term changes in soil carbon and nitrogen pools during tillage management transition. *Soil Sci. Soc. Am. J.* **62**:1564-1571.
- Meixner, F. X., and Eugster, W. 1999.** Effects of landscape pattern and topography on emission and transport. Pages *in* Tenhunen, J. D. and Kabut, P., eds. *Integrating Hydrology, Ecosystem Dynamics and Biogeochemistry in Complex Landscapes.* John Wiley & Sons Ltd.
- Meyer, L. D., Bauer, A. and Heil, R. D. 1985.** Experimental approaches for quantifying the effect of soil erosion on productivity. Pages 213-234 *in* Follet, R. F. and Stewart, B. A., eds. *Soil Erosion and Crop Productivity.* ASA-CSSS-SSSA.
- Mielke, L. N. and Schepers, J. S. 1986.** Plant response to topsoil thickness on an eroded loess soil. *J. of Soil and Water Cons.* **January-February**: 59-63.
- Mosier, A. R. 1998.** Soil processes and global change. *Biol Fertil Soils.* **27**: 221-229.
- Paul, E. A., and Clark, F.E. (Eds.). 1996.** *Soil microbiology and biochemistry.* Academic Press. San Diego.
- Pekrun, C., Hans-Peter, K. and Claupein, W. 2003.** Soil tillage for sustainable nutrient management. Pages *in* El Titi, A., ed. *Soil Tillage in Agroecosystems.* CRC Press.
- Pennock, D. J., van Kessel, C., Farrel, R. E. and Sutherland, R. A. 1992.** Landscape-scale variations in dentirification. *Soil Sci. Soc. Am. J.* **56**: 770-776.
- Podolsky, G. P., and Schindler, D. 1994.** *Soils of the Manitoba Zero Tillage Research Association Research Farm.* Manitoba Land Resource Unit, Agriculture and Agri-Food Canada.
- Quansah, C. 1981.** The effect of soil type, slope, rain intensity and their interaction on splash detachment and transport. *J. of Soil Sci.* **32**: 215-224.

- Reicosky, D. C. 1997a.** Tillage-induced CO₂ emission from soil. *Nutrient Cycling in Agroecosystems*. **49**: 273-285.
- Reicosky, D. C., Dugas, W.A., and Torbert, H.A. 1997b.** Tillage-induced soil carbon dioxide loss from different cropping systems. *Soil and Tillage Res.* **41**:105-118.
- Robbins, C. W., Mackey, B. E. and Freeborn, L. L. 1997.** Improving exposed subsoils with fertilizers and crop rotations. *Soil Sci. Soc. Am. J.* **61**: 1221-1225.
- Sadler, J. M. 1984.** Effects of topsoil loss and intensive cropping on soil properties related to the crop production potential of a podzolic grey luvisol. *Can. J. Soil Sci.* **64**: 533-543.
- Salinas-Garcia, J. R., Hons, F.M. and Matocha, J.E. 1997.** Long-term effects of tillage and fertilization on soil organic matter dynamics. *Soil Sci. Soc. Am. J.* **61**: 152-159.
- SAS. 2000.** SAS User's Guide: Statistics, SAS Institute Inc., Cary, NC., U.S.A
- Schumacher, T. E., Lindstrom, M. J., Schumacher, J. A. and Lemme, G. D. 1999.** Modeling spatial variation in productivity due to tillage and water erosion. *Soil and Tillage Res.* **51**: 331-339.
- Smith, E. G., Peng, Y., Lerohl, M. and Larney, F. J. 2000.** Economics of N and P fertilization to restore wheat yields on three artificially eroded sites in southern Alberta. *Can. J. Soil Sci.* **80**: 165-169.
- Smith, K. A., Ball, T., Conen, F., Dobbie, K.E., Massheder, J. and A. Rey. 2003.** Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes. *European Journal of Soil Science.* **54**: 779-791.
- Stevenson, F. C., Legere, A., Simard, R.R., Angers, D. A., Pageau, D. and Lafond, J. 1998.** Manure, tillage and crop rotation: effects on residual weed interference in spring barley cropping systems. *Agron. J.* **90**: 496-504.
- Stevenson, F. C., and van Kessel, C. 1996.** A landscape-scale assessment of the nitrogen and non-nitrogen rotation benefits of pea. *Soil Sci. Soc. Am. J.* **60**: 1979-1805.
- Tanaka, D. L. and Aase, J. K. 1989.** Influence of topsoil removal and fertilizer application on spring wheat yields. *Soil Sci. Soc. Am. J.* **53**: 228-232.
- Troeh, F., Hobbs, J. A. and Donahue, R. L. 1980.** Water erosion and sedimentation. *in* Troeh, M. R., ed. *Soil and Water Conservation*.

- Van Cleemput, O. 1998.** Subsoils: chemo- and biological denitrification, N₂O and N₂ emissions. *Nutrient Cycling in Agroecosystems*. **52**: 187-194.
- van den Pol-van Dasselaar, A., Corre, W.J., Prieme, A., Klemedtsson, A. K., Weslien, P., Stein, A., Klemedtsson, L. and Oenema, O. 1998.** Spatial variability of methane, nitrous oxide and carbon dioxide emissions from drained grasslands. *Soil Sci. Soc. Am. J.* **62**: 810-817.
- Van Muysen, W. and Govers, G. 2002.** Soil displacement and tillage erosion during secondary tillage operations: the case of rotary harrow and seeding equipment. *Soil and Tillage Res.* **65**: 185-191.
- Van Muysen, W., Govers, G. and Van Oost, K. 2002.** Identification of important factors in the process of tillage erosion: the case of mouldboard tillage. *Soil and Tillage Res.* **65**: 77-93.
- Vance, E. D., Brookes, P.C., and Jenkinson, D.S. 1987.** An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.* **19**: 703-707.
- Verity, G. E. and Anderson, D. W. 1990.** Soil erosion effects on soil quality and yield. *Can. J. Soil Sci.* **70**: 471-484.

9.0 APPENDICES

Appendix A

Soil Addition and Removal Depth Measurements, Field Plans and Topographical Survey Maps

Table A.1 Soil addition depth measurements from nine places within each plot following disking in 2003 in FE1

Plot #	Treatment ID	Measurement (cm)									Mean
		1	2	3	4	5	6	7	8	9	
Block A – Upper Slope Area											
1	U1A	7	9	12	9	10	7	9	8	10	9.0
3	U2A	8	10	9	8	9	9	8	10	7	8.7
5	U1A	11	10	9	9	9	10	9	10	11	9.8
7	U2A	8	9	10	9	9	9	8	10	11	9.2
9	U1A	7	11	9	10	14	9	7	12	11	10.0
11	U2A	8	7	9	13	10	9	8	11	9	9.3
Block B – Upper Slope Area											
1	U1A	9	10	8	10	11	9	13	9	9	9.8
3	U2A	8	8	6	11	8.5	10	10	9	8	8.7
5	U1A	8	7	12	10	11	10	7	8	9	9.1
7	U2A	9	10	9	11	11	12	10	11	11	10.4
9	U1A	9	9	10	8	9	9	7	9	11	9.0
11	U2A	8	10	9	9	12	10	6	11	10	9.4

Table A.2 Soil removal depth measurements from nine places within each plot following disking in 2003 in FE1

Plot #	Treatment ID	Measurement (cm)									Mean
		1	2	3	4	5	6	7	8	9	
Block A – Lower Slope Area											
1	L1R	17	18	17	25	26	28	33	32	34	26
3	L2R	18	18	15	24	23	26	15	15	14	19
5	L1R	19	20	22	26	25	24	23	22	23	23
7	L2R	18	14	14	16	18	18	22	21	18	18
9	L1R	23	23	21	29	28	26	21	17	18	23
11	L2R	17	16	19	18	18	19	18	14	17	17
Block B – Lower Slope Area											
1	L1R	18	12	14	24	26	26	20	21	19	20
3	L2R	24	17	21	24	23	24	19	20	18	21
5	L1R	19	18	13	22	21	23	20	19	20	19
7	L2R	23	24	24	22	28	28	20	19	13	22
9	L1R	21	20	21	17	16	17	21	20	21	19
11	L2R	24	20	21	23	23	22	17	14	17	20

Table A.3 Soil addition depth measurements from nine places within each plot following disking in 2004 in FE2

Plot #	Measurement (cm)									Mean
	1	2	3	4	5	6	7	8	9	
1	10	12	13	11	6	13	13	11	9	10.9
5	17	13	11	11	11	16	6	10	16	12.3
7	13	12	13	15	15	20	7	18	10	13.7
12	11	14	13	14	9	18	14	10	9	12.4
14	8	18	11	12	18	18	14	18	13	14.4
18	9	13	17	18	12	15	16	16	14	14.4
19	13	10	10	14	14	12	13	12	15	12.6
22	12	15	10	15	13	10	14	15	10	12.7
26	10	17	13	16	14	14	13	15	9	13.4

Block A	Upper Slope		Lower Slope	
P12	Control	P12	Control	N →
P11	Addition	P11	Removal	
P10	Disked	P10	Mowed	
P9	Addition	P9	Removal	
P8	Control	P8	Control	
P7	Addition	P7	Removal	
P6	Disked	P6	Mowed	
P5	Addition	P5	Removal	
P4	Control	P4	Control	
P3	Addition	P3	Removal	
P2	Disked	P2	Mowed	
P1	Addition	P1	Removal	

Block B	Upper Slope		Lower Slope
P12	Control	P12	Removal
P11	Addition	P11	Control
P10	Disked	P10	Removal
P9	Addition	P9	Mowed
P8	Control	P8	Removal
P7	Addition	P7	Control
P6	Disked	P6	Removal
P5	Addition	P5	Mowed
P4	Control		
P3	Addition		
P2	Disked		
P1	Addition	P4	Control
		P3	Removal
		P2	Mowed
		P1	Removal

Treatment Descriptions:

U1A/U2A Addition addition of soil to a depth of 10 cm plus disking
 U1D Disked disking only
 U2C Control no addition or disking

L1R/L2R Removal removal of soil to a depth of 20 cm following the removal of
 standing plant biomass
 L1D Mowed removal of standing plant biomass
 L2C Control no removal or mowing

Figure A.1 Field layout for FE1.

**Upper Slope
Block* Treatment**

1	Addition
	Control
	Disked
2	Disked
	Control
	Addition
3	Addition
	Disked
	Control
4	Disked
	Control
	Addition
5	Control
	Addition
	Disked
6	Disked
	Control
	Addition
7	Addition
	Disked
	Control
8	Control
	Disked
	Addition
9	Control
	Addition
	Disked

Depressions (three)**

East	Removal
	Control

Centre	Removal
	Control

West	Removal
	Control

Treatment Descriptions:

Addition addition of soil to a depth of 10 cm plus disking

Disked disking only

Control no addition or disking

Removal removal of soil to a depth of 20 cm

Control no removal

* blocks are not all adjacent as in FE1 plots, these are arranged in nine different areas along the eroded ridge i.e. each block is completely independent

Figure A.2 Field layout for FE2.





Figure A.3 Topographic maps of Block A.





Figure A.4 Topographic maps of Block B and FO3.

Appendix B

Gas Chromatograph Specifications

Gas Chromatograph Specifications (taken from MacLeod et al. 2005)

Gas analysis was performed using a Varian Star 3800 Gas Chromatograph (Varian, Mississauga, ON) fitted with three detectors (electron capture - ECD, flame ionization - FID and thermal conductivity - TCD) and a Combi-PAL autosampler. The autosampler removes a 2.5 mL volume from the sample tube and injects this into a sample valve that delivers 0.25 mL to the ECD, TCD and FID in series. The ECD was operated at 300 °C, 10%Ar, 90%CH₄ carrier gas at flow rate of 30 mL min⁻¹ (13.0 psi), Porapak QS 80/100 precolumn (0.32 cm diameter x 46 cm length) and analytical columns (0.32 cm diameter x 183 cm length) in a column oven operated at 70 °C. A precolumn was used in combination with a four-port valve to remove water from the sample. The FID was operated at 250 °C, carrier gas was prepurified helium at 10 mL min⁻¹ passing through a Porapak N analytical column (0.32 cm x 46 cm) operated at 70 °C. The TCD was operated at 130 °C with a prepurified helium carrier gas at 30 mL min⁻¹ (20 psi), Haysep D 80/100 analytical column (0.32 cm diameter x 183 cm length) maintained at 70 °C. Five replicates of two concentrations of standard gas mixtures (same concentrations as those used during sampling) were included in each run and were used to construct standard curves. The standard gases collected during sampling were used to confirm sample integrity during sampling and storage.

Appendix C

Soil Microbial Biomass Extraction Methodology

Equipment per soil sample:

- 2 square French bottles
- 2 #5 stoppers
- 2 Whatman No. 5 filter papers
- 4 30mL scintillation vials
- CHCl_3
- must be purified, dried, and distilled in glass
- must not be stabilised with ethanol
- boiling chips
- 150mL beaker
- desiccator able to withstand a high vacuum without implosion
- vacuum pump
- fumehood
- 10% HCl acid bath
- 0.5 M K_2SO_4 solution

Procedure

Solutions

- Prepare 0.5 M K_2SO_4 solution by dissolving 87.135g K_2SO_4 crystals in 1L distilled water.
 - Use moderate heat to help expediate the process.

Soil

- Weigh out two 25g portions of soil into the square French bottles.
 - one sample will be fumigated for 24h and then extracted
 - one sample will be extracted immediately

Fumigation

1. Prepare the desiccator for fumigation by lining it with moistened paper towels.
2. Place samples in desiccator with a 150mL beaker containing approximately 50mL of CHCl_3 and boiling chips.
3. Seal and evacuate the desiccator, taking care to vent the fumes released by the vacuum pump into the fumehood, until the CHCl_3 boils vigorously, and continue evacuating for approximately 1 minute.
4. Seal the desiccator under vacuum by turning collar, and leave for 24 hours.
5. After 24 hours, release the vacuum by turning the desiccator collar; a hissing noise should be heard. Remove the beaker of CHCl_3 and the paper towels.
6. Remove the residual CHCl_3 vapour from the soil samples by repeatedly evacuating the chamber 3 times for about 30 seconds each time.

Extraction

- Unfumigated samples are extracted immediately after weighing, while fumigated samples are extracted after 24 hours of fumigation.
1. Add 50mL of 0.5M K₂SO₄ to the square French bottles using a repipettor. Stopper the bottles using #5 stoppers. For each set of extractions, prepare two solution blanks containing only K₂SO₄.
 2. Place the bottles on a lateral shaker set at high speed for 1 hour.
 3. After shaking, pass the soil suspension through Whatman No. 5 filter paper.
 - Funnels are not necessary; filter paper is folded and placed in the funnel rack directly.
 - Filter paper should be rinsed with approximately 50mL deionised water prior to filtration.
 4. Collect filtrate in 30mL scintillation vials.
 - Two sets of filtrate samples are collected – A set and B set.
 - Vials should be switched when the A vial is about 3/4 full.
 - Vials should be labelled with the site name, original date of sampling, sample code, “F” or “U” for fumigated or unfumigated, and “A” or “B.”
 5. Cap vials and placed in the freezer as soon as possible. The B set may be left overnight if the filtration time prolonged.

Analysis

- Analyse filtrate for N, C, NO₃⁻, and NH₄⁺ using the Technicon Auto-Analyser.

Calculations

1. Calculate the mass of C (μg g⁻¹ soil)

$$= \frac{[(\mu\text{g mL}^{-1} \text{ C})_{\text{fumigated}} - (\mu\text{g mL}^{-1} \text{ C})_{\text{blank}}] \cdot [\text{mL K}_2\text{SO}_4 + (\text{mass wet soil} \cdot \text{GMC})]}{(\text{mass of wet soil}/(1 + \text{GMC}))}$$

- μg mL⁻¹ C comes from the Auto-Analyser
- mass of wet soil/(1 + GMC) gives the mass of oven dry soil
- mass wet soil • GMC gives the mass of water in the sample
- * to calculate the mass of C in the unfumigated sample, substitute the appropriate data value for (μg mL⁻¹ C)_{unfumigated}

2. Calculate the mass of microbial biomass C (μg C g⁻¹ soil)

$$= \frac{\text{CO}_2\text{-C (fumigated)} - \text{CO}_2\text{-C (unfumigated)}}{(0.25) \cdot (\text{mass dry soil})}$$

- where 0.25 is a correction factor

- * To calculate microbial biomass N, substitute the Auto-Analyser N data into the above steps, and use 0.18 as the correction factor in Step 2.

Safety

- All technicians are responsible for familiarising themselves with the Materials Safety Data Sheets for all chemicals used in this procedure.
- If WHMIS control products must be stored in containers other than their originals, a workplace label must be prepared for the new container. Control products include both pure decanted chemicals and prepared solutions.

Notes

- Square French bottles should be acid-washed for 24 hours, then rinsed with distilled water and allowed to dry prior to use.
- If more or less soil is used in the analysis, adjust the amount of K_2SO_4 added so that the ratio of soil:solution remains 1:2.

Appendix D

Soil Test Analysis Methodologies

Table D.1 Soil test methodologies

Nutrient		Lab	Method
Nitrate	NO ₃ ⁻	AgVise	Water + 0.001 M CaCl ₂
		SDSU	Lachat quikchem method 12-107-04-1-b.(determination of nitrate in 2m KCl soil extracts by flow injection analysis)
Phosphorus	P	AgVise	Olsen P
		SDSU	Lachat QuikChem method (determination of phosphorus in 0.5 m sodium bicarbonate soil extracts by flow injection analysis)
Potassium	K	AgVise	Ammonium acetate method
		SDSU	The displaced Potassium ion was measured using a GBC Avanta Sigma Atomic Absorption Spectrometer
Sulphate	SO ₄	AgVise	Water + 0.001 M CaCl ₂
		SDSU	Lachat quikchem method 12-116-10-1-d. (determination of sulfate by flow injection analysis)
Total Carbon	TC	AgVise	Using leco instrumentation, test TC and test inorganic carbon (CaCO ₃)
		SDSU	The Elementar Vario MAX CNS Macro Elemental Analyzer
Total Organic Carbon	TOC	AgVise	Subtract inorganic carbon from TC
Calcium Carbonate	CaCO ₃	AgVise	
		SDSU	Inorganic Carbon was determined using the Inorganic Carbon by Modified Pressure Calcimeter Method
Electrical Conductivity	EC	AgVise	1:1 extraction
		SDSU	
pH		SDSU	pH was determined using the "SOIL TESTING PROCEDURES" South Dakota State Soil Testing and Plant Analysis Laboratory

Appendix E

Soil Temperature Data

Table E.1 Soil surface temperatures (°C) in the upper slope area of FE1

Date	Block A						Block B					
	U1A	U1D	Δ	U2A	U2C	Δ	U1A	U1D	Δ	U2A	U2C	Δ
a) 2003												
27-May	21.4	20.7	0.7	21.0	18.6	2.4**	20.6	21.7	-1.1	20.9	20.0	1.0**
3-Jun	17.6	17.2	0.4*	17.8	15.6	2.1***	16.9	17.0	0.0	17.0	15.3	1.8***
17-Jun	30.1	30.2	-0.1	30.9	24.3	6.6***	30.3	30.3	0.0	29.6	26.8	2.8**
25-Jun	19.7	19.8	-0.1	19.9	17.9	2.0*	20.8	20.1	0.7	19.8	19.0	0.8
3-Jul	29.4	33.2	-3.8	32.3	28.7	3.6	30.0	33.5	-3.5	32.4	33.2	-0.9
15-Jul	26.9	25.8	1.1	28.3	24.7	3.6	20.5	22.2	-1.7	21.0	21.5	-0.5
31-Jul	29.8	27.2	2.6	30.2	24.7	5.5	23.5	22.6	0.8	24.0	22.0	2.1**
12-Aug	26.4	27.5	-1.1	28.3	25.8	2.5	22.5	22.4	0.1	21.9	22.0	-0.1
17-Sep	10.4	10.7	-0.2	10.6	10.9	-0.3	13.8	13.8	0.0	13.7	13.0	0.7
CV %	28	30		31	29		24	26		25	27	
b) 2004												
21-Apr	19.0	21.6	-2.6	24.7	19.8	4.8	19.1	18.3	0.8	18.8	13.3	5.4**
17-May	24.4	21.7	2.7**	24.8	25.2	-0.4	22.0	18.8	3.2	26.7	23.9	2.8
28-May	26.0	25.7	0.3	25.4	24.5	0.9	23.7	24.2	-0.5	23.7	22.7	1.0
9-Jun	31.0	31.4	-0.4	26.0	29.3	-3.3	29.4	31.5	-2.1	29.0	28.4	0.6
25-Jun	22.4	21.6	0.8	22.4	21.0	1.4	18.3	18.5	-0.2	18.5	17.0	1.5**
15-Jul	33.1	29.0	4.1	28.8	25.5	3.4	30.5	32.2	-1.8	29.5	28.8	0.7
16-Aug	19.5	20.3	-0.8	20.2	19.5	0.7	20.0	20.4	-0.4	19.9	19.8	0.1
CV %	24	18		15	18		24	27		21	27	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table E.2 Soil temperatures (°C) 5 cm below the surface in the upper slope area of FE1

Date	Block A						Block B					
	U1A	U1D	Δ	U2A	U2C	Δ	U1A	U1D	Δ	U2A	U2C	Δ
a) 2003												
27-May	17.1	17.2	-0.1	18.3	16.7	1.6	16.1	17.3	-1.2	16.2	14.7	1.4
3-Jun	15.8	16.3	-0.5	16.7	15.1	1.6***	15.2	15.5	-0.3	15.4	15.2	0.2
17-Jun	23.7	23.5	0.2	23.1	20.4	2.7***	24.9	25.2	-0.3	24.9	20.9	4.0***
25-Jun	18.5	18.2	0.3	18.7	17.6	1.1*	18.1	18.2	-0.1	17.6	17.3	0.3
3-Jul	21.4	20.9	0.5	21.8	19.8	2.0	23.9	24.4	-0.5	24.4	21.6	2.8*
15-Jul	21.6	21.1	0.6	21.4	19.8	1.6**	18.9	18.9	0.0	19.7	19.1	0.6
31-Jul	26.1	25.0	1.1	25.4	23.3	2.1	22.6	23.4	-0.8	23.5	22.1	1.4**
12-Aug	24.0	25.0	-1.1	25.2	22.8	2.4*	21.5	21.4	0.1	21.5	21.3	0.2**
17-Sep	10.6	11.0	-0.4	10.9	11.1	-0.3	12.3	12.5	-0.2	12.2	12.2	0.0
CV %	24	23		22	20		22	21		22	19	
b) 2004												
21-Apr	11.9	11.0	0.9	12.3	9.4	2.8	9.5	9.4	0.1	8.0	6.2	1.8***
17-May	14.3	12.4	1.9	14.1	11.5	2.6	11.5	12.6	-1.1*	12.6	10.2	2.4*
28-May	19.1	19.8	-0.7	19.2	18.4	0.8	14.5	15.1	-0.6	16.1	14.2	1.9
9-Jun	23.6	23.4	0.3	25.4	20.3	5.1	19.9	20.0	-0.1	19.6	18.4	1.2
25-Jun	15.5	17.5	-2.0***	15.8	15.6	0.2	22.4	22.2	0.2	22.7	20.9	1.8
15-Jul	28.3	24.8	3.6	25.8	22.1	3.7	26.3	26.3	-0.1	26.9	24.4	2.5
16-Aug	18.2	18.5	-0.3	18.1	18.2	-0.1	19.1	18.6	0.5*	18.5	18.9	-0.4
CV %	31	29		30	27		34	32		34	38	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table E.3 Soil surface temperatures (°C) in the lower slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	22.4	18.7	3.7***	24.9	13.5	11.4*	20.7	19.3	1.4	22.0	16.2	5.9*
3-Jun	15.6	13.1	2.5***	15.4	11.2	4.1***	16.3	14.9	1.5**	16.3	14.5	1.8**
17-Jun	26.1	20.4	5.7*	25.2	18.5	6.7***	24.6	23.2	1.4	27.3	20.4	6.8**
25-Jun	18.9	15.8	3.1***	18.1	15.6	2.5**	17.3	16.8	0.4	18.2	15.6	2.6*
3-Jul	29.1	22.9	6.2***	28.0	21.0	6.9**	30.0	24.6	5.4	33.2	22.1	11.1***
15-Jul	24.1	18.2	5.8***	23.3	18.4	4.8**	20.9	18.6	2.2*	20.9	17.8	3.1
31-Jul	27.9	21.4	6.5**	27.6	21.5	6.1	27.0	23.6	3.4	26.4	22.1	4.3*
12-Aug	21.4	20.3	1.1**	20.6	19.7	0.9	21.0	21.1	-0.1	21.6	20.0	1.6**
17-Sep	11.0	11.9	-0.9***	11.3	11.7	-0.4	12.7	13.0	-0.3	13.0	13.2	-0.1
CV %	26	21		27	23		28	20		27	19	
b) 2004												
21-Apr	19.6	16.0	19.6	17.6	12.7	4.9*	18.0	12.3	5.6	20.0	15.7	4.3
17-May	18.7	20.2	18.7	19.3	27.5	-8.2	20.2	19.2	1.0*	24.1	19.0	5.1*
28-May	20.1	17.9	20.1	21.5	18.5	2.9	13.7	11.8	1.8	15.8	14.8	1.0
9-Jun	26.9	31.0	26.9	25.8	32.0	-6.2***	27.1	23.1	4.0*	30.0	21.7	8.3
25-Jun	19.2	17.8	19.2	18.3	16.0	2.3*	15.6	16.0	-0.4	15.5	16.5	-0.9
15-Jul	30.6	28.6	30.6	29.7	29.1	0.7	31.7	33.0	-1.3	31.2	32.7	-1.5
16-Aug	24.0	22.1	24.0	23.0	21.8	1.2	23.5	22.4	1.1	22.2	23.6	-1.4
CV %	20	28		19	34		34	37		31	31	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table E.4 Soil temperatures (°C) 5 cm below the surface in the lower slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	18.9	12.6	6.4***	18.1	10.6	7.6**	16.5	13.1	3.5*	16.1	10.5	5.5***
3-Jun	15.1	11.8	3.3**	14.7	10.3	4.4**	15.1	13.7	1.4**	14.8	13.3	1.5*
17-Jun	22.4	15.7	6.7**	21.9	15.4	6.6***	22.7	18.0	4.8**	21.6	18.0	3.6**
25-Jun	17.0	14.3	2.8***	16.5	13.5	2.9**	15.7	15.1	0.6	14.8	14.3	0.5
3-Jul	20.3	16.1	4.2	18.9	15.4	3.5***	24.4	21.4	3.0	25.4	18.9	6.5***
15-Jul	20.2	16.9	3.4***	19.8	17.2	2.6**	17.9	17.1	0.8	17.9	16.8	1.0*
31-Jul	24.4	19.7	4.7**	24.4	19.0	5.4**	25.8	21.8	4.0**	25.2	20.5	4.6*
12-Aug	20.8	18.4	2.4**	20.2	18.4	1.8**	20.0	20.1	-0.1	20.7	19.0	1.7
17-Sep	10.9	12.4	-1.55***	10.9	12.3	-1.4**	12.1	12.7	-0.7	12.2	12.6	-0.4***
CV %	22	18		21	21		25	21		25	21	
b) 2004												
21-Apr	9.9	2.6	7.2***	8.1	4.2	3.9	5.4	6.0	-0.6	3.6	5.4	-1.8
17-May	11.3	9.0	2.4	10.3	7.8	2.5*	11.2	9.8	1.4	9.8	10.0	-0.3
28-May	17.7	9.7	7.9*	13.6	9.5	4.1**	19.6	21.0	-1.4	16.4	18.6	-2.2
9-Jun	22.8	15.5	7.3**	21.4	16.1	5.3**	16.8	19.2	-2.4	15.8	17.2	-1.4
25-Jun	16.2	12.3	3.9**	15.0	12.5	2.5*	18.8	19.0	-0.2	18.4	19.3	-0.9
15-Jul	28.7	20.5	8.1***	26.1	20.1	6.1**	26.4	26.5	-0.1	24.4	26.7	-2.3
16-Aug	21.4	17.2	4.2***	19.7	18.7	1.0	21.2	20.7	0.5	21.1	21.8	-0.7
CV %	37	46		38	45		40	40		46	42	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table E.5 Soil temperatures (°C) in the upper slope area of FE2 in 2004

Treatment	21-Apr	17-May	28-May	9-Jun	25-Jun	15-Jul	16-Aug	CV %
<i>a) surface</i>								
Addition	28.1	23.8a	18.8	20.0	20.6	32.5	21.6	21.0
Control	27.4	19.3b	18.9	21.1	19.6	31.2	19.9	21.4
Disturbed	28.8	23.2a	18.9	19.7	20.3	30.8	20.7	20.3
<i>b) 5 cm below the surface</i>								
Addition	11.2a	9.0a	11.7b	14.5	14.8	26.9	19.2	39.9
Control	8.7b	7.1b	11.1c	14.3	14.2	25.0	17.5	43.5
Disturbed	12.7a	8.8a	12.6a	14.9	14.5	24.9	18.4	33.9

a-b Mean values followed by the same letter (within columns) are not significantly different

Note: LSD groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$; values indicate the means of 9 replicates

Appendix F

Soil Gravimetric Moisture Content Data

Table F.1 Soil gravimetric moisture content (%) in the upper slope area of FE1

Date	Block A						Block B					
	U1A	UID	Δ	U2A	U2C	Δ	U1A	UID	Δ	U2A	U2C	Δ
a) 2003												
27-May	35	26	9	31	28	3	27	24	3	25	24	1**
17-Jun	26	21	5	27	25	2	29	24	5***	27	23	3*
3-Jul	26	21	5	27	25	2	24	21	3*	24	19	5
31-Jul	15	15	0	15	16	-1	18	17	2	16	19	-3
17-Sep	31	29	2	31	28	4	28	24	3	30	25	5***
CV %	23	24		19	29		19	15		21	14	
b) 2004												
21-Apr	30	30	0	29	30	-1	25	23	2*	24	26	-1
17-May	27	28	-1	33	31	2*	25	24	1	19	26	-7*
28-May	29	27	1	30	26	4	29	26	3	27	24	3
9-Jun	31	28	3*	33	29	3	28	22	6***	26	22	4***
25-Jun	28	24	3	27	27	1	22	21	2	24	25	0
15-Jul	43	33	10	33	39	-6	30	27	3*	30	28	2
16-Aug	22	23	-1	24	29	-6	24	23	0	24	23	1
CV %	23	13		14	19		12	12		14	16	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table F.2 Soil gravimetric moisture content (%) in the lower slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	33	41	-8	45	38	6	48	35	12*	50	35	15****
17-Jun	31	34	-3	39	34	5	37	27	10	44	24	20****
3-Jul	35	41	-6	47	38	9	45	25	20**	49	26	23****
31-Jul	31	24	7	35	23	12**	39	25	14**	44	26	18*
17-Sep	26	35	-8	35	39	-4	34	33	1	39	32	7**
CV %	24	20		20	11		16	27		13	17	
b) 2004												
21-Apr	33	38	-5	30	38	-8**	32	41	-9*	26	33	-7
17-May	38	48	-10****	41	45	-4*	29	40	-11**	32	45	-13
28-May	29	40	-11	38	44	-6*	38	38	0	36	49	-13
9-Jun	37	45	-8	43	44	-1	37	41	-4	37	42	-5
25-Jun	40	38	2	42	48	-6	34	41	-7	30	49	-19
15-Jul	32	98.7	-66*	40	61	-21	36	43	-7	42	47	-5
16-Aug	32	33	-1	36	34	2	33	40	-7	33	43	-11
CV %	16	49		12	25		27	15		24	21	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ****Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table F.3 Soil gravimetric moisture content (%) in the upper slope area of FE2 in 2004

Treatment	21-Apr	17-May	28-May	9-Jun	25-Jun	15-Jul	16-Aug	CV %
Addition	27a	27	31	30a	29	34	29	8
Control	25a	26	28	26b	28	36	28	12
Disturbed	21b	26	28	29ab	27	36	29	16

a-b Mean values followed by the same letter (within columns) are not significantly different

Note: LSD groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$; values indicate the means of 9 replicates

Appendix G

Greenhouse Gas Flux Data

Table G.1 Soil respiration (carbon dioxide flux) (mg CO₂-C m⁻² hr⁻¹) from the upper slope area of FE1												
Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	0.5	0.2	0.3	0.2	0.5	-0.3	11.2	5.7	5.5	10.6	7.0	3.7
3-Jun	12.7	4.8	7.9	9.8	9.6	0.2	12.4	4.4	8.0	11.0	11.6	-0.5**
17-Jun	41.3	10.1	31.2	21.8	21.5	0.3	18.9	15.9	3.0	14.8	28.1	-13.3
25-Jun	27.0	9.0	18.0	9.5	16.7	-7.2	19.6	13.7	5.9	15.3	21.3	-6.0
3-Jul	26.4	15.1	11.3	22.1	29.8	-7.6	25.0	23.5	1.5	22.0	26.9	-4.9
15-Jul	16.0	16.4	-0.3	24.8	41.2	-16.4**	33.3	40.9	-7.5	32.5	48.8	-16.2*
31-Jul	37.1	16.5	20.7	26.6	46.2	-19.6*	17.4	21.5	-4.2	22.0	45.8	-23.8*
12-Aug	10.4	6.2	4.2	12.3	9.8	2.6	3.8	3.3	0.5	6.1	5.6	0.4
17-Sep	11.9	11.3	0.6	10.0	17.7	-7.7***	6.3	5.1	1.2	6.7	9.4	-2.8
CV %	97	69		85	75		63	85		65	76	
b) 2004												
21-Apr	7.6	6.6	1.0	5.1	4.4	0.68*	3.7	3.7	0.0	4.7	3.2	1.51*
17-May	12.9	9.8	3.05*	9.7	6.2	3.5	3.3	4.9	-1.6	3.3	5.7	-2.42**
28-May	15.5	10.5	4.9	22.6	20.0	2.6	6.4	6.9	-0.5	8.1	13.8	-5.8
9-Jun	18.6	19.3	-0.7	15.3	21.6	-6.3	8.3	6.7	1.6	11.1	13.9	-2.8
25-Jun	6.9	11.3	-4.4	6.4	11.0	-4.6	4.9	3.8	1.1	4.6	9.1	-4.5
15-Jul	80.6	81.5	-0.8	92.1	80.6	11.4	45.2	47.8	-2.6	45.4	46.5	-1.1
16-Aug	38.4	38.1	0.3	33.3	37.7	-4.5	27.9	30.3	-2.3	28.5	33.8	-5.3
CV %	102	105		117	114		117	112		102	90	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table G.2 Soil respiration (carbon dioxide flux) ($\text{mg CO}_2\text{-C m}^{-2} \text{hr}^{-1}$) from the lower slope area of FE1

Date	Block A						Block B					
	LIR	LID	Δ	L2R	L2C	Δ	LIR	LID	Δ	L2R	L2C	Δ
a) 2003												
27-May	1.8	3.6	-1.8	1.3	2.1	-0.8	35.1	32.2	2.9	32.9	25.1	7.8
3-Jun	16.6	46.2	-29.6	17.7	33.1	-15.4	16.6	10.5	6.1	23.2	11.2	11.9
17-Jun	53.8	51.4	2.4	41.0	34.9	6.1	37.6	28.2	9.4**	35.4	32.1	3.3
25-Jun	33.8	30.7	3.0	29.8	31.6	-1.8	17.7	24.9	-7.2	398.9	32.1	366.9
3-Jul	60.1	55.6	4.5	61.0	78.0	-17.0	54.1	58.1	-4.0	71.7	62.2	9.5
15-Jul	41.1	52.9	-11.8	41.7	67.3	-25.6	48.7	32.4	16.3	43.0	43.2	-0.2
31-Jul	61.4	87.3	-25.9*	46.9	92.6	-45.7**	78.0	38.5	39.5**	57.3	54.2	3.1
12-Aug	11.8	27.9	-16.1	17.4	50.1	-32.8*	27.5	28.9	-1.5	39.3	39.3	0.0
17-Sep	16.7	29.0	-12.3	8.8	31.0	-22.2**	19.9	14.4	5.5	15.5	20.2	-4.7
CV %	78	64		80	66		62	62		272	66	
b) 2004												
21-Apr	13.3	19.0	-5.7**	9.3	14.9	-5.6**	10.4	6.1	4.3	7.9	5.6	2.2
17-May	5.5	32.2	-26.7**	6.3	29.4	-23.1***	16.1	9.0	7.0	13.7	8.2	5.5
28-May	15.0	25.1	-10.0	13.9	7.6	6.3	30.8	14.3	16.5	18.0	13.3	4.7
9-Jun	21.5	46.0	-24.5**	22.3	40.6	-18.3*	29.5	16.1	13.4	28.4	23.9	4.5
25-Jun	25.0	46.6	-21.6**	27.6	51.3	-23.7	33.9	29.3	4.6	42.9	39.2	3.7
15-Jul	56.6	101.7	-45.4**	71.1	66.8	4.3	85.8	78.6	7.2	91.2	83.2	8.0
16-Aug	64.8	111.5	-46.7	54.3	98.1	-43.8*	61.8	41.0	20.8	76.9	63.0	13.9
CV %	87	66		85	81		75	90		87	92	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table G.3 Soil methane flux ($\mu\text{g CH}_4\text{-C m}^{-2} \text{ hr}^{-1}$) from the upper slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	4.6	3.7	1.0	5.9	10.8	-4.9	1.4	-1.3	2.7	0.5	4.7	-4.2
3-Jun	-0.3	-2.1	1.8	-1.4	-5.7	4.3	1.0	-1.8	2.9	3.3	0.9	2.4
17-Jun	-0.5	-0.5	0.0	-0.3	-0.1	-0.2	0.5	-0.8	1.3	-1.6	-1.3	-0.3
25-Jun	-0.4	-1.3	1.0	-0.6	-1.3	0.8	-3.7	-0.2	-3.5	-0.6	4.0	-4.6
3-Jul	-4.8	-0.4	-4.4	-1.2	-4.5	3.3	-21.6	-25.2	3.6	-19.0	-39.1	20.0**
15-Jul	-1.7	-4.7	3.0	-3.3	-14.1	10.7	-0.8	-0.4	-0.4	1.4	-0.4	1.8
31-Jul	-13.5	-1.1	-12.5**	-0.9	-7.2	6.2***	-3.9	-4.9	1.0	-1.5	1.8	-3.3
12-Aug	-12.7	-5.9	-6.8***	-9.6	-15.6	6.0	-3.6	-9.0	5.4	-7.6	-10.4	2.8
17-Sep	-1.4	0.0	-1.4	-2.4	5.2	-7.6	2.3	5.8	-3.5	1.4	1.4	0.0
CV %	200	256		348	278		265	225		299	334	
b) 2004												
21-Apr	-6.7	-7.2	0.6*	-3.0	-1.5	-1.5	-3.9	-5.0	1.1	-5.7	-2.7	-3.0
17-May	-5.3	-2.8	-2.5	-3.1	-3.1	0.0	-3.6	-5.6	1.9	-2.7	-6.4	3.7**
28-May	-18.2	-23.4	5.2	-18.5	-23.3	4.8	-5.0	-6.6	1.7	-3.8	-3.3	-0.5
9-Jun	-0.9	-1.9	0.9	-2.5	-2.7	0.2	-1.5	-1.6	0.1	-2.2	-2.2	0.0
25-Jun	-4.0	-7.3	3.4	-2.4	-2.1	-0.2	4.2	-0.4	4.6	3.4	-1.0	4.4
15-Jul	1.1	-4.9	6.0	-1.8	-2.0	0.3	-1.3	-2.3	1.0	-1.9	-0.2	-1.7
16-Aug	-9.9	-15.8	5.9*	-10.0	-10.9	0.9	-7.2	-8.2	1.0	-8.3	-6.1	-2.2
CV %	103	87		114	133		147	83		152	102	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table G.4 Soil methane flux ($\mu\text{g CH}_4\text{-C m}^{-2} \text{hr}^{-1}$) from the lower slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	1.3	-3.8	5.1**	0.5	-10.5	11.0	-0.8	-0.5	-0.4	-0.7	-1.8	1.1***
3-Jun	-1.2	-6.1	4.9	-3.1	0.6	-3.7	-1.2	-1.7	0.5	-0.9	-2.1	1.1
17-Jun	-2.0	-2.2	0.2	-1.1	-2.7	1.7	0.7	0.1	0.6	0.5	0.8	-0.4
25-Jun	-1.7	1.5	-3.3	-0.3	-1.4	1.1	55.3	0.1	55.2	-1.8	2.3	-4.1
3-Jul	-1.2	-2.1	0.9	-3.7	-7.4	3.6	-4.8	-12.5	7.7	-22.9	-5.2	-17.7
15-Jul	-2.0	6.1	-8.2	0.1	-0.6	0.7	-9.5	-5.1	-4.3	-2.3	-6.3	3.9
31-Jul	-21.2	-17.7	-3.5	-17.4	-18.9	1.5	1.0	-2.5	3.6	0.1	-2.7	2.8
12-Aug	2.3	3.7	-1.4	-4.0	2.2	-6.2	-4.8	-5.1	0.3	-5.7	-2.2	-3.5
17-Sep	1.5	4.6	-3.2	2.0	11.4	-9.4*	7.6	2.1	5.5	9.7	-0.7	10.4***
CV %	319	385		266	465		682	259		377	219	
b) 2004												
21-Apr	-3.1	-0.9	-2.2	-2.0	0.9	-2.9***	-6.6	-5.5	-1.1	-1.4	-7.1	5.7
17-May	0.7	-2.5	3.2	0.3	-1.8	2.2	-17.5	-16.4	-1.1	-20.9	-19.3	-1.6
28-May	44.6	-32.9	77.5	62.0	74.9	-12.9	-0.5	-5.6	5.1	-1.8	-5.0	3.2
9-Jun	-5.8	-6.0	0.3	-5.1	-7.6	2.4	-4.6	-1.5	-3.1	-2.1	-3.8	1.7
25-Jun	-2.2	-0.4	-1.8	-2.7	-6.4	3.7***	-5.2	-5.8	0.5	-6.9	7.9	-14.8
15-Jul	3.6	0.1	3.4*	-0.4	6.2	-6.6	4.9	3.6	1.3	5.0	1.5	3.5
16-Aug	-10.1	-11.1	1.0	-7.5	-10.4	2.9	-8.7	-5.5	-3.2	-6.7	-6.6	-0.1
CV %	973	162		942	893		127	121		166	243	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table G.5 Soil nitrous oxide flux ($\mu\text{g N}_2\text{O-N m}^{-2} \text{hr}^{-1}$) from the upper slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	6.4	6.0	0.4	3.4	34.6	-31.2	2.8	0.9	1.9	2.9	0.6	2.3
3-Jun	10.9	2.2	8.7**	11.9	10.9	1.0	3.3	1.1	2.2	6.4	1.1	5.2
17-Jun	23.5	9.9	13.7	12.2	13.8	-1.6	10.7	2.6	8.1	3.8	3.6	0.1
25-Jun	9.3	2.3	7.0	1.6	1.9	-0.2	5.5	2.6	2.8	2.9	3.2	-0.3
3-Jul	4.9	1.0	3.8	2.6	1.6	1.0	2.6	2.8	-0.3	2.0	1.3	0.7
15-Jul	3.7	1.9	1.8	3.6	3.0	0.6	1.8	1.2	0.5	1.5	2.4	-0.9
31-Jul	11.2	1.6	9.6	3.3	2.5	0.7	0.5	3.4	-2.9	0.7	0.8	0.0
12-Aug	2.8	1.6	1.2	1.6	0.8	0.8	-0.7	-0.5	-0.1	-0.3	-0.9	0.6*
17-Sep	6.3	6.5	-0.2	3.7	4.3	-0.6	1.1	1.4	-0.3	1.3	0.5	0.8
CV %	111	136		137	181		128	116		143	122	
b) 2004												
21-Apr	6.6	4.2	2.4	4.7	4.2	0.6	1.2	2.2	-1.0***	3.3	1.9	1.4
17-May	6.1	1.7	4.4	3.7	1.1	2.6	1.1	2.4	-1.4	0.7	4.5	-3.8*
28-May	-0.7	-1.0	0.3	-0.3	-0.3	0.0	31.2	30.1	1.2	53.4	101.8	-48.4
9-Jun	87.3	36.9	50.5	55.8	35.5	20.3	65.3	8.3	57.1	34.7	70.8	-36.1
25-Jun	3.2	6.2	-3.0	3.0	4.0	-1.0	35.8	-3.2	39.0	27.6	5.3	22.3
15-Jul	44.4	7.5	36.8	10.5	5.3	5.2	1.9	0.4	1.5*	0.7	1.6	-0.9
16-Aug	2.2	2.1	0.1	1.8	1.9	-0.1	2.6	2.4	0.3	1.3	2.6	-1.3
CV %	179	164		189	220		222	221		160	187	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table G.6 Soil nitrous oxide flux ($\mu\text{g N}_2\text{O-N m}^{-2} \text{ hr}^{-1}$) from the lower slope area of FE1

Date	Block A						Block B					
	L1R	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
27-May	0.7	0.7	0.0	3.0	0.7	2.3	4.6	4.3	0.4	4.7	11.2	-6.4
3-Jun	3.7	0.1	3.6	7.5	0.9	6.6	2.6	5.0	-2.5	7.2	12.9	-5.7
17-Jun	3.4	-0.9	4.3	5.9	-1.6	7.5	1.9	3.9	-2.0	6.4	6.2	0.2
25-Jun	3.5	0.8	2.68*	6.7	-0.3	7.1	4.7	3.6	1.1	773.0	3.5	769.5
3-Jul	2.5	0.3	2.21*	1.0	-0.1	1.0	11.5	4.4	7.1	22.4	15.3	7.2
15-Jul	3.6	-0.2	3.8	0.7	-0.1	0.8	3.8	0.7	3.1	3.4	6.2	-2.8
31-Jul	0.5	1.2	-0.7	1.1	3.0	-2.0	2.2	0.9	1.3	2.4	5.3	-2.9
12-Aug	0.1	1.1	-1.00**	0.4	0.7	-0.30*	-1.1	-1.4	0.3	-0.7	1.5	-2.2
17-Sep	1.9	2.1	-0.2	1.4	2.9	-1.43**	1.8	1.5	0.4	1.7	1.3	0.4
CV %	102	271		174	295		182	168		485	144	
b) 2004												
21-Apr	1.1	1.5	-0.38*	1.0	1.7	-0.7	2.1	0.9	1.2	1.0	1.1	-0.1
17-May	0.3	1.2	-0.9	0.4	1.4	-1.0	-2.7	-2.7	0.1	-2.6	-2.9	0.3
28-May	15.9	0.0	15.9	20.6	24.7	-4.1	0.5	0.3	0.2	0.2	0.0	0.1
9-Jun	0.3	2.4	-2.2	0.1	1.2	-1.1	2.0	0.8	1.21*	2.9	0.5	2.33**
25-Jun	-2.0	66.0	-68.0	-0.1	0.4	-0.5	0.0	-0.5	0.5	-0.2	-0.8	0.6
15-Jul	1.9	1.9	0.0	0.5	4.5	-4.0	2.9	0.9	2.0	3.2	1.4	1.76**
16-Aug	0.7	2.4	-1.7	0.8	1.8	-1.0	1.6	0.2	1.4	0.7	0.4	0.3
CV %	418	393		378	272		222	4413		266	6055	

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table G.7 Soil gas flux in the upper slope area of FE2 in 2004

Treatment	21-Apr	17-May	28-May	9-Jun	25-Jun	15-Jul	16-Aug	CV %
a) CO_2 ($mg\ CO_2-C\ m^{-2}\ hr^{-1}$)								
Addition	4.0 ^a	4.1 ^a	3.9	5.5	2.9	20.8 ^b	24.5	123
Control	1.1 ^b	1.4 ^b	3.3	4.7	2.7	60.5 ^a	30.7	157
Disturbed	2.5 ^a	3.1 ^a	2.3	3.0	2.5	24.0 ^b	23.8	140
b) CH_4 ($\mu g\ CH_4-C\ m^{-2}\ hr^{-1}$)								
Addition	-3.0 ^b	-3.7 ^b	-2.5	-3.2	-1.6	-5.8	3.6 ^b	160
Control	-0.1 ^a	-0.5 ^a	-0.5	-4.1	-1.7	-4.4	-6.6 ^a	160
Disturbed	-3.3 ^b	-3.9 ^b	-1.7	-3.1	-1.4	-5.4	-4.5 ^b	114
c) N_2O ($\mu g\ N_2O-N\ m^{-2}\ hr^{-1}$)								
Addition	1.6	1.0	5.2	29.3	0.3	8.4	2.8	351
Control	0.2	-0.1	9.6	27.5	7.9	10.4	1.9	287
Disturbed	0.8	1.1	3.1	4.1	17.3	36.4	2.1	403

a-b Mean values followed by the same letter (within columns) are not significantly different

Note: LSD groupings are associated with significant treatment effects found using linear mixed ANOVA and $P < 0.10$; values indicate the means of 9 replicates

Appendix H

Calculation of the Expected Percent Increase in Organic Carbon in the Addition Soil (FE1)

Upper Slope Area - Actual Percent Organic Carbon

Block A

Depth	Upper Slope Control Soil	Addition Soil (mean of U1A and U2A)
0	3.8%	5.1%
15 cm	3.1%	2.0%
30 cm		

Block B

Depth	Upper Slope Control Soil	Addition Soil (mean of U1A and U2A)
0	2.9%	3.7%
15 cm	2.1%	2.6%
30 cm		

Lower Slope Area - Actual Percent Organic Carbon

Block A

Depth	Lower Slope Control Soil
0	6.9%
15 cm	4.2%
30 cm	

Block B

Depth	Lower Slope Control Soil
0	5.6%
15 cm	4.9%
30 cm	

Calculation of the o.c. (%) of the soil that was added to the hilltop: 20 cm of soil was removed; therefore, the 0-15 cm surface layer was removed plus one third of the 15-30 cm layer was removed. The O.C. (%) that was added to the upper slope areas was:

Block A: $((3 \times 6.9\%) + (4.2\%)) / 4 = 6.2\%$

Block B: $((3 \times 5.6\% + (4.9\%)) / 4 = 5.4\%$

The 20 cm of soil was added to a larger area to give an added depth of 10 cm. When the o.c. was measured, the top 15 cm of soil was sampled and therefore the 10 cm of added material would have been diluted with 5 cm of the original surface material. **The**

Block A: $((2 \times 6.2\%) + 3.8) / 3 = 5.4\%$

Block B: $((2 \times 5.4\%) + 2.9) / 3 = 4.5\%$

Appendix I

Crop and Weed Emergence Data for FE2

Table I.1 Crop and weed emergence (plants m⁻²)

Block	Plot No.	Treatment	Crop Emergence	Weed Emergence
1	1	ADD	14	14
	2	CON	30	7
	3	DIST	17	4
2	4	DIST	18	10
	5	CON	20	7
	6	ADD	47	13
3	7	ADD	13	31
	8	DIST	4	19
	9	CON	61	10
4	10	DIST	16	11
	11	CON	42	20
	12	ADD	30	4
5	13	CON	8	26
	14	ADD	25	10
	15	DIST	31	3
6	16	DIST	52	2
	17	CON	39	15
	18	ADD	34	25
7	19	ADD	1	11
	20	DIST	23	11
	21	CON	53	8
8	22	CON	69	14
	23	DIST	23	36
	24	ADD	56	10
9	25	CON	14	9
	26	ADD	4	35
	27	DIST	21	11

Appendix J

Soil Seed Bank Experiment Data with Treatment Means and Standard Deviations




Figure J.1 Mean total plant emergence by site and soil depth. Error bars represent standard deviations.




Figure J.2 Mean total species emergence by site and soil depth. Error bars represent standard deviations.

Table J.1 Soil seed bank experiment data with plant and species emergence, means and standard deviations

Tray #	Site	Replicate	Depth	Plants per kg soil	Mean plants per kg soil	Standard Deviation	Species per kg soil	Mean species per kg soil	Standard Deviation
10	A	01	0-5	16	34	16	4	6	2
26	A	02	0-5	46			7		
5	A	03	0-5	40			8		
8	A	01	5-10	3	6	3	2	3	1
16	A	02	5-10	9			3		
12	A	03	5-10	5			3		
27	A	01	10-15	0	3	3	0	1	1
11	A	02	10-15	6			3		
24	A	03	10-15	1			1		
17	A	01	15-20	0	0	1	0	0	1
6	A	02	15-20	1			1		
25	A	03	15-20	0			0		
3	A	01	20-25	0	0	0	0	0	0
21	A	02	20-25	1			1		
7	A	03	20-25	0			0		
36	B	01	0-5	74	52	30	4	5	1
19	B	02	0-5	64			4		
28	B	03	0-5	18			6		
22	B	01	5-10	11	7	3	3	3	1
44	B	02	5-10	8			2		
38	B	03	5-10	4			3		
15	B	01	10-15	6	5	2	3	2	1
2	B	02	10-15	8			1		
18	B	03	10-15	3			2		
45	B	01	15-20	3	2	1	1	1	0
41	B	02	15-20	1			1		
23	B	03	15-20	3			1		
30	B	01	20-25	1	1	0	1	1	0
20	B	02	20-25	1			1		
33	B	03	20-25	1			1		
43	C	01	0-5	5	8	7	5	3	2
29	C	02	0-5	16			3		
40	C	03	0-5	3			2		
13	C	01	5-10	1	1	0	1	1	0
37	C	02	5-10	1			1		
39	C	03	5-10	1			1		
14	C	01	10-15	0	0	0	0	0	0
31	C	02	10-15	0			1		
4	C	03	10-15	0			0		
42	C	01	15-20	1	0	0	1	0	0
34	C	02	15-20	1			1		
1	C	03	15-20	0			0		
35	C	01	20-25	0	1	1	0	0	0
9	C	02	20-25	1			1		
32	C	03	20-25	1			1		

Appendix K

Core Method for Bulk Density Determination

Equipment

- aluminum soil cores
- trowel/shovel
- knife/spatula
- scale

In the Field

1. Record the number of core being used
2. Place insertion ring on top of core and press into soil with foot
 - if insertion ring is not available, another aluminum core can be substituted
3. Remove insertion ring
4. Use a trowel to extract core, keeping soil intact on both ends
5. Square ends with a knife or spatula, and place core contents in a plastic bag
6. Tie plastic bag and place in a cooler with ice packs

In the Lab

1. Weigh bag, core and soil and record mass in grams
2. Weigh several bags to determine an approximate mass per bag
3. Subtract mass of bag from mass of soil
4. Determine Gravimetric Moisture Content as per the GMC SOP

Calculations

- calculate mass of soil and core volume
- mass of dry soil = (mass of full core – mass of empty core)/(1+GMC)
- volume of core = $\pi r^2 h$
- $r = 2.54\text{cm}$ for all cores (diameter = 2")
- $h =$ length in data file (convert length to cm before entering it is the spreadsheet calculation)
- $BD =$ mass of oven dry soil (g)/volume of core (cm^3)

Original Author: J.L.B. Culley

Source: Chapter 50, Soil Sampling and Methods of Analysis, M.R. Carter, Ed., 1993

Appendix L

Nitrous Oxide, Carbon Dioxide and Methane Flux Equations

L.1 Nitrous Oxide Flux

$$\frac{1 \mu\text{LN}_2\text{O}}{\text{L} \cdot \text{hr}} \times \frac{323.7(\text{cm}^2) \cdot h(\text{cm})}{0.0324\text{m}^2} \times \frac{1\text{L}}{1000\text{cm}^3} = 9.99 \cdot h(\text{cm}) \frac{\mu\text{LN}_2\text{O}}{\text{m}^2 \cdot \text{hr}}$$

$$9.99 \cdot h(\text{cm}) \frac{\mu\text{LN}_2\text{O}}{\text{m}^2 \cdot \text{hr}} \times \frac{1 \mu\text{mole N}_2\text{O}}{0.0821 \cdot T(\text{K}) \mu\text{LN}_2\text{O}} \times \frac{44 \mu\text{g N}_2\text{O}}{1 \mu\text{mole N}_2\text{O}} = 5,353 \frac{h(\text{cm}) \mu\text{g N}_2\text{O}}{T(\text{K}) \text{m}^2 \cdot \text{h}}$$

$$5,353 \frac{h(\text{cm}) \mu\text{g N}_2\text{O}}{T(\text{K}) \text{m}^2 \cdot \text{h}} \times \frac{10^3 \text{ng N}_2\text{O}}{1 \mu\text{g N}_2\text{O}} \times \frac{1\text{h}}{60\text{min}} \times \frac{1\text{min}}{60\text{sec}} = 1487 \frac{h(\text{cm}) \text{ng N}_2\text{O}}{T(\text{K}) \text{m}^2 \cdot \text{sec}}$$

$$5,353 \frac{h(\text{cm}) \mu\text{g N}_2\text{O}}{T(\text{K}) \text{m}^2 \cdot \text{h}} \times \frac{1 \text{g N}_2\text{O}}{10^6 \mu\text{g N}_2\text{O}} \times \frac{28 \text{g N}}{44 \text{g N}_2\text{O}} \times \frac{10^4 \text{m}^2}{1\text{ha}} \times \frac{24\text{h}}{1\text{d}} = 817.6 \frac{h(\text{cm}) \text{g N}_2\text{O} - \text{N}}{T(\text{K}) \text{ha} \cdot \text{d}}$$

L.2 Carbon Dioxide Flux

$$\frac{1 \mu\text{LCO}_2}{\text{L} \cdot \text{hr}} \times \frac{323.7(\text{cm}^2) \cdot h(\text{cm})}{0.0324\text{m}^2} \times \frac{1\text{L}}{1000\text{cm}^3} = 9.99 \cdot h(\text{cm}) \frac{\mu\text{LCO}_2}{\text{m}^2 \cdot \text{hr}}$$

$$9.99 \cdot h(\text{cm}) \frac{\mu\text{LCO}_2}{\text{m}^2 \cdot \text{hr}} \times \frac{1 \mu\text{mole CO}_2}{0.0821 \cdot T(\text{K}) \mu\text{LCO}_2} \times \frac{44 \mu\text{g CO}_2}{1 \mu\text{mole CO}_2} = 5,353 \frac{h(\text{cm}) \mu\text{g CO}_2}{T(\text{K}) \text{m}^2 \cdot \text{h}}$$

$$5,353 \frac{h(\text{cm}) \mu\text{g CO}_2}{T(\text{K}) \text{m}^2 \cdot \text{h}} \times \frac{1\text{h}}{60\text{min}} \times \frac{1\text{min}}{60\text{sec}} = 1.487 \frac{h(\text{cm}) \mu\text{g CO}_2}{T(\text{K}) \text{m}^2 \cdot \text{sec}}$$

$$5,353 \frac{h(\text{cm}) \mu\text{g CO}_2}{T(\text{K}) \text{m}^2 \cdot \text{h}} \times \frac{1 \text{kg CO}_2}{10^9 \mu\text{g CO}_2} \times \frac{12 \text{kg C}}{44 \text{kg CO}_2} \times \frac{10^4 \text{m}^2}{1\text{ha}} \times \frac{24\text{h}}{1\text{d}} = 0.3504 \frac{h(\text{cm}) \text{kg CO}_2 - \text{C}}{T(\text{K}) \text{ha} \cdot \text{d}}$$

L.3 Methane Flux

$$\frac{1 \text{ uL CH}_4}{\text{L} \cdot \text{hr}} \times \frac{323.7(\text{cm}^2) \cdot h(\text{cm})}{0.0324 \text{ m}^2} \times \frac{1 \text{ L}}{1000 \text{ cm}^3} = 9.99 \cdot h(\text{cm}) \frac{\text{uL CH}_4}{\text{m}^2 \cdot \text{hr}}$$

$$9.99 \cdot h(\text{cm}) \frac{\text{uL CH}_4}{\text{m}^2 \cdot \text{hr}} \times \frac{1 \text{ umole CH}_4}{0.0821 \cdot T(\text{K}) \text{ uL CH}_4} \times \frac{16 \text{ ug CH}_4}{1 \text{ umole CH}_4} = 1,947 \frac{h(\text{cm})}{T(\text{K})} \frac{\text{CH}_4}{\text{m}^2 \cdot \text{h}}$$

$$1,947 \frac{h(\text{cm})}{T(\text{K})} \frac{\text{ug CH}_4}{\text{m}^2 \cdot \text{h}} \times \frac{10^3 \text{ ng CH}_4}{1 \text{ ug CH}_4} \times \frac{1 \text{ h}}{60 \text{ min}} \times \frac{1 \text{ min}}{60 \text{ sec}} = 540.8 \frac{h(\text{cm})}{T(\text{K})} \frac{\text{ng CH}_4}{\text{m}^2 \cdot \text{sec}}$$

$$1,947 \frac{h(\text{cm})}{T(\text{K})} \frac{\text{ug CH}_4}{\text{m}^2 \cdot \text{h}} \times \frac{1 \text{ g CH}_4}{10^6 \text{ ug CH}_4} \times \frac{12 \text{ g C}}{16 \text{ g CH}_4} \times \frac{10^4 \text{ m}^2}{1 \text{ ha}} \times \frac{24 \text{ h}}{1 \text{ d}} = 350.5 \frac{h(\text{cm})}{T(\text{K})} \frac{\text{g CH}_4 - \text{C}}{\text{ha} \cdot \text{d}}$$

Appendix M

Bulk Density Measurement Data

Table M.1 Bulk density in the upper slope area of FE1, July 15, 2004

Nutrient	Block A ^a						Block B					
	U1A ^b	UID	Δ	U2A	U2C	Δ	U1A	UID	Δ	U2A	U2C	Δ
<i>a) 0-10 cm Upper Slope Area</i>												
Bulk Density (g cm ⁻³)	0.86	0.97	-0.10	0.90	1.04	-0.14	0.95	0.98	-0.03	0.92	0.91	0.01

^a Flax in Block A and Peas in Block B

^b U1A = Addition treatment pair 1; UID = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table M.2 Bulk density in the lower slope area of FE1, July 15, 2004

Nutrient	Block A ^a						Block B					
	L1R ^b	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ

a) 0-10 cm Lower Slope Area

Bulk Density (g cm ⁻³)	1.25	0.36	0.89	1.17	0.57	0.60	1.03	0.93	0.10	0.75	0.78	-0.02
------------------------------------	------	------	------	------	------	------	------	------	------	------	------	-------

^a Flax in Block A and Peas in Block B

^bU1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

**Table M.3 Bulk density in
the upper slope are of FE2,
July 15, 2004**

	Bulk Density (g cm ⁻³)
Addition	0.86
Control	0.84
Disturbed	0.86

Appendix N

Cumulative Greenhouse Gas Flux Data

Table N.1 Cumulative greenhouse gas flux for the upper slope area of FE1 over a 115 day sampling period in 2003 and over a 118 day sampling period in 2004

	Block A ^a						Block B					
	U1A ^b	UID	Δ	U2A	U2C	Δ	U1A	UID	Δ	U2A	U2C	Δ
a) 2003												
CO ₂ (mg CO ₂ -C m ⁻²)	2295	1211	1084	1852	2570	-718	1816	1710	106	1845	2654	-809
CH ₄ (μg CH ₄ -C m ⁻²)	-591	-252	-339**	-350	-654	304	-345	-458	113**	-222	-451	673
N ₂ O (μg N ₂ O-N m ⁻²)	919	394	525***	517	613	-96	263	164	99	216	150	66
a) 2004												
CO ₂ (mg CO ₂ -C m ⁻²)	3633	3611	22	3780	3417	363	2026	2125	-99	2107	2433	-326
CH ₄ (μg CH ₄ -C m ⁻²)	-616	-964	348**	-600	-595	-5	-275	-473	198	-329	-348	19
N ₂ O (μg N ₂ O-N m ⁻²)	2664	935	1729	1265	801	464	2047	530	1517	1694	2452	-758

^a Flax in Block A and Peas in Block B

^bU1A = Addition treatment pair 1; UID = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table N.2 Cumulative greenhouse gas flux for the lower slope area of FE1 over a 115 day sampling period in 2003 and over a 118 day sampling period in 2004

	Block A ^a						Block B					
	L1R ^b	L1D	Δ	L2R	L2C	Δ	L1R	L1D	Δ	L2R	L2C	Δ
a) 2003												
CO ₂ (mg CO ₂ -C m ⁻²)	3685	5068	-1383	3324	5828	-2504*	4236	3305	931	4170	4158	12
CH ₄ (μg CH ₄ -C m ⁻²)	-296	-142	-154	-383	-152	-231	295	-335	630	-229	-246	17
N ₂ O (μg N ₂ O-N m ⁻²)	228	83	145***	282	105	177	312	203	109	571	649	-78
a) 2004												
CO ₂ (mg CO ₂ -C m ⁻²)	3746	7166	-3420*	3966	5781	-1815	5009	3887	1122	5376	4626	750*
CH ₄ (μg CH ₄ -C m ⁻²)	289	-598	878	191	310	-119	-641	-565	-76	-561	-494	-67
N ₂ O (μg N ₂ O-N m ⁻²)	231	1357	-1126	204	415	-211	111	-8	119***	99	-1	100***

^a Flax in Block A and Peas in Block B

^bU1A = Addition treatment pair 1; U1D = Disturbed treatment pair 1; U2A = Addition treatment pair 2; U2C = Control treatment pair 2

*Significant at $P < 0.10$; ** Significant at $P < 0.05$; ***Significant at $P < 0.01$ using a paired t-test

Note: Values indicate the means of 3 replicates

Table N.3 Cumulative greenhouse gas flux over a 118 day sampling period in 2004 for the upper slope area of FE2

	CO ₂ (mg CO ₂ -C m ⁻²)	CH ₄ (μg CH ₄ -C m ⁻²)	N ₂ O (μg N ₂ O-N m ⁻²)
Addition	1068b	-329	760
Control	1908a	-287	885
Disturbed	1020b	-378	1348

a-c Mean values followed by the same letter (within columns) are not significantly different based on an LSMeans comparison at $P < 0.01$

Note: Values indicate the means of 9 replicates

Appendix O

Column Experiment Part 1 (Addition Experiment) and Column Experiment Part 2

(Removal Experiment) Cumulative Flux Data




Figure O.1 Cumulative (a) carbon dioxide, (b) methane and (c) nitrous oxide flux for a 30-day sampling period from Column Experiment Part 1 (LSD groupings based $P < 0.10$).




Figure O.2 Cumulative (a) carbon dioxide, (b) methane and (c) nitrous oxide rates per mass of soil from Column Experiment Part 1 (LSD groupings based $P < 0.05$).






Figure O.3 Cumulative (a) carbon dioxide, (b) methane and (c) nitrous oxide flux results from a 36-day sampling period for Column Study Part 2 – Removal Experiment (LSD groupings based on a P<0.05).